

# Document made available under the Patent Cooperation Treaty (PCT)

International application number: PCT/US04/034693

International filing date: 15 October 2004 (15.10.2004)

Document type: Certified copy of priority document

Document details: Country/Office: US  
Number: 60/615,548  
Filing date: 01 October 2004 (01.10.2004)

Date of receipt at the International Bureau: 06 June 2005 (06.06.2005)

Remark: Priority document submitted or transmitted to the International Bureau in compliance with Rule 17.1(a) or (b)



World Intellectual Property Organization (WIPO) - Geneva, Switzerland  
Organisation Mondiale de la Propriété Intellectuelle (OMPI) - Genève, Suisse

110685



# THE UNITED STATES OF AMERICA

TO ALL TO WHOM THESE PRESENTS SHALL COME:

UNITED STATES DEPARTMENT OF COMMERCE

United States Patent and Trademark Office

May 26, 2005

THIS IS TO CERTIFY THAT ANNEXED HERETO IS A TRUE COPY FROM THE RECORDS OF THE UNITED STATES PATENT AND TRADEMARK OFFICE OF THOSE PAPERS OF THE BELOW IDENTIFIED PATENT APPLICATION THAT MET THE REQUIREMENTS TO BE GRANTED A FILING DATE.

APPLICATION NUMBER: 60/615,548

FILING DATE: October 01, 2004

RELATED PCT APPLICATION NUMBER: PCT/US04/34693



Certified by

*Don W. Dudas*

Under Secretary of Commerce  
for Intellectual Property  
and Director of the United States  
Patent and Trademark Office

**COVER SHEET FOR PROVISIONAL APPLICATION FOR PATENT**

Commissioner for Patents  
P.O. Box 1450  
Mail Stop Provisional Patent Application  
Alexandria, VA 22313-1450

Sir:

This is a request for filing a PROVISIONAL APPLICATION under 37 CFR 1.53(c).

17513 U.S. PTO  
60/615548  
100104

Docket Number		10271-143-888		Type a plus sign (+) inside this box →	+
INVENTOR(S) APPLICANT(S)					
LAST NAME	FIRST NAME	MIDDLE INITIAL	RESIDENCE (CITY AND EITHER STATE OR FOREIGN COUNTRY)		
Kinch	Michael	S.	Laytonsville, MD		
TITLE OF THE INVENTION (280 characters max)					
EPA2 VACCINES					
JONES DAY CORRESPONDENCE ADDRESS :20583					
ENCLOSED APPLICATION PARTS (check all that apply)					
<input checked="" type="checkbox"/> Specification	Number of Pages	120	<input type="checkbox"/> Applicant claims small entity status, see 37 CFR §1.27		
<input checked="" type="checkbox"/> Drawing(s)	Number of Sheets	30	<input type="checkbox"/> Other (specify)		
METHOD OF PAYMENT (check one)					
<input type="checkbox"/> A check or money order is enclosed to cover the Provisional filing fees.				ESTIMATED PROVISIONAL FILING FEE AMOUNT	
<input checked="" type="checkbox"/> The Commissioner is hereby authorized to charge the required filing fee to Deposit Account Number 50-3013.				<input checked="" type="checkbox"/> \$160 <input type="checkbox"/> \$80	

The invention was made by an agency of the United States Government or under a contract with an agency of the United States Government.  
☒ No. ☐ Yes, the name of the U.S. Government agency and the Government contract number are:

Respectfully submitted,

Signature

*Margaret B. Brivanlou*  
Margaret B. Brivanlou  
JONES DAY

REGISTRATION NO.  
(if appropriate)

40,922

Date

October 1, 2004

☐ Additional inventors are being named on separately numbered sheets attached hereto.

Total number of cover sheet pages.

1

**PROVISIONAL APPLICATION FILING ONLY**

## **EphA2 VACCINES**

---

### **1. FIELD OF THE INVENTION**

[0001] The present invention relates to methods and compositions designed for the treatment, management, or prevention of proliferative cell disease. The present invention further relates to methods and compositions for eliciting an immune response against hyperproliferative cells. The methods of the invention comprise the administration of an effective amount of an EphA2 vaccine, comprising, for example, EphA2 antigenic peptides or an EphA2 antigenic peptide expression vehicle. The invention also provides pharmaceutical compositions comprising one or more EphA2 antigenic peptides or peptide expression vehicles of the invention either alone or in combination with one or more other agents useful for therapy of proliferative disorders.

### **2. BACKGROUND OF THE INVENTION**

#### **2.1 HYPERPROLIFERATIVE DISEASES**

##### **2.1.1 Cancer**

[0002] A neoplasm, or tumor, is a neoplastic mass resulting from abnormal uncontrolled cell growth which can be benign or malignant. Benign tumors generally remain localized. Malignant tumors are collectively termed cancers. The term "malignant" generally means that the tumor can invade and destroy neighboring body structures and spread to distant sites to cause death (for review, *see* Robbins and Angell, 1976, Basic Pathology, 2d Ed., W.B. Saunders Co., Philadelphia, pp. 68-122). Cancer can arise in many sites of the body and behaves differently depending upon its origin. Cancerous cells destroy the part of the body in which they originate and then spread to other part(s) of the body where they start new growth and cause more destruction.

[0003] More than 1.2 million Americans develop cancer each year. Cancer is the second leading cause of death in the United States and, if current trends continue, cancer is expected to be the leading cause of death by the year 2010. Lung and prostate cancer are the top cancer killers for men in the United States. Lung and breast cancer are the top cancer killers for women in the United States. One in two men in the United States will be diagnosed with cancer at some time during his lifetime. One in three women in the United States will be diagnosed with cancer at some time during her lifetime.

[0004] A cure for cancer has yet to be found. Current treatment options, such as surgery, chemotherapy and radiation treatment, are often either ineffective or present serious side effects.

### 2.1.2 Metastasis

5 [0005] The most life-threatening forms of cancer often arise when a population of tumor cells gains the ability to colonize distant and foreign sites in the body. These metastatic cells survive by overriding restrictions that normally constrain cell colonization into dissimilar tissues. For example, typical mammary epithelial cells will generally not grow or survive if transplanted to the lung, yet lung metastases are a major cause of breast  
10 cancer morbidity and mortality. Recent evidence suggests that dissemination of metastatic cells through the body can occur long before clinical presentation of the primary tumor. These micrometastatic cells may remain dormant for many months or years following the detection and removal of the primary tumor. Thus, a better understanding of the mechanisms that allow for the growth and survival of metastatic cells in a foreign  
15 microenvironment is critical for the improvement of therapeutics designed to fight metastatic cancer and diagnostics for the early detection and localization of metastases.

### 2.1.3 Cancer Cell Signaling

[0006] Cancer is a disease of aberrant signal transduction. Aberrant cell signaling overrides anchorage-dependent constraints on cell growth and survival (Rhim *et al.*, 1997,  
20 *Crit. Rev. in Oncogenesis* 8:305; Patarca, 1996, *Crit. Rev. in Oncogenesis* 7:343; Malik *et al.*, 1996, *Biochimica et Biophysica Acta* 1287:73; Cance *et al.*, 1995, *Breast Cancer Res. Treat.* 35:105). Tyrosine kinase activity is induced by extracellular matrix (ECM) anchorage and indeed, the expression or function of tyrosine kinases is usually increased in malignant cells (Rhim *et al.*, 1997, *Critical Reviews in Oncogenesis* 8:305; Cance *et al.*,  
25 1995, *Breast Cancer Res. Treat.* 35:105, 1995; Hunter, 1997, *Cell* 88:333). Based on evidence that tyrosine kinase activity is necessary for malignant cell growth, tyrosine kinases have been targeted with new therapeutics (Levitzi *et al.*, 1995, *Science* 267:1782; Kondapaka *et al.*, 1996, *Mol. & Cell. Endocrinol.* 117:53; Fry *et al.*, 1995, *Curr. Opin. in BioTechnology* 6:662). Unfortunately, obstacles associated with specific targeting to tumor  
30 cells often limit the application of these drugs. In particular, tyrosine kinase activity is often vital for the function and survival of benign tissues (Levitzi *et al.*, 1995, *Science* 267:1782). To minimize collateral toxicity, it is critical to first identify and then target tyrosine kinases that are selectively overexpressed in tumor cells.

#### 2.1.4 Cancer Therapy

[0007] Barriers to the development of anti-metastasis agents have been the assay systems that are used to design and evaluate these drugs. Most conventional cancer therapies target rapidly growing cells. However, cancer cells do not necessarily grow more rapidly but instead survive and grow under conditions that are non-permissive to normal cells (Lawrence and Steeg, 1996, *World J. Urol.* 14:124-130). These fundamental differences between the behavior of normal and malignant cells provide opportunities for therapeutic targeting. The paradigm that micrometastatic tumors have already disseminated throughout the body emphasizes the need to evaluate potential chemotherapeutic drugs in the context of a foreign and three-dimensional microenvironment. Many standard cancer drug assays measure tumor cell growth or survival under typical cell culture conditions (*i.e.*, monolayer growth). However, cell behavior in two-dimensional assays often does not reliably predict tumor cell behavior *in vivo*.

[0008] Currently, cancer therapy may involve surgery, chemotherapy, hormonal therapy and/or radiation treatment to eradicate neoplastic cells in a patient (*see, e.g.*, Stockdale, 1998, "Principles of Cancer Patient Management," in *Scientific American: Medicine*, vol. 3, Rubenstein and Federman, eds., ch. 12, sect. IV). Recently, cancer therapy may also involve biological therapy or immunotherapy. All of these approaches can pose significant drawbacks for the patient. Surgery, for example, may be contraindicated due to the health of the patient or may be unacceptable to the patient. Additionally, surgery may not completely remove the neoplastic tissue. Radiation therapy is only effective when the neoplastic tissue exhibits a higher sensitivity to radiation than normal tissue, and radiation therapy can also often elicit serious side effects. Hormonal therapy is rarely given as a single agent and, although it can be effective, is often used to prevent or delay recurrence of cancer after other treatments have removed the majority of the cancer cells. Biological therapies/immunotherapies are limited in number and each therapy is generally effective for only a very specific type of cancer.

[0009] With respect to chemotherapy, there are a variety of chemotherapeutic agents available for treatment of cancer. A significant majority of cancer chemotherapeutics act by inhibiting DNA synthesis, either directly, or indirectly by inhibiting the biosynthesis of the deoxyribonucleotide triphosphate precursors, to prevent DNA replication and concomitant cell division (*see, e.g.*, Gilman *et al.*, 1990, Goodman and Gilman's: *The Pharmacological Basis of Therapeutics*, 8th Ed. (Pergamom Press, New York)). These agents, which include alkylating agents, such as nitrosourea, anti-metabolites, such as methotrexate and hydroxyurea, and other agents, such as etoposides, campathecins, bleomycin, doxorubicin,

daunorubicin, *etc.*, although not necessarily cell cycle specific, kill cells during S phase because of their effect on DNA replication. Other agents, specifically colchicine and the vinca alkaloids, such as vinblastine and vincristine, interfere with microtubule assembly resulting in mitotic arrest. Chemotherapy protocols generally involve administration of a combination of chemotherapeutic agents to increase the efficacy of treatment.

[0010] Despite the availability of a variety of chemotherapeutic agents, chemotherapy has many drawbacks (*see, e.g.*, Stockdale, 1998, "Principles Of Cancer Patient Management" in Scientific American Medicine, vol. 3, Rubenstein and Federman, eds., ch. 12, sect. X). Almost all chemotherapeutic agents are toxic, and chemotherapy causes significant, and often dangerous, side effects, including severe nausea, bone marrow depression, immunosuppression, *etc.* Additionally, even with administration of combinations of chemotherapeutic agents, many tumor cells are resistant or develop resistance to the chemotherapeutic agents. In fact, those cells resistant to the particular chemotherapeutic agents used in the treatment protocol often prove to be resistant to other drugs, even those agents that act by mechanisms different from the mechanisms of action of the drugs used in the specific treatment; this phenomenon is termed pleiotropic drug or multidrug resistance. Thus, because of drug resistance, many cancers prove refractory to standard chemotherapeutic treatment protocols.

[0011] There is a significant need for alternative cancer treatments, particularly for treatment of cancer that has proved refractory to standard cancer treatments, such as surgery, radiation therapy, chemotherapy, and hormonal therapy. Further, it is uncommon for cancer to be treated by only one method. Thus, there is a need for development of new therapeutic agents for the treatment of cancer and new, more effective, therapy combinations for the treatment of cancer.

## **2.1.5 OTHER HYPERPROLIFERATIVE DISORDERS**

### **2.1.5.1 Asthma**

[0012] Asthma is a disorder characterized by intermittent airway obstruction. In western countries, it affects 15% of the pediatric population and 7.5% of the adult population (Strachan *et al.*, 1994, *Arch. Dis. Child* 70:174-178). Most asthma in children and young adults is initiated by IgE mediated allergy (atopy) to inhaled allergens such as house dust mite and cat dander allergens. However, not all asthmatics are atopic, and most atopic individuals do not have asthma. Thus, factors in addition to atopy are necessary to induce the disorder (Fraser *et al.*, eds., 1994, *Synopsis of Diseases of the Chest*: 635-53 (WB Saunders Company, Philadelphia); Djukanovic *et al.*, 1990, *Am. Rev. Respir. Dis.*

142:434-457). Asthma is strongly familial, and is due to the interaction between genetic and environmental factors. The genetic factors are thought to be variants of normal genes ("polymorphisms") which alter their function to predispose to asthma.

[0013] Asthma may be identified by recurrent wheeze and intermittent air flow limitation. An asthmatic tendency may be quantified by the measurement of bronchial hyper-responsiveness in which an individual's dose-response curve to a broncho-constrictor such as histamine or methacholine is constructed. The curve is commonly summarized by the dose which results in a 20% fall in air flow (PD20) or the slope of the curve between the initial air flow measurement and the last dose given (slope).

[0014] In the atopic response, IgE is produced by B-cells in response to allergen stimulation. These antibodies coat mast cells by binding to the high affinity receptor for IgE and initiate a series of cellular events leading to the destabilization of the cell membrane and release of inflammatory mediators. This results in mucosal inflammation, wheezing, coughing, sneezing and nasal blockage.

[0015] Atopy can be diagnosed by (i) a positive skin prick test in response to a common allergen; (ii) detecting the presence of specific serum IgE for allergen; or (iii) by detecting elevation of total serum IgE.

#### **2.5.1.2 COPD**

[0016] Chronic obstructive pulmonary disease (COPD) is an umbrella term frequently used to describe two conditions of fixed airways disorders, chronic bronchitis and emphysema. Chronic bronchitis and emphysema are most commonly caused by smoking; approximately 90% of patients with COPD are or were smokers. Although approximately 50% of smokers develop chronic bronchitis, only 15% of smokers develop disabling airflow obstruction. Certain animals, particularly horses, suffer from COPD as well.

[0017] The airflow obstruction associated with COPD is progressive, may be accompanied by airway hyperactivity, and may be partially reversible. Non-specific airway hyper-responsiveness may also play a role in the development of COPD and may be predictive of an accelerated rate of decline in lung function.

[0018] COPD is a significant cause of death and disability. It is currently the fourth leading cause of death in the United States and Europe. Treatment guidelines advocate early detection and implementation of smoking cessation programs to help reduce morbidity and mortality due to the disorder. However, early detection and diagnosis has been difficult for a number of reasons. COPD takes years to develop and acute episodes of bronchitis often are not recognized by the general practitioner as early signs of COPD. Many patients



exhibit features of more than one disorder (e.g., chronic bronchitis or asthmatic bronchitis) making precise diagnosis a challenge, particularly early in the etiology of the disorder. Also, many patients do not seek medical help until they are experiencing more severe symptoms associated with reduced lung function, such as dyspnea, persistent cough, and sputum production. As a consequence, the vast majority of patients are not diagnosed or treated until they are in a more advanced stage of the disorder.

### 2.1.5.3 Mucin

[0019] Mucins are a family of glycoproteins secreted by the epithelial cells including those at the respiratory, gastrointestinal and female reproductive tracts. Mucins are responsible for the viscoelastic properties of mucus (Thornton *et al.*, 1997, *J. Biol. Chem.* 272:9561-9566). Nine mucin genes are known to be expressed in man: MUC 1, MUC 2, MUC 3, MUC 4, MUC 5AC, MUC 5B, MUC 6, MUC 7 and MUC 8 (Bobek *et al.*, 1993, *J. Biol. Chem.* 268:20563-9; Dusseyn *et al.*, 1997, *J. Biol. Chem.* 272:3168-78; Gendler *et al.*, 1991, *Am. Rev. Resp. Dis.* 144:S42-S47; Gum *et al.*, 1989, *J. Biol. Chem.* 264:6480-6487; Gum *et al.*, 1990, *Biochem. Biophys. Res. Comm.* 171:407-415; Lesuffleur *et al.*, 1995, *J. Biol. Chem.* 270:13665-13673; Meerzaman *et al.*, 1994, *J. Biol. Chem.* 269:12932-12939; Porchet *et al.*, 1991, *Biochem. Biophys. Res. Comm.* 175:414-422; Shankar *et al.*, 1994, *Biochem. J.* 300:295-298; Toribara *et al.*, 1997, *J. Biol. Chem.* 272:16398-403). Many airway disorders such chronic bronchitis, chronic obstructive pulmonary disease, bronchiectasis, asthma, cystic fibrosis and bacterial infections are characterized by mucin overproduction (Prescott *et al.*, *Eur. Respir. J.*, 1995, 8:1333-1338; Kim *et al.*, *Eur. Respir. J.*, 1997, 10:1438; Steiger *et al.*, 1995, *Am. J. Respir. Cell Mol. Biol.*, 12:307-314). Mucociliary impairment caused by mucin hypersecretion leads to airway mucus plugging which promotes chronic infection, airflow obstruction and sometimes death. For example, COPD, a disorder characterized by slowly progressive and irreversible airflow limitation, is a major cause of death in developed countries. The respiratory degradation consists mainly of decreased luminal diameters due to airway wall thickening and increased mucus caused by goblet cell hyperplasia and hypersecretion. Epidermal growth factor (EGF) is known to upregulate epithelial cell proliferation, and mucin production/secretion (Takeyama *et al.*, 1999, *Proc. Natl. Acad. Sci. USA* 96:3081-6; Burgel *et al.*, 2001, *J. Immunol.* 167:5948-54). EGF also causes mucin-secreting cells, such as goblet cells, to proliferate and increase mucin production in airway epithelia (Lee *et al.*, 2000, *Am. J. Physiol. Lung Cell. Mol. Physiol.* 278:185-92; Takeyama *et al.*, 2001, *Am. J. Respir. Crit. Care. Med.* 163:511-6; Burgel *et al.*, 2000, *J. Allergy Clin. Immunol.* 106:705-12). Historically, mucus hypersecretion has been treated in two ways: physical methods to

increase clearance and mucolytic agents. Neither approach has yielded significant benefit to the patient or reduced mucus obstruction. Therefore, it would be desirable to have methods for reducing mucin production and treating the disorders associated with mucin hypersecretion.

5                                   **2.1.5.4 Restenosis**

[0020]           Vascular interventions, including angioplasty, stenting, atherectomy and grafting are often complicated by undesirable effects. Exposure to a medical device which is implanted or inserted into the body of a patient can cause the body tissue to exhibit adverse physiological reactions. For instance, the insertion or implantation of certain  
10 catheters or stents can lead to the formation of emboli or clots in blood vessels. Other adverse reactions to vascular intervention include endothelial cell proliferation which can lead to hyperplasia, restenosis, *i.e.* the re-occlusion of the artery, occlusion of blood vessels, platelet aggregation, and calcification. Treatment of restenosis often involves a second angioplasty or bypass surgery. In particular, restenosis may be due to endothelial cell injury  
15 caused by the vascular intervention in treating a restenosis.

[0021]           Angioplasty involves insertion of a balloon catheter into an artery at the site of a partially obstructive atherosclerotic lesion. Inflation of the balloon is intended to rupture the intima and dilate the obstruction. About 20 to 30% of obstructions reocclude in just a few days or weeks (Elтчaninoff *et al.*, 1998, *J. Am Coll. Cardiol.* 32: 980-984). Use  
20 of stents reduces the re-occlusion rate, however a significant percentage continues to result in restenosis. The rate of restenosis after angioplasty is dependent upon a number of factors including the length of the plaque. Stenosis rates vary from 10% to 35% depending the risk factors present. Further, repeat angiography one year later reveals an apparently normal lumen in only about 30% of vessels having undergone the procedure.

25 [0022]           Restenosis is caused by an accumulation of extracellular matrix containing collagen and proteoglycans in association with smooth muscle cells which is found in both the atheroma and the arterial hyperplastic lesion after balloon injury or clinical angioplasty. Some of the delay in luminal narrowing with respect to smooth muscle cell proliferation may result from the continuing elaboration of matrix materials by neointimal smooth  
30 muscle cells. Various mediators may alter matrix synthesis by smooth muscle cells *in vivo*.

**2.1.5.5 Neointimal Hyperplasia**

[0023]           Neointimal hyperplasia is the pathological process that underlies graft atherosclerosis, stenosis, and the majority of vascular graft occlusion. Neointimal hyperplasia is commonly seen after various forms of vascular injury and a major component

of the vein graft's response to harvest and surgical implantation into high-pressure arterial circulation.

[0024] Smooth muscle cells in the middle layer (*i.e.* media layer) of the vessel wall become activated, divide, proliferate and migrate into the inner layer (*i.e.* intima layer). The resulting abnormal neointimal cells express pro-inflammatory molecules, including cytokines, chemokines and adhesion molecules that further trigger a cascade of events that lead to occlusive neointimal disease and eventually graft failure.

[0025] The proliferation of smooth muscle cells is a critical event in the neointimal hyperplastic response. Using a variety of approaches, studies have clearly demonstrated that blockade of smooth muscle cell proliferation resulted in preservation of normal vessel phenotype and function, causing the reduction of neointimal hyperplasia and graft failure.

[0026] Existing treatments for the indications discussed above is inadequate; thus, there exists a need for improved treatments for the above indications.

## 2.2 EphA2

[0027] EphA2 is a 130 kDa receptor tyrosine kinase that is expressed in adult epithelia, where it is found at low levels and is enriched within sites of cell-cell adhesion (Zantek *et al.*, 1999, *Cell Growth & Differentiation* 10:629; Lindberg *et al.*, 1990, *Molecular & Cellular Biology* 10:6316). This subcellular localization is important because EphA2 binds ligands (known as Ephrins A1 to A5) that are anchored to the cell membrane (Eph Nomenclature Committee, 1997, *Cell* 90:403; Gale *et al.*, 1997, *Cell & Tissue Research* 290: 227). The primary consequence of ligand binding is EphA2 autophosphorylation (Lindberg *et al.*, 1990, *supra*). However, unlike other receptor tyrosine kinases, EphA2 retains enzymatic activity in the absence of ligand binding or phosphotyrosine content (Zantek *et al.*, 1999, *supra*). EphA2 is upregulated on a large number hyperproliferating cells, including aggressive carcinoma cells.

## 3. SUMMARY OF THE INVENTION

[0028] EphA2 is overexpressed and functionally altered in a large number of malignant carcinomas. EphA2 is an oncoprotein and is sufficient to confer metastatic potential to cancer cells. EphA2 is also associated with other hyperproliferating cells and is implicated in diseases caused by cell hyperproliferation. The present invention stems from the inventors' discovery that administration of an expression vehicle for an EphA2 antigenic peptide to a subject provides beneficial therapeutic and prophylactic benefits against hyperproliferative disorders involving EphA2 overexpressing cells. Without being

bound by any mechanism or theory, it is believed that the therapeutic and prophylactic benefit is the result of an immune response elicited against the EphA2 antigenic peptide.

**[0029]** The present invention thus provides EphA2 vaccines and methods for their use. The EphA2 vaccines of the present invention can elicit or mediate a cellular immune response, a humoral immune response, or both. Where the immune response is a cellular immune response, it can be a Tc, Th1 or a Th2 immune response. In a preferred embodiment, the immune response is a Th2 cellular immune response.

**[0030]** In a preferred embodiment, an EphA2 vaccine of the invention comprises or encodes one or more epitopes on EphA2 that is selectively exposed or increased on cancer cells relative to not non-cancer cells. In one embodiment, the cancer is of an epithelial cell origin. In other embodiments, the cancer is a cancer of the skin, lung, colon, prostate, breast, ovary, esophageal, bladder, or pancreas or is a renal cell carcinoma or a melanoma. In another embodiment, the cancer is of a T cell origin. In yet other embodiments, the cancer is a leukemia or a lymphoma.

**[0031]** In a preferred embodiment, the methods and compositions of the invention are used to prevent, treat or manage EphA2-expressing tumor metastases. In a preferred embodiment, the EphA2-expressing cells against which an immune response is sought ("target cells") overexpress EphA2. In a preferred embodiment, some EphA2 on the target cells is not bound to ligand, either as a result of decreased cell-cell contacts, altered subcellular localization, or increases in amount of EphA2 relative to ligand.

**[0032]** Thus, the present invention provides methods of eliciting an immune response against an EphA2-expressing cell, said method comprising administering to an individual an EphA2 vaccine in an amount effective to elicit an immune response against an EphA2-expressing cell.

**[0033]** The present invention further provides a method of treating or preventing a hyperproliferative disorder of EphA2-expressing cells, said method comprising administering to an individual an EphA2 vaccine in an amount effective to treat or prevent the hyperproliferative disorder.

**[0034]** The present invention yet further provides EphA2 vaccines useful for eliciting an immune response against an EphA2-expressing cell and/or for treating or preventing a hyperproliferative disorder of EphA2-expressing cells.

**[0035]** The EphA2 vaccines may comprise an EphA2 antigenic peptide, an EphA2 antigenic peptide expression vehicle, an antigen presenting cell sensitized with an EphA2 antigenic peptide, or an anti-idiotypic antibody or antigen-binding fragment thereof which immunospecifically binds to an idiotype of an anti-EphA2 antibody.

[0036] In embodiments where an EphA2 vaccine comprises an EphA2 antigenic peptide, the vaccine may further comprise an adjuvant, or a heat shock protein bound to the EphA2 antigenic peptide.

[0037] In certain embodiments, the EphA2 antigenic peptide comprises a protein transduction domain, for example the protein transduction domain is the Antennapedia or the HIV tat protein transduction domain.

[0038] In certain embodiments in which an EphA2 vaccine comprises an EphA2 antigenic peptide expression vehicle, the expression vehicle can be a nucleic acid, preferably DNA, encoding said EphA2 antigenic polypeptide operably linked to a promoter.

The DNA can be conjugated to a carrier, including but not limited to an asialoglycoprotein carrier, a transferrin carrier, or a polymeric IgA carriers.

[0039] In other embodiments, the expression vehicle is an infectious agent comprising a nucleic acid, said nucleic acid comprising a nucleotide sequence encoding said EphA2 antigenic polypeptide operably linked to a promoter. Preferably, the sequence encoding said EphA2 antigenic polypeptide is codon-optimized for expression in said infectious agent and/or the infectious agent is coated with a reagent that targets the infectious agent to EphA2-expressing cells (such as an EphA2 antibody) or to antigen-presenting cells.

[0040] A preferred infectious agent for use as an EphA2 antigenic peptide expression vehicle in accordance with the methods and compositions of the invention is a bacterium. Preferred bacteria for administration to human subjects are attenuated, for example are attenuated in their tissue tropism or ability to spread from cell to cell. Preferably, the nucleic acid encoding the EphA2 antigenic peptide comprises a nucleotide sequence encoding a secretory signal, e.g., the SecA secretory signal, operatively linked to the sequence encoding the EphA2 antigenic polypeptide. A preferred strain of bacteria for use in the methods and compositions of the invention is *Pseudomonas aeruginosa*. In certain specific embodiments, the bacteria is not *Listeria*, and more preferably is not *Listeria monocytogenes*.

[0041] Another preferred infectious agent for use as an EphA2 antigenic peptide expression vehicle in accordance with the methods and compositions of the invention is a virus, for example a retrovirus, including but not limited to lentivirus, an adenovirus, an adeno-associated virus, or a herpes simplex virus. Preferred viruses for administration to human subjects are attenuated viruses.

[0042] As an alternative to an infectious agent or nucleic acid, an EphA2 antigenic peptide expression vehicle can be a mammalian cell comprising a recombinant nucleic acid,

said nucleic acid comprising a nucleotide sequence encoding said EphA2 antigenic polypeptide. Preferably, the mammalian cell is a human cell. Mammalian cells for use in the methods and compositions of the invention may be encapsulated within a membrane, for example a THERACYTE membrane, and/or administered by means of implantation.

5     **[0043]**       Compositions of the present invention useful as EphA2 vaccines also include anti-idiotypes of anti-EphA2 antibodies. In certain specific embodiments, the EphA2 vaccines comprise anti-idiotypes of the anti-EphA2 monoclonal antibodies secreted by the hybridoma clones deposited in the ATCC as PTA-4572, PTA-4573, and PTA-4574.

10     **[0044]**       With respect to EphA2 vaccines comprising sensitized antigen presenting cells, such as macrophages and dendritic cells, in certain embodiments, the antigen presenting cells are sensitized prior to their administration. For example, the antigen presenting cells may be sensitized by a method comprising: (a) contacting the cells with a composition comprising one or more EphA2 antigenic peptides, and optionally comprising one or more heat shock proteins such as hsp70, gp96, or hsp90, in an amount effective to  
15     sensitize the cells. In a preferred embodiment, the antigen presenting cells are autologous to the individual to whom they are administered; however, the cells need not be autologous.

20     **[0045]**       The compositions and methods of the present invention are useful in the treatment of hyperproliferative diseases. In certain embodiments, the hyperproliferative disease is cancer. In certain embodiments, the cancer is of an epithelial cell origin and/or involves cells that overexpress EphA2 relative to non-cancer cells having the tissue type of said cancer cells. In specific embodiments, the cancer is a cancer of the skin, lung, colon, breast, ovarian, esophageal, prostate, bladder or pancreas or is a renal cell carcinoma or melanoma. In yet other embodiments, the cancer is of a T cell origin. In specific  
25     embodiments, the cancer is a leukemia or a lymphoma. In yet other embodiments, the hyperproliferative disorder is non-neoplastic. In specific embodiments, the non-neoplastic hyperproliferative disorder is an epithelial cell disorder. Exemplary non-neoplastic hyperproliferative disorders are asthma, chronic pulmonary obstructive disease, lung fibrosis, bronchial hyper responsiveness, psoriasis, and seborrheic dermatitis.

30     **[0046]**       The EphA2 antigenic polypeptide for use in accordance with the methods and compositions of the present invention may comprise full length EphA2 or an antigenic fragment or derivative thereof. In certain embodiments, the EphA2 antigenic polypeptide comprises the extracellular domain of EphA2 or the intracellular domain of EphA2. In certain embodiments the EphA2 antigenic polypeptide is a chimeric polypeptide comprising at least an antigenic portion of EphA2 and a second polypeptide.

**[0047]** A vaccine of the invention may have one or a plurality of EphA2 antigenic polypeptides, a plurality of EphA2 antigenic polypeptide expression vehicles, or antigen presenting cells sensitized with a plurality of EphA2 antigenic polypeptides. A vaccine of the invention may also have one or more classes of immune response-inducing or -

mediating reagents, for example both an EphA2 antigenic polypeptide and an EphA2 antigenic polypeptide expression vehicle, both an EphA2 antigenic polypeptide and an antigen-presenting cell sensitized with an EphA2 antigenic polypeptide, or both an EphA2 antigenic polypeptide expression vehicle and an antigen-presenting cell sensitized with an EphA2 antigenic polypeptide.

**[0048]** The methods of the present invention encompass combination therapy with an EphA2 vaccine and one or more additional therapeutics, for example an additional anti-cancer therapy. In certain embodiments, the additional anti-cancer therapy is an agonistic EphA2 antibody. In other embodiments, the additional anti-cancer therapy is chemotherapy, biological therapy, immunotherapy, radiation therapy, hormonal therapy, or surgery.

**[0049]** The vaccines of the invention can be administered by mucosal, intranasal, parenteral, intramuscular, or intraperitoneal routes.

**[0050]** In other embodiments, the EphA2 vaccines of the invention are used to treat, prevent and/or manage a non-cancer disease or disorder associated with cell

hyperproliferation, such as but not limited to asthma, chronic obstructive pulmonary disease, restenosis (smooth muscle and/or endothelial), psoriasis, etc. In preferred embodiments, the hyperproliferative cells are epithelial. In preferred embodiments, the hyperproliferative cells overexpress EphA2. In a preferred embodiment, some EphA2 is not bound to ligand, either as a result of decreased cell-cell contacts, altered subcellular localization, or increases in amount of EphA2 relative to EphA2-ligand.

**[0051]** In yet other aspects of the invention, the EphA2 vaccines are used to treat, prevent and/or manage a disease or disorder associated involving aberrant angiogenesis. The EphA2 vaccines are used to elicit an immune response against EphA2 expressed on neovasculature. Thus, the present invention provides methods comprising administering to

the subject a composition comprising an EphA2 vaccine in an amount effective to treat disease involving aberrant angiogenesis. Examples of such diseases include but are not limited to macular degeneration, diabetic retinopathy, retinopathy of prematurity, vascular restenosis, infantile hemangioma, verruca vulgaris, psoriasis, Kaposi's sarcoma, neurofibromatosis, recessive dystrophic epidermolysis bullosa, rheumatoid arthritis, ankylosing spondylitis, systemic lupus, psoriatic arthropathy, Reiter's syndrome, and

Sjogren's syndrome, endometriosis, preeclampsia, atherosclerosis and coronary artery disease.

[0052] The methods and compositions of the invention are useful not only in untreated patients but are also useful in the treatment of patients partially or completely refractory to current standard and experimental cancer therapies, including but not limited to chemotherapies, hormonal therapies, biological therapies, radiation therapies, and/or surgery as well as to improve the efficacy of such treatments. In particular, EphA2 expression has been implicated in increasing levels of the cytokine IL-6, which has been associated with the development of cancer cell resistance to different treatment regimens, such as chemotherapy and hormonal therapy. In addition, EphA2 overexpression can override the need for estrogen receptor activity thus contributing to tamoxifen resistance in breast cancer cells. Accordingly, in a preferred embodiment, the invention provides therapeutic and prophylactic methods for the treatment or prevention of cancer that has been shown to be or may be refractory or non-responsive to therapies other than those comprising administration of EphA2 antibodies of the invention. In a specific embodiment, one or more EphA2 vaccines of the invention are administered to a patient refractory or non-responsive to a non-EphA2-based treatment, particularly tamoxifen treatment or a treatment in which resistance is associated with increased IL-6 levels, to render the patient non-refractory or responsive. The treatment to which the patient had previously been refractory or non-responsive can then be administered with therapeutic effect.

[0053] In another embodiment, kits comprising the vaccines or vaccine components of the invention are provided.

### **3.1 DEFINITIONS**

[0054] As used herein, the term "EphA2 vaccine" can be any reagent that elicits or mediates an immune response against EphA2 on hyperproliferative cells. In certain embodiments, an EphA2 vaccine is an EphA2 antigenic peptide of the invention, an expression vehicle (e.g., a naked nucleic acid or a viral or bacterial vector or a cell) for an EphA2 antigenic peptide, or T cells or antigen presenting cells (e.g., dendritic cells or macrophages) that have been primed with the EphA2 antigenic peptide of the invention.

[0055] As used herein, the term "EphA2 antigenic peptide" or "EphA2 antigenic polypeptide" refers to an EphA2 polypeptide, preferably of SEQ ID NO:2, or a fragment or derivative thereof comprising one or more B cell epitopes or T cell epitopes of EphA2. In certain embodiments, the EphA2 antigenic peptides are not one or more of the following peptides: TLADFDPRV (SEQ ID NO:3); VLLLVLAGV (SEQ ID NO:4); VLAGVGFFI (SEQ ID NO:5); IMNDMPIYM (SEQ ID NO:6); SLLGLKDQV (SEQ ID NO:7);



WLVPIGQCL (SEQ ID NO:8); LLWGCALAA (SEQ ID NO:9); GLTRTSVTV (SEQ ID NO:10); NLYYAESDL (SEQ ID NO:11); KLNVEERSV (SEQ ID NO:12); IMGQFSHHN (SEQ ID NO:13); YSVCNVMSG (SEQ ID NO:14); MQNIMNDMP (SEQ ID NO:15); EAGIMGQFSHHNIIR (SEQ ID NO:16); PIYMYSVCNVMSG (SEQ ID NO:17);

- 5 DLMQNIMNDMPIYMYS (SEQ ID NO:18). In certain specific embodiments, the EphA2 antigenic peptide is not any of SEQ ID NO:3-12, is not SEQ ID NO:13-15, and/or is not SEQ ID NO:16-18. In yet another specific embodiment, the EphA2 antigenic peptide is not SEQ ID NO:3-18.

- [0056]** The term “derivative” as used herein refers to a polypeptide that comprises  
10 an amino acid sequence of an EphA2 polypeptide or a fragment of an EphA2 polypeptide that has been altered by the introduction of amino acid residue substitutions, deletions or additions (*i.e.*, mutations). The term “derivative” as used herein also refers to an EphA2 polypeptide or a fragment of an EphA2 polypeptide which has been modified, *i.e.*, by the covalent attachment of any type of molecule to the polypeptide. For example, but not by  
15 way of limitation, an EphA2 polypeptide or a fragment of an EphA2 polypeptide may be modified, *e.g.*, by glycosylation, acetylation, pegylation, phosphorylation, amidation, derivatization by known protecting/blocking groups, proteolytic cleavage, linkage to a cellular ligand or other protein, *etc.* A derivative of an EphA2 polypeptide or a fragment of an EphA2 polypeptide may be modified by chemical modifications using techniques known  
20 to those of skill in the art, including, but not limited to, specific chemical cleavage, acetylation, formylation, metabolic synthesis of tunicamycin, *etc.* Further, a derivative of an EphA2 polypeptide or a fragment of an EphA2 polypeptide may contain one or more non-classical amino acids. In one embodiment, a polypeptide derivative possesses a similar or identical function as an EphA2 polypeptide or a fragment of an EphA2 polypeptide  
25 described herein. In another embodiment, a derivative of EphA2 polypeptide or a fragment of an EphA2 polypeptide has an altered activity when compared to an unaltered polypeptide. For example, a derivative of an EphA2 polypeptide or fragment thereof can differ in phosphorylation relative to an EphA2 polypeptide or fragment thereof.

- [0057]** The term “B cell epitope” as used herein refers to a portion of an EphA2  
30 polypeptide having antigenic or immunogenic activity in an animal, preferably in a mammal, and most preferably in a mouse or a human. An epitope having immunogenic activity is a portion of an EphA2 polypeptide that elicits an antibody response in an animal. An epitope having antigenic activity is a portion of an EphA2 polypeptide to which an antibody immunospecifically binds as determined by any method well known in the art, for  
35 example, by immunoassays. Antigenic epitopes need not necessarily be immunogenic.

**[0058]** The term “T cell epitope” as used herein refers to at least a portion of an EphA2 polypeptide, preferably an EphA2 polypeptide of SEQ ID NO:2, that is recognized by a T cell receptor. The term “T cell epitope” encompasses helper T cell (Th) epitopes and cytotoxic T cell (Tc) epitopes. The term “helper T cell epitopes” encompasses Th1 and Th2 epitopes.

**[0059]** The “fragments” described herein include an EphA2 antigenic peptide or polypeptide comprising an amino acid sequence of at least 5 contiguous amino acid residues, at least 10 contiguous amino acid residues, at least 15 contiguous amino acid residues, at least 20 contiguous amino acid residues, at least 25 contiguous amino acid residues, at least 40 contiguous amino acid residues, at least 50 contiguous amino acid residues, at least 60 contiguous amino acid residues, at least 70 contiguous amino acid residues, at least 80 contiguous amino acid residues, at least 90 contiguous amino acid residues, at least 100 contiguous amino acid residues, at least 125 contiguous amino acid residues, at least 150 contiguous amino acid residues, at least 175 contiguous amino acid residues, at least 200 contiguous amino acid residues, or at least 250 contiguous amino acid residues of the amino acid sequence of an EphA2 polypeptide.

**[0060]** As used herein, the term “in combination” refers to the use of more than one prophylactic and/or therapeutic agents. The use of the term “in combination” does not restrict the order in which prophylactic and/or therapeutic agents are administered to a subject with a hyperproliferative cell disorder, especially cancer. A first prophylactic or therapeutic agent can be administered prior to (e.g., 1 minute, 5 minutes, 15 minutes, 30 minutes, 45 minutes, 1 hour, 2 hours, 4 hours, 6 hours, 12 hours, 24 hours, 48 hours, 72 hours, 96 hours, 1 week, 2 weeks, 3 weeks, 4 weeks, 5 weeks, 6 weeks, 8 weeks, or 12 weeks before), concomitantly with, or subsequent to (e.g., 1 minute, 5 minutes, 15 minutes, 30 minutes, 45 minutes, 1 hour, 2 hours, 4 hours, 6 hours, 12 hours, 24 hours, 48 hours, 72 hours, 96 hours, 1 week, 2 weeks, 3 weeks, 4 weeks, 5 weeks, 6 weeks, 8 weeks, or 12 weeks after) the administration of a second prophylactic or therapeutic agent to a subject which had, has, or is susceptible to a hyperproliferative cell disorder, especially cancer. The prophylactic or therapeutic agents are administered to a subject in a sequence and within a time interval such that the agent of the invention can act together with the other agent to provide an increased benefit than if they were administered otherwise. Any additional prophylactic or therapeutic agent can be administered in any order with the other additional prophylactic or therapeutic agents.

**[0061]** As used herein, the phrase “low tolerance” refers to a state in which the patient suffers from side effects from treatment so that the patient does not benefit from

and/or will not continue therapy because of the adverse effects and/or the harm from the side effects outweighs the benefit of the treatment.

**[0062]** As used herein, the terms “manage,” “managing” and “management” refer to the beneficial effects that a subject derives from administration of a prophylactic or therapeutic agent, which does not result in a cure of the disease. In certain embodiments, a subject is administered one or more prophylactic or therapeutic agents to “manage” a disease so as to prevent the progression or worsening of the disease.

**[0063]** As used herein, the phrase “non-responsive/refractory” is used to describe patients treated with one or more currently available therapies (e.g., cancer therapies) such as chemotherapy, radiation therapy, surgery, hormonal therapy and/or biological therapy/immunotherapy, particularly a standard therapeutic regimen for the particular cancer, wherein the therapy is not clinically adequate to treat the patients such that these patients need additional effective therapy, e.g., remain unsusceptible to therapy. The phrase can also describe patients who respond to therapy yet suffer from side effects, relapse, develop resistance, etc. In various embodiments, “non-responsive/refractory” means that at least some significant portion of the cancer cells are not killed or their cell division arrested. The determination of whether the cancer cells are “non-responsive/refractory” can be made either *in vivo* or *in vitro* by any method known in the art for assaying the effectiveness of treatment on cancer cells, using the art-accepted meanings of “refractory” in such a context. In various embodiments, a cancer is “non-responsive/refractory” where the number of cancer cells has not been significantly reduced, or has increased during the treatment.

**[0064]** As used herein, the term “potentiate” refers to an improvement in the efficacy of a therapeutic agent at its common or approved dose.

**[0065]** As used herein, the terms “prevent,” “preventing” and “prevention” refer to the prevention of the onset, recurrence, or spread of a disease in a subject resulting from the administration of a prophylactic or therapeutic agent.

**[0066]** As used herein, the term “prophylactic agent” refers to any agent that can be used in the prevention of the onset, recurrence or spread of a disease or disorder associated with EphA2 overexpression and/or cell hyperproliferative disease, particularly cancer. In certain embodiments, the term “prophylactic agent” refers to an EphA2 vaccine of the invention, such as a composition comprising an EphA2 antigenic peptide, an EphA2 antigenic peptide expression vehicle, or an antigen presenting cell sensitized with an EphA2 antigenic peptide. In certain other embodiments, the term “prophylactic agent” refers to cancer chemotherapeutics, radiation therapy, hormonal therapy, biological therapy (e.g.,

immunotherapy). In other embodiments, more than one prophylactic agent may be administered in combination.

**[0067]** As used herein, a “prophylactically effective amount” refers to that amount of the prophylactic agent sufficient to result in the prevention of the onset, recurrence or spread of cell hyperproliferative disease, preferably, cancer. A prophylactically effective amount may refer to the amount of prophylactic agent sufficient to prevent the onset, recurrence or spread of hyperproliferative disease, particularly cancer, including but not limited to those predisposed to hyperproliferative disease, for example, those genetically predisposed to cancer or previously exposed to carcinogens. A prophylactically effective amount may also refer to the amount of the prophylactic agent that provides a prophylactic benefit in the prevention of hyperproliferative disease. Further, a prophylactically effective amount with respect to a prophylactic agent of the invention means that amount of prophylactic agent alone, or in combination with other agents, that provides a prophylactic benefit in the prevention of hyperproliferative disease. Used in connection with an amount of an EphA2 vaccine of the invention, the term can encompass an amount that improves overall prophylaxis or enhances the prophylactic efficacy of or synergies with another prophylactic agent.

**[0068]** As used herein, a “protocol” includes dosing schedules and dosing regimens.

**[0069]** As used herein, the phrase “side effects” encompasses unwanted and adverse effects of a prophylactic or therapeutic agent. Adverse effects are always unwanted, but unwanted effects are not necessarily adverse. An adverse effect from a prophylactic or therapeutic agent might be harmful or uncomfortable or risky. Side effects from chemotherapy include, but are not limited to, gastrointestinal toxicity such as, but not limited to, early and late-forming diarrhea and flatulence, nausea, vomiting, anorexia, leukopenia, anemia, neutropenia, asthenia, abdominal cramping, fever, pain, loss of body weight, dehydration, alopecia, dyspnea, insomnia, dizziness, mucositis, xerostomia, and kidney failure, as well as constipation, nerve and muscle effects, temporary or permanent damage to kidneys and bladder, flu-like symptoms, fluid retention, and temporary or permanent infertility. Side effects from radiation therapy include but are not limited to fatigue, dry mouth, and loss of appetite. Side effects from biological therapies/immunotherapies include but are not limited to rashes or swellings at the site of administration, flu-like symptoms such as fever, chills and fatigue, digestive tract problems and allergic reactions. Side effects from hormonal therapies include but are not limited to nausea, fertility problems, depression, loss of appetite, eye problems, headache, and weight fluctuation. Additional undesired effects typically experienced by patients are numerous

and known in the art. Many are described in the *Physicians' Desk Reference* (56<sup>th</sup> ed., 2002).

**[0070]** As used herein, the terms “subject” and “patient” are used interchangeably.

As used herein, a subject is preferably a mammal such as a non-primate (e.g., cows, pigs, horses, cats, dogs, rats *etc.*) and a primate (e.g., monkey and human), most preferably a human.

**[0071]** As used herein, the terms “treat,” “treating” and “treatment” refer to the eradication, reduction or amelioration of symptoms of a disease or disorder, particularly, the eradication, removal, modification, or control of primary, regional, or metastatic cancer tissue that results from the administration of one or more therapeutic agents. In certain embodiments, such terms refer to the minimizing or delaying the spread of cancer resulting from the administration of one or more therapeutic agents to a subject with such a disease.

**[0072]** As used herein, the term “therapeutic agent” refers to any agent that can be used in the prevention, treatment, or management of a disease or disorder associated with overexpression of EphA2 and/or cell hyperproliferative diseases or disorders, particularly, cancer. In certain embodiments, the term “therapeutic agent” refers to an EphA2 vaccine of the invention, such as a composition comprising an EphA2 antigenic peptide, an EphA2 antigenic peptide expression vehicle, or an antigen presenting cell sensitized with an EphA2 antigenic peptide. In certain other embodiments, the term “therapeutic agent” refers to cancer chemotherapeutics, radiation therapy, hormonal therapy, and/or biological therapy/immunotherapy. In other embodiments, more than one therapeutic agent may be administered in combination.

**[0073]** As used herein, a “therapeutically effective amount” refers to that amount of the therapeutic agent sufficient to treat or manage a disease or disorder associated with EphA2 overexpression and/or cell hyperproliferative disease and, preferably, the amount sufficient to destroy, modify, control or remove primary, regional or metastatic cancer tissue. A therapeutically effective amount may refer to the amount of therapeutic agent sufficient to delay or minimize the onset of the hyperproliferative disease, e.g., delay or minimize the spread of cancer. A therapeutically effective amount may also refer to the amount of the therapeutic agent that provides a therapeutic benefit in the treatment or management of cancer. Further, a therapeutically effective amount with respect to a therapeutic agent of the invention means that amount of therapeutic agent alone, or in combination with other therapies, that provides a therapeutic benefit in the treatment or management of hyperproliferative disease or cancer. Used in connection with an amount of an EphA2 vaccine of the invention, the term can encompass an amount that improves

overall therapy, reduces or avoids unwanted effects, or enhances the therapeutic efficacy of or synergies with another therapeutic agent.

**[0074]** As used herein, the terms “T cell malignancies” and “T cell malignancy” refer to any T cell lymphoproliferative disorder, including thymic and post-thymic

malignancies. T cell malignancies include tumors of T cell origin. T cell malignancies refer to tumors of lymphoid progenitor cell, thymocyte, T cell, NK-cell, or antigen presenting cell origin. T cell malignancies include, but are not limited to, leukemias, including acute lymphoblastic leukemias, thymomas, acute lymphoblastic leukemias, and lymphomas, including Hodgkin’s and non-Hodgkin’s disease, with the proviso that T cell

malignancies are not cutaneous T cell malignancies, in particular cutaneous-cell lymphomas. In a preferred embodiment, T cell malignancies are systemic, non-cutaneous T cell malignancies.

### **3.2 SEQUENCES**

**[0075]** Below is a brief summary of the sequence presented in the accompanying sequence listing, which is incorporated by reference herein in its entirety:

**[0076]** **SEQ ID NO:1**

Human EphA2 cDNA (full length)

**[0077]** **SEQ ID NO:2**

Human EphA2 polypeptide (full length)

**[0078]** **SEQ ID NOs:3-18**

Human EphA2 peptides

**[0079]** **SEQ ID NO:19**

Construct: LLOss-PEST-hEphA2

Native LLO signal peptide + PEST fused to full-length human EphA2

Not Codon optimized

No epitope tags (e.g., myc or FLAG used in this construct)

Fusion protein coding sequence shown

**[0080]** **SEQ ID NO:20**

Construct: LLOss-PEST-hEphA2

Native LLO signal peptide + PEST fused to full-length human EphA2

Not Codon optimized

No epitope tags (e.g., myc or FLAG used in this construct)

Predicted fusion protein shown

**[0081]** **SEQ ID NO:21**

EphA2 EX2 domain

		Native nucleotide sequence
	[0082]	<b>SEQ ID NO:22</b>
		EphA2 EX2 domain
		Nucleotide sequence for optimal codon usage in <i>Listeria</i>
5	[0083]	<b>SEQ ID NO:23</b>
		EphA2 EX2 domain
		Primary Amino Acid Sequence
	[0084]	<b>SEQ ID NO:24</b>
		Construct: LLOss-PEST-EX2_hEphA2
10		Native LLO signal peptide + PEST fused to external domain of human EphA2
		Not Codon optimized
		No epitope tags (e.g., myc or FLAG used in this construct)
	[0085]	<b>SEQ ID NO:25</b>
15		Construct: LLOss-PEST-EX2_hEphA2
		Native LLO signal peptide + PEST fused to external domain of human EphA2
		Not Codon optimized
		No epitope tags (e.g., myc or FLAG used in this construct)
20		Predicted fusion protein shown
	[0086]	<b>SEQ ID NO:26</b>
		NativeLLOss-PEST-FLAG-EX2_EphA2-myc-CodonOp
		(Native <i>L. monocytogenes</i> LLO signal peptide + PEST-Codon optimized - FLAG-EX-2 EphA2-Myc)
25		Nucleotide Sequence (including <i>hly</i> promoter)
	[0087]	<b>SEQ ID NO:27</b>
		NativeLLOss-PEST-FLAG-EX2_EphA2-myc-CodonOp
		(Native <i>L. monocytogenes</i> LLO signal peptide + PEST-Codon optimized - FLAG-EX-2 EphA2-Myc)
30		Primary Amino Acid Sequence
	[0088]	<b>SEQ ID NO:28</b>
		Codon Optimized LLOss-PEST-FLAG-EX2_EphA2-myc-CodonOp
		(Codon Optimized <i>L. monocytogenes</i> LLO signal peptide + PEST-Codon optimized -FLAG-EX-2 EphA2-Myc)
35		Nucleotide Sequence (including <i>hly</i> promoter)

	[0089]	<b>SEQ ID NO:29</b>
		Codon Optimized LLOss-PEST-FLAG-EX2_EphA2-myc-CodonOp (Codon Optimized <i>L. monocytogenes</i> LLO signal peptide + PEST-Codon optimized -FLAG-EX-2 EphA2-Myc)
5		Primary Amino Acid Sequence
	[0090]	<b>SEQ ID NO:30</b>
		PhoD-FLAG-EX2_EphA2-myc-CodonOp (Codon optimized <i>B. subtilis</i> phoD Tat signal peptide-FLAG-EX-2 EphA2- Myc)
10		Nucleotide Sequence (including <i>hly</i> promoter)
	[0091]	<b>SEQ ID NO:31</b>
		PhoD-FLAG-EX2_EphA2-myc-CodonOp (Codon optimized <i>B. subtilis</i> phoD Tat signal peptide-FLAG-EX-2 EphA2- Myc)
15		Amino acid sequence
	[0092]	<b>SEQ ID NO:32</b>
		EphA2 CO domain
		Native nucleotide sequence
	[0093]	<b>SEQ ID NO:33</b>
20		EphA2 CO domain
		Nucleotide sequence for optimal codon usage in <i>Listeria</i>
	[0094]	<b>SEQ ID NO:34</b>
		EphA2 CO domain
		Primary Amino Acid Sequence
25	[0095]	<b>SEQ ID NO:35</b>
		Construct: LLOss-PEST-CO-huEphA2 Native LLO signal peptide + PEST fused to cytoplasmic domain of human EphA2 Not Codon optimized
30		No epitope tags (e.g., myc or FLAG used in this construct) Fusion protein coding sequence shown
	[0096]	<b>SEQ ID NO:36</b>
		Construct: LLOss-PEST-CO-huEphA2 Native LLO signal peptide + PEST fused to cytoplasmic domain of human EphA2
35		



	Not Codon optimized
	No epitope tags (e.g., myc or FLAG used in this construct)
	Predicted fusion protein shown
[0097]	<b>SEQ ID NO:37</b>
5	NativeLLOss-PEST-FLAG-CO_EphA2-myc-CodonOp (Native L. monocytogenes LLO signal peptide + PEST-Codon optimized - FLAG-CO_EphA2-Myc) Nucleotide Sequence (including <i>hly</i> promoter)
[0098]	<b>SEQ ID NO:38</b>
10	NativeLLOss-PEST-FLAG-CO_EphA2-myc-CodonOp (Native L. monocytogenes LLO signal peptide + PEST-Codon optimized - FLAG-CO_EphA2-Myc) Primary Amino Acid Sequence
[0099]	<b>SEQ ID NO:39</b>
15	Codon Optimized LLOss-PEST-FLAG-CO_EphA2-myc-CodonOp (Codon Optimized L. monocytogenes LLO signal peptide + PEST-Codon optimized -FLAG-CO_EphA2-Myc) Nucleotide Sequence (including <i>hly</i> promoter)
[00100]	<b>SEQ ID NO:40</b>
20	Codon Optimized LLOss-PEST-FLAG-CO_EphA2-myc-CodonOp (Codon Optimized L. monocytogenes LLO signal peptide + PEST-Codon optimized -FLAG-CO_EphA2-Myc) Primary Amino Acid Sequence
[00101]	<b>SEQ ID NO:41</b>
25	PhoD-FLAG-CO_EphA2-myc-CodonOp (Codon optimized B. subtilis phoD Tat signal peptide-FLAG-CO_EphA2- Myc) Nucleotide Sequence (including <i>hly</i> promoter)
[00102]	<b>SEQ ID NO:42</b>
30	PhoD-FLAG-CO_EphA2-myc-CodonOp (Codon optimized B. subtilis phoD Tat signal peptide-FLAG-CO_EphA2- Myc) Amino acid sequence
[00103]	<b>SEQ ID NO:43</b>
35	Construct: pAM401-MCS

Plasmid pAM401 containing multiple cloning site (MCS) from pPL2 vector  
 Insertion of small *Aat II* MCS fragment from pPL2 inserted into pAM401  
 plasmid between blunted *Xba I* and *Nru I* sites.  
 Complete pAM401-MCS plasmid sequence shown

#### 4. BRIEF DESCRIPTION OF THE FIGURES

**[00104]** **Figure 1.** *Listeria* intracellular life cycle, antigen presenting cell activation, and antigen presentation.

**[00105]** **Figure 2.** Western blot analysis of secreted protein from recombinant *Listeria* encoding native EphA2 CO domain sequence.

**[00106]** **Figure 3.** Western blot analysis of secreted protein from recombinant *Listeria* encoding native or codon-optimized LLO secA1 signal peptide fused with codon-optimized EphA2 EX2 domain sequence signal peptide.

**[00107]** **Figure 4.** Western blot analysis of secreted protein from recombinant *Listeria* encoding native or codon-optimized LLO secA2 signal peptide or codon-optimized Tat signal peptide fused with codon-optimized EphA2 CO domain sequence.

**[00108]** **Figure 5.** Flow cytometry analysis of human EphA2 expression in CT2 murine carcinoma cells. Single cell FACS sorting assays were performed by standard techniques to identify CT26 cell clones expressing high levels of human EphA2.

**[00109]** **Figure 6.** Western blot analysis of pooled populations CT26 murine colon carcinoma cells expressing high levels of human EphA2 protein.

**[00110]** **Figure 7.** Flow Cytometry of B16F10 cells expressing huEphA2.

**[00111]** **Figure 8.** Western blot analysis of lysate from 293 cells 48 hr. following transfection with pCDNA4 plasmid DNA encoding full-length native EphA2 sequence.

**[00112]** **Figure 9A-9B.** In the CT26 tumor model, therapeutic immunization with positive control *Listeria* expressing AH1-A5.

**[00113]** **Figure 10A-10B.** Preventative immunization with *Listeria* expressing ECD of hEphA2 suppresses CT26-hEphA2 tumor growth (**Figure 10A**) and increases survival (**Figure 10B**).

**[00114]** **Figures 11A-11D.** Preventive studies following i.v. administration of L4029EphA2-exFlag, *Listeria* control (L4029), or *Listeria* positive control containing the AH1 protein (L4029-AH1) ( $5 \times 10^5$  cells in 100  $\mu$ l volume) either subcutaneously or intravenously. **Figure 11A** demonstrates tumor volume of mice inoculated with CT26 cells expressing the ECD of huEphA2, vehicle (HBSS), *Listeria* (L4029) or *Listeria* positive (L4029-AH1) controls. **Figure 11B** demonstrates mean tumor volume of mice inoculated with CT26 cells expressing the ECD of huEphA2 (L4029-EphA2 exFlag) compared to the

- Listeria* (L4029) control. **Figure 11C** illustrates results of the prevention study in the s.c. model, measuring percent survival of the mice post CT26 tumor cell inoculation. **Figure 11D** illustrates the results of the prevention study in the lung metastases model, measuring percent survival of the mice post tumor cell inoculation.
- 5 [00115] **Figure 12.** Preventative immunization with *Listeria* expressing ECD of hEphA2 increases survival following RenCa-hEphA2 tumor challenge.
- [00116] **Figure 13A-13C.** Figures 13A-13C illustrate results of a typical therapeutic study of animals inoculated with CT26 murine colon carcinoma cells transfected with human EphA2 (L4029-EphA2 exFlag), *Listeria* control (L4029-control) or vehicle (HBSS).
- 10 In **Figure 13A**, tumor volume was measured at several intervals post inoculation. **Figure 13B** illustrates the mean tumor volume of mice inoculated with CT26 cells containing either *Listeria* control or the ECD of huEphA2. **Figure 13C** represents the results of a therapeutic study using the lung metastases model, measuring percent survival of mice post inoculation with CT26 cells with either HBSS or *Listeria* control, or *Listeria* expressing the ECD of
- 15 huEphA2.
- [00117] **Figure 14A-F.** **Figure 14A.** Therapeutic immunization in Balb/C mice with *Listeria* expressing ICD of hEphA2 suppresses established CT26-hEphA2 tumor growth. **Figure 14B.** Immunization of Balb/C mice bearing CT26.24 (huEphA2+) lung tumors with recombinant *Listeria* encoding EphA2 CO domain confers long-term survival.
- 20 **Figure 14C.** Long-term survival of Balb/C mice bearing CT26.24 (huEphA2+) lung tumors immunized with recombinant *Listeria* encoding OVA.AH1 or OVA.AH1-A5. **Figure 14D.** Increased survival of Balb/C mice bearing CT26.24 (huEphA2+) lung tumors when immunized with recombinant *Listeria* encoding codon-optimized or native EphA2 CO domain sequence. **Figure 14E.** Increased survival of Balb/C mice bearing CT26.24 (huEphA2+) lung tumors when immunized with recombinant *Listeria* encoding codon-optimized secA1 signal peptide fused with codon-optimized EphA2 EX2 domain sequence. **Figure 14F.** Immunization of Balb/C mice bearing CT26.24 (huEphA2+) lung tumors with recombinant *Listeria* encoding EphA2 CO domain but not with plasmid DNA encoding full-length EphA2 confers long-term survival.
- 30 [00118] **Figure 15.** Long-term suppression of CT26-hEphA2 tumor growth upon rechallenge.
- [00119] **Figure 16.** Immunization with *Listeria* expressing hEphA2 elicits a specific CD8+ T cell response.
- [00120] **Figure 17.** Both CD4+ and CD8+ T cell responses are required for optimal
- 35 hEphA2-directed anti-tumor efficacy.

[00121] **Figures 18A-B.** Therapeutic vaccination with *Listeria* expressing human EphA2 ICD enhances CD45+ tumor infiltrate. **Figure 18A** depicts images of tumor sections stained with biotinylated rat anti-mouse CD45/B200. **Figure 18B** is a bar graph normalizing the image data to tumor volume.

## 5. DETAILED DESCRIPTION OF THE INVENTION

[00122] The present invention is based, in part, on the inventors' discovery that a vaccine that comprises an EphA2 antigenic peptide can confer beneficial immune response against EphA2-expressing cells in involved in a hyperproliferative disease such as cancer. In particular, such a vaccine can contain an EphA2 antigenic peptide, an expression vehicle  
10 for an EphA2 antigenic peptide, or an antigen-presenting cell that is sensitized with an EphA2 antigenic peptide.

[00123] Accordingly, the present invention relates to methods and compositions that provide for the treatment, inhibition, and management of diseases and disorders associated with overexpression of EphA2 and/or cell hyperproliferative diseases and disorders. A  
15 particular aspect of the invention relates to methods and compositions containing compounds that, when administered to a subject with a hyperproliferative disorder involving EphA2-expressing cells, either elicit or mediate an immune response against EphA2, resulting in a growth inhibition of the EphA2-expressing cells involved in the hyperproliferative disorder. The present invention further relates to methods and  
20 compositions for the treatment, inhibition, or management of metastases of cancers of epithelial cell origin, especially human cancers of the breast, ovary, oesophagus, lung, skin, prostate, bladder, and pancreas, and renal cell carcinomas and melanomas. The invention further relates to methods and compositions for the treatment, inhibition, or management of cancers of T cell origin, especially leukemias and lymphomas. Further compositions and  
25 methods of the invention include other types of active ingredients in combination with the EphA2 vaccines of the invention. In other embodiments, the methods of the invention are used to treat, prevent or manage other diseases or disorders associated with cell hyperproliferation, for example but not limited to asthma, psoriasis, restenosis, COPD, *etc.*

[00124] The present invention also relates to methods for the treatment, inhibition,  
30 and management of cancer or other hyperproliferative cell disorders or diseases that have become partially or completely refractory to current or standard cancer treatment, such as chemotherapy, radiation therapy, hormonal therapy, and biological-/immuno-therapy.

### 5.1 EphA2 Antigenic Peptides

- [00125]** As discussed above, the invention encompasses administration of exogenous EphA2 antigenic peptides that are capable of eliciting an immune response to EphA2, resulting in a cellular or humoral immune response against endogenous EphA2. Additionally, the present invention encompasses the use of an EphA2 antigenic peptide expression vehicle.
- [00126]** In principle, an EphA2 antigenic peptide (sometimes referred to as an “EphA2 antigenic polypeptide”) for use in the methods and compositions of the present invention can be any EphA2 antigenic peptide that is capable of eliciting an immune response against EphA2-expressing cells involved in a hyperproliferative disorder. Thus, an EphA2 antigenic peptide can be an EphA2 polypeptide, preferably an EphA2 polypeptide of SEQ ID NO:2, or a fragment or derivative of an EphA2 polypeptide that (1) displays antigenicity of EphA2 (ability to bind or compete with EphA2 for binding to an anti-EphA2 antibody), (2) displays immunogenicity of EphA2 (ability to generate antibody which binds to EphA2), or (3) contains one or more T cell epitopes of EphA2.
- [00127]** In certain embodiments, the EphA2 antigenic peptide is full length human EphA2 (SEQ ID NO:2).
- [00128]** In other embodiments, the EphA2 antigenic peptide comprises the intracellular domain of EphA2 (residue 22 to 554 of SEQ ID NO:2).
- [00129]** In yet other embodiments, the EphA2 antigenic peptide comprises the intracellular domain EphA2 (residue 558 to 976 of SEQ ID NO:2).
- [00130]** In certain embodiments, the peptide corresponds to or comprises an EphA2 epitope that is exposed in a cancer cell but occluded in a non-cancer cell. In a preferred embodiment, the EphA2 antigenic peptides preferentially include epitopes on EphA2 that are selectively exposed or increased on cancer cells but not non-cancer cells (“exposed EphA2 epitope peptides”).
- [00131]** The present invention further encompasses the use of a plurality of EphA2 antigenic peptides in the compositions and methods of the present invention. In certain embodiments, the plurality of EphA2 antigenic peptides are multimerized or polyvalent.
- [00132]** Fragments of EphA2 that are useful in the methods and compositions present invention may contain deletions, additions or substitutions of amino acid residues within the amino acid sequence encoded by an EphA2 gene. Preferably mutations result in a silent change, thus producing a functionally equivalent EphA2 gene product.
- [00133]** An EphA2 antigenic polypeptide sequence preferably comprises an amino acid sequence that exhibits at least about 65% sequence similarity to human EphA2, more preferably exhibits at least 70% sequence similarity to human EphA2, yet more preferably

exhibits at least about 75% sequence similarity human EphA2. In other embodiments, the EphA2 polypeptide sequence preferably comprises an amino acid sequence that exhibits at least 85% sequence similarity to human EphA2, yet more preferably exhibits at least 90% sequence similarity to human EphA2, and most preferably exhibits at least about 95%

sequence similarity to human EphA2.

**[00134]** Additional polypeptides suitable in the present methods are those encoded by the nucleic acids described in Section 5.2 below.

**[00135]** The determination of percent identity between two sequences can be accomplished using a mathematical algorithm. A preferred, non-limiting example of a mathematical algorithm utilized for the comparison of two sequences is the algorithm of Karlin and Altschul, 1990, *Proc Natl Acad Sci. USA* 87:2264-2268, modified as in Karlin and Altschul, 1993, *Proc Natl Acad Sci. USA* 90:5873-5877. Such an algorithm is incorporated into the NBLAST and XBLAST programs of Altschul *et al.*, 1990, *J. Mol. Biol.* 215:403-410. BLAST nucleotide searches can be performed with the NBLAST program, score = 100, wordlength = 12 to obtain nucleotide sequences homologous to a nucleic acid molecules of the invention. BLAST protein searches can be performed with the XBLAST program, score = 50, wordlength = 3 to obtain amino acid sequences homologous to a protein molecules of the invention. To obtain gapped alignments for comparison purposes, Gapped BLAST can be utilized as described in Altschul *et al.*, 1997, *Nucleic Acids Res.* 25:3389-3402. Alternatively, PSI-Blast can be used to perform an iterated search which detects distant relationships between molecules (*Id.*). When utilizing BLAST, Gapped BLAST, and PSI-Blast programs, the default parameters of the respective programs (e.g., XBLAST and NBLAST) can be used.

**[00136]** Another preferred, non limiting example of a mathematical algorithm utilized for the comparison of sequences is the algorithm of Myers and Miller, 1988, *CABIOS* 4:11 17. Such an algorithm is incorporated into the ALIGN program (version 2.0) which is part of the GCG sequence alignment software package. When utilizing the ALIGN program for comparing amino acid sequences, a PAM120 weight residue table, a gap length penalty of 12, and a gap penalty of 4 can be used. Additional algorithms for sequence analysis are known in the art and include ADVANCE and ADAM as described in Torellis and Robotti, 1994, *Comput. Appl. Biosci.* 10:3-5; and FASTA described in Pearson and Lipman, 1988, 85:2444-8. Within FASTA, ktup is a control option that sets the sensitivity and speed of the search. If ktup = 2, similar regions in the two sequences being compared are found by looking at pairs of aligned residues; if ktup = 1, single aligned amino acids are examined. ktup can be set to 2 or 1 for protein sequences, or from 1 to 6

for DNA sequences. The default if ktup is not specified is 2 for proteins and 6 for DNA. For a further description of FASTA parameters, *see* <http://bioweb.pasteur.fr/docs/man/man/fasta.1.html#sect2>.

[00137] The percent identity between two sequences can be determined using techniques similar to those described above, with or without allowing gaps. In calculating percent identity, only exact matches are counted. However, conservative substitutions should be considered in evaluating sequences that have a low percent identity with the EphA2 sequences disclosed herein.

[00138] In a specific embodiment, EphA2 antigenic polypeptides comprising at least 10, 20, 30, 40, 50, 75, 100, or 200 amino acids of an EphA2 polypeptide of SEQ ID NO:2 are used in the present invention. In a preferred embodiment, such a polypeptide comprises all or a portion of the extracellular domain of an EphA2 polypeptide of SEQ ID NO:2.

## **5.2 Methods of Identifying EphA2 Antigenic Peptides**

### **5.2.1 Assays for EphA2 Antigenic Peptides**

[00139] The present invention provides *Listeria*-based EphA2 vaccines comprising *Listeria* bacteria engineered to express an EphA2 antigenic peptide. Any assay known in the art for determining whether a peptide is a T cell epitope or a B cell epitope may be employed in testing EphA2 peptides for suitability in the present methods and compositions.

[00140] For example, ELISPOT assays and methods for intracellular cytokine staining can be used for enumeration and characterization of antigen-specific CD4<sup>+</sup> and CD8<sup>+</sup> T cells. Lalvani *et al.* (1997) *J. Exp. Med.* 186:859-865; Waldrop *et al.* (1997) *J. Clin. Invest.* 99:1739-1750.

[00141] EphA2 antigenic peptides can be determined by screening synthetic peptides corresponding to portions of EphA2. Candidate antigenic peptides can be identified on the basis of their sequence or predicted structure. A number of algorithms are available for this purpose.

[00142] Exemplary protocols for such assays are presented below.

### **5.2.2 Peptides That Display Immunogenicity of EphA2**

[00143] The ability of EphA2 peptides to elicit EphA2-specific antibody responses in mammals can be examined, for example, by immunizing animals (*e.g.*, mice, guinea pigs or rabbits) with individual EphA2 peptides emulsified in Freund's adjuvant.

[00145] EphA2 peptides which produce antisera that react specifically with the EphA2 peptides and also recognized full length EphA2 protein in immunoblots are said to display the antigenicity of EphA2.

### 5.2.3 CD4<sup>+</sup> T-Cell Proliferation Assay

- 5 [00146] For example, such assays include *in vitro* cell culture assays in which peripheral blood mononuclear cells ("PBMCs") are obtained from fresh blood of a patient with a disease involving overexpression of EphA2, and purified by centrifugation using FICOLL-PLAQUE PLUS (Pharmacia, Upsalla, Sweden) essentially as described by Kruse and Sebald, 1992, *EMBO J.* 11:3237-3244. The peripheral blood mononuclear cells are
- 10 incubated for 7-10 days with candidate EphA2 antigenic peptides. Antigen presenting cells may optionally be added to the culture 24 to 48 hours prior to the assay, in order to process and present the antigen. The cells are then harvested by centrifugation, and washed in RPMI 1640 media (GibcoBRL, Gaithersburg, MD).  $5 \times 10^4$  activated T cells/well are in RPMI 1640 media containing 10% fetal bovine serum, 10 mM HEPES, pH 7.5, 2 mM L-
- 15 glutamine, 100 units/ml penicillin G, and 100  $\mu$ g/ml streptomycin sulphate in 96 well plates for 72 hrs at 37°C, pulsed with 1  $\mu$ Ci <sup>3</sup>H-thymidine (DuPont NEN, Boston, MA)/well for 6 hrs, harvested, and radioactivity measured in a TOPCOUNT scintillation counter (Packard Instrument Co., Meriden, CT).

### 5.2.4 Intracellular Cytokine Staining (ICS)

- 20 [00147] Measurement of antigen-specific, intracellular cytokine responses of T cells can be performed essentially as described by Waldrop *et al.*, 1997, *J. Clin. Invest.* 99:1739-1750; Openshaw *et al.*, 1995, *J. Exp. Med.* 182:1357-1367; or Estcourt *et al.*, 1997, *Clin. Immunol. Immunopathol.* 83:60-67. Purified PBMCs from patients with a disease involving EphA2-overexpressing cells are placed in 12x75 millimeter polystyrene tissue culture tubes
- 25 (Becton Dickinson, Lincoln Park, N.J.) at a concentration of  $1 \times 10^6$  cells per tube. A solution comprising 0.5 milliliters of HL-1 serum free medium, 100 units per milliliter of penicillin, 100 units per milliliter streptomycin, 2 millimolar L glutamine (Gibco BRL), varying amounts of individual EphA2 antigenic candidate peptides, and 1 unit of anti-CD28 mAb (Becton-Dickinson, Lincoln Park, N.J.) is added to each tube. Anti-CD3 mAb is
- 30 added to a duplicate set of normal PBMC cultures as positive control. Culture tubes are incubated for 1 hour. Brefeldin A is added to individual tubes at a concentration of 1 microgram per milliliter, and the tubes are incubated for an additional 17 hours.
- [00148] PBMCs stimulated as described above are harvested by washing the cells twice with a solution comprising Dulbecco's phosphate-buffered saline (dPBS) and 10 units
- 35 of Brefeldin A. These washed cells are fixed by incubation for 10 minutes in a solution



comprising 0.5 milliliters of 4% paraformaldehyde and dPBS. The cells are washed with a solution comprising dPBS and 2% fetal calf serum (FCS). The cells are then either used immediately for intracellular cytokine and surface marker staining or are frozen for no more than three days in freezing medium, as described (Waldrop *et al.*, 1997, *J. Clin. Invest.*

99:1739-1750).

**[00149]** The cell preparations were rapidly thawed in a 37°C water bath and washed once with dPBS. Cells, either fresh or frozen, are resuspended in 0.5 milliliters of permeabilizing solution (Becton Dickinson Immunocytometry systems, San Jose, Calif.) and incubated for 10 minutes at room temperature with protection from light.

Permeabilized cells are washed twice with dPBS and incubated with directly conjugated mAbs for 20 minutes at room temperature with protection from light. Optimal concentrations of antibodies are predetermined according to standard methods. After staining, the cells were washed, refixed by incubation in a solution comprising dPBS 1% paraformaldehyde, and stored away from light at 4°C for flow cytometry analysis.

### **5.2.5 ELISPOT Assays**

**[00150]** The ELISPOT assay measures Th1-cytokine specific induction in murine splenocytes following *Listeria* vaccination. ELISPOT assays are performed to determine the frequency of T lymphocytes in response to endogenous antigenic peptide stimulation, and are as described in Geginat, et al., 2001, *J. Immunol.* 166:1877-1884. Balb/c mice (3 per group) are vaccinated with *L. monocytogenes* expressing candidate EphA2 antigenic peptides or HBSS as control. Whole mouse spleens are harvested and pooled five days after vaccination. Single cell suspensions of murine splenocytes are plated in the presence of various antigens overnight in a 37°C incubator.

**[00151]** Assays are performed in nitrocellulose-backed 96-well microtiter plates coated with rat anti-mouse IFN-  $\gamma$  mAb. For the testing of the candidate EphA2 antigenic peptide, a  $1 \times 10^{-5}$  M peptide solution is prepared. In round-bottom 96-well microtiter plates per well  $6 \times 10^5$  unseparated splenocytes in 135  $\mu$ l culture medium (a modification of Eagle's medium (Life Technologies, Eggenstein, Germany) supplemented with 10% FCS, 100 U/ml penicillin, 100  $\mu$ g/ml streptomycin,  $1 \times 10^{-5}$  M 2-ME, and 2 mM glutamine) are mixed with 15  $\mu$ l of the  $1 \times 10^{-5}$  M peptide solution to yield a final peptide concentration of  $1 \times 10^{-6}$  M. After 6 h of incubation at 37°C, cells are resuspended by vigorous pipetting, and 100  $\mu$ l or 10  $\mu$ l of cell suspension ( $4 \times 10^5$ /well or  $4 \times 10^4$ /well, respectively) is transferred to Ab-coated ELISPOT plates and incubated overnight at 37°C. In the ELISPOT plates, the final volume was adjusted to 150  $\mu$ l to ensure homogenous distribution of cells.

[00152] Purified CD4<sup>+</sup> or CD8<sup>+</sup> T cells are tested in a modified assay as follows: 15  
 μl prediluted peptide (1 x 10<sup>-5</sup> M) is directly added to Ab-coated ELISPOT plates and  
 mixed with 4 x 10<sup>5</sup> splenocytes from nonimmune animals as APC to yield a final volume of  
 100 μl. After 4 h of preincubation of APC at 37°C, 1 x 10<sup>5</sup> CD4<sup>+</sup> or CD8<sup>+</sup> cells purified  
 5 from *L. monocytogenes*-immune mice are added per well in a volume of 50 μl and plates are  
 incubated overnight at 37°C. The ELISPOT-based *ex vivo* MHC restriction analysis is  
 performed after loading of cell lines expressing specific MHC class I molecules with 1 x 10<sup>-7</sup>  
 6 M peptide for 2 h at 37°C. Subsequently, unbound peptides are washed off (four times) to  
 prevent binding of peptides to responder splenocytes. Per well of the ELISPOT plate, 1 x  
 10 10<sup>5</sup> peptide-loaded APC are mixed with 4 x 10<sup>5</sup> or 4 x 10<sup>4</sup> responder splenocytes in a final  
 volume of 150 μl. After overnight incubation at 37°C, ELISPOT plates are developed with  
 biotin-labeled rat anti-mouse IFN- γ mAb, HRP streptavidin conjugate, and  
 aminoethylcarbazole dye of spots per splenocytes seeded. The specificity and sensitivity of  
 the ELISPOT assay is controlled with IFN- γ secreting CD8 T cell lines specific for a  
 15 control antigen.

### 5.3 Fusion Proteins

[00153] In certain embodiments of the present invention, an EphA2 vaccine  
 comprises, expresses or is an antigen-presenting cell that is sensitized with an EphA2  
 antigenic peptide that is a fusion protein. Thus, the present invention encompasses  
 20 compositions and methods in which the EphA2 antigenic peptides are fusion proteins  
 comprising all or a fragment or derivative of EphA2 operatively associated to a  
 heterologous component, *e.g.*, a heterologous peptide. Heterologous components can  
 include, but are not limited to sequences which facilitate isolation and purification of the  
 fusion protein. Heterologous components can also include sequences which confer stability  
 25 to EphA2 antigenic peptides. Such fusion partners are well known to those of skill in the  
 art.

[00154] The present invention encompasses the use of fusion proteins comprising an  
 EphA2 polypeptide (*e.g.*, a polypeptide of SEQ ID NO:2) and a heterologous polypeptide  
 (*i.e.*, an unrelated polypeptide or fragment thereof, preferably at least 10, at least 20, at least  
 30 30, at least 40, at least 50, at least 60, at least 70, at least 80, at least 90 or at least 100 amino  
 acids of the polypeptide). The fusion can be direct, but may occur through linker  
 sequences. The heterologous polypeptide may be fused to the N-terminus or C-terminus of  
 the EphA2 antigenic polypeptide. Alternatively, the heterologous polypeptide may be  
 flanked by EphA2 polypeptide sequences.

- [00155] A fusion protein can comprise an EphA2 antigenic polypeptide fused to a heterologous signal sequence at its N-terminus. Various signal sequences are commercially available. Eukaryotic heterologous signal sequences include, but are not limited to, the secretory sequences of melittin and human placental alkaline phosphatase (Stratagene, La Jolla, CA). Prokaryotic heterologous signal sequences useful in the methods of the invention include, but are not limited to, the phoA secretory signal (Sambrook *et al.*, eds., 1989, Molecular Cloning: A Laboratory Manual, 2nd ed., Cold Spring Harbor Laboratory, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY) and the protein A secretory signal (Pharmacia Biotech, Piscataway, NJ).
- 10 [00156] The EphA2 antigenic polypeptide can be fused to tag sequences, *e.g.*, a hexahistidine peptide, such as the tag provided in a pQE vector (QIAGEN, Inc., Chatsworth, CA), among others, many of which are commercially available for use in the methods of the invention. As described in Gentz *et al.*, 1989, *Proc. Natl. Acad. Sci. USA*, 86:821-824, for instance, hexahistidine provides for convenient purification of the fusion protein. Other
- 15 examples of peptide tags are the hemagglutinin "HA" tag, which corresponds to an epitope derived from the influenza hemagglutinin protein (Wilson *et al.*, 1984, *Cell*, 37:767) and the "flag" tag (Knappik *et al.*, 1994, *Biotechniques*, 17(4):754-761). These tags are especially useful for purification of recombinantly produced EphA2 antigenic polypeptides.
- [00157] Any fusion protein may be readily purified by utilizing an antibody specific or selective for the fusion protein being expressed. For example, a system described by Janknecht *et al.* allows for the ready purification of non-denatured fusion proteins expressed in human cell lines (Janknecht *et al.*, 1991, *Proc. Natl. Acad. Sci. USA* 88:8972). In this system, the gene of interest is subcloned into a vaccinia recombination plasmid such that the open reading frame of the gene is translationally fused to an amino-terminal tag consisting
- 20 of six histidine residues. Extracts from cells infected with recombinant vaccinia virus are loaded onto Ni<sup>2+</sup> nitriloacetic acid-agarose columns and histidine-tagged proteins are selectively eluted with imidazole-containing buffers.
- [00158] An affinity label can also be fused at its amino terminal to the carboxyl terminal of the EphA2 antigenic polypeptide for use in the methods of the invention. The
- 30 precise site at which the fusion is made in the carboxyl terminal is not critical. The optimal site can be determined by routine experimentation. An affinity label can also be fused at its carboxyl terminal to the amino terminal of the EphA2 antigenic polypeptide for use in the methods and compositions of the invention.
- [00159] A variety of affinity labels known in the art may be used, such as, but not
- 35 limited to, the immunoglobulin constant regions (*see also* Petty, 1996, Metal-chelate

affinity chromatography, in *Current Protocols in Molecular Biology*, Vol. 2, Ed. Ausubel *et al.*, Greene Publish. Assoc. & Wiley Interscience), glutathione S-transferase (GST; Smith, 1993, *Methods Mol. Cell Bio.* 4:220-229), the *E. coli* maltose binding protein (Guan *et al.*, 1987, *Gene* 67:21-30), and various cellulose binding domains (U.S. Patent Nos. 5,496,934; 5,202,247; 5,137,819; Tomme *et al.*, 1994, *Protein Eng.* 7:117-123), *etc.* Other affinity labels are recognized by specific binding partners and thus facilitate isolation by affinity binding to the binding partner which can be immobilized onto a solid support. Some affinity labels may afford the EphA2 antigenic polypeptide novel structural properties, such as the ability to form multimers. These affinity labels are usually derived from proteins that normally exist as homopolymers. Affinity labels such as the extracellular domains of CD8 (Shiue *et al.*, 1988, *J. Exp. Med.* 168:1993-2005), or CD28 (Lee *et al.*, 1990, *J. Immunol.* 145:344-352), or fragments of the immunoglobulin molecule containing sites for interchain disulfide bonds, could lead to the formation of multimers.

**[00160]** As will be appreciated by those skilled in the art, many methods can be used to obtain the coding region of the above-mentioned affinity labels, including but not limited to, DNA cloning, DNA amplification, and synthetic methods. Some of the affinity labels and reagents for their detection and isolation are available commercially.

**[00161]** In certain embodiments, the affinity label is a non-variable portion of the immunoglobulin molecule. Typically, such portions comprise at least a functionally operative CH2 and CH3 domain of the constant region of an immunoglobulin heavy chain. Fusions are also made using the carboxyl terminus of the Fc portion of a constant domain, or a region immediately amino-terminal to the CH1 of the heavy or light chain. Suitable immunoglobulin-based affinity label may be obtained from IgG-1, -2, -3, or -4 subtypes, IgA, IgE, IgD, or IgM, but preferably IgG1. Many DNA encoding immunoglobulin light or heavy chain constant regions are known or readily available from cDNA libraries. See, for example, Adams *et al.*, *Biochemistry*, 1980, 19:2711-2719; Gough *et al.*, 1980, *Biochemistry*, 19:2702-2710; Dolby *et al.*, 1980, *Proc. Natl. Acad. Sci. U.S.A.*, 77:6027-6031; Rice *et al.*, 1982, *Proc. Natl. Acad. Sci. U.S.A.*, 79:7862-7865; Falkner *et al.*, 1982, *Nature*, 298:286-288; and Morrison *et al.*, 1984, *Ann. Rev. Immunol.*, 2:239-256. Because many immunological reagents and labeling systems are available for the detection of immunoglobulins, the EphA2 antigenic polypeptide-Ig fusion protein can readily be detected and quantified by a variety of immunological techniques known in the art, such as the use of enzyme-linked immunosorbent assay (ELISA), immunoprecipitation, fluorescence activated cell sorting (FACS), *etc.* Similarly, if the affinity label is an epitope with readily available antibodies, such reagents can be used with the techniques mentioned

above to detect, quantitate, and isolate the EphA2 antigenic polypeptide containing the affinity label. In many instances, there is no need to develop specific or selective antibodies to the EphA2 antigenic polypeptide for the purposes of purification.

**[00162]** A fusion protein can comprise an EphA2 antigenic polypeptide fused to the Fc domain of an immunoglobulin molecule or a fragment thereof for use in the methods and compositions of the invention. A fusion protein can also comprise an EphA2 antigenic polypeptide fused to the CH2 and/or CH3 region of the Fc domain of an immunoglobulin molecule. Furthermore, a fusion protein can comprise an EphA2 antigenic polypeptide fused to the CH2, CH3, and hinge regions of the Fc domain of an immunoglobulin molecule (see Bowen *et al.*, 1996, *J. Immunol.* 156:442-49). This hinge region contains three cysteine residues which are normally involved in disulfide bonding with other cysteines in the Ig molecule. Since none of the cysteines are required for the peptide to function as a tag, one or more of these cysteine residues may optionally be substituted by another amino acid residue, such as for example, serine.

**[00163]** Various leader sequences known in the art can be used for the efficient secretion of the EphA2 antigenic polypeptide from bacterial and mammalian cells (von Heijne, 1985, *J. Mol. Biol.* 184:99-105). Leader peptides are selected based on the intended host cell, and may include bacterial, yeast, viral, animal, and mammalian sequences. For example, the herpes virus glycoprotein D leader peptide is suitable for use in a variety of mammalian cells. A preferred leader peptide for use in mammalian cells can be obtained from the V-J2-C region of the mouse immunoglobulin kappa chain (Bernard *et al.*, 1981, *Proc. Natl. Acad. Sci.* 78:5812-5816). Preferred leader sequences for targeting EphA2 antigenic polypeptide expression in bacterial cells include, but are not limited to, the leader sequences of the *E. coli* proteins OmpA (Hobom *et al.*, 1995, *Dev. Biol. Stand.* 84:255-262), Pho A (Oka *et al.*, 1985, *Proc. Natl. Acad. Sci.* 82:7212-16), OmpT (Johnson *et al.*, 1996, *Protein Expression* 7:104-113), LamB and OmpF (Hoffman & Wright, 1985, *Proc. Natl. Acad. Sci. USA* 82:5107-5111),  $\beta$ -lactamase (Kadonaga *et al.*, 1984, *J. Biol. Chem.* 259:2149-54), enterotoxins (Morioka-Fujimoto *et al.*, 1991, *J. Biol. Chem.* 266:1728-32), and the *Staphylococcus aureus* protein A (Abrahmsen *et al.*, 1986, *Nucleic Acids Res.* 14:7487-7500), and the *B. subtilis* endoglucanase (Lo *et al.*, *Appl. Environ. Microbiol.* 54:2287-2292), as well as artificial and synthetic signal sequences (MacIntyre *et al.*, 1990, *Mol. Gen. Genet.* 221:466-74; Kaiser *et al.*, 1987, *Science*, 235:312-317).

**[00164]** In certain embodiments, the fusion partner comprises a non-EphA2 polypeptide corresponding to an antigen associated with the cell type against which a therapeutic or prophylactic immune is desired. For example, the non-EphA2 polypeptide

can comprise an epitope of a tumor-associated antigen, such as, but not limited to, MAGE-1, MAGE-2, MAGE-3, gp100, TRP-2, tyrosinase, MART-1,  $\beta$ -HCG, CEA, Ras,  $\beta$ -catenin, gp43, GAGE-1, BAGE-1, PSA, and MUC-1, 2, 3.

[00165] Fusion proteins can be produced by standard recombinant DNA techniques or by protein synthetic techniques, e.g., by use of a peptide synthesizer. For example, a nucleic acid molecule encoding a fusion protein can be synthesized by conventional techniques including automated DNA synthesizers. Alternatively, PCR amplification of gene fragments can be carried out using anchor primers which give rise to complementary overhangs between two consecutive gene fragments which can subsequently be annealed and reamplified to generate a chimeric gene sequence (see, e.g., *Current Protocols in Molecular Biology*, Ausubel *et al.*, eds., John Wiley & Sons, 1992).

[00166] The nucleotide sequence coding for a fusion protein can be inserted into an appropriate expression vector, i.e., a vector which contains the necessary elements for the transcription and translation of the inserted protein-coding sequence. The expression of a fusion protein may be regulated by a constitutive, inducible or tissue-specific or -selective promoter. It will be understood by the skilled artisan that fusion proteins, which can facilitate solubility and/or expression, and can increase the *in vivo* half-life of the EphA2 antigenic polypeptide and thus are useful in the methods of the invention. The EphA2 antigenic polypeptides or peptide fragments thereof, or fusion proteins can be used in any assay that detects or measures EphA2 antigenic polypeptides or in the calibration and standardization of such assay.

[00167] The methods of invention encompass the use of EphA2 antigenic polypeptides or peptide fragments thereof, which may be produced by recombinant DNA technology using techniques well known in the art. Thus, methods for preparing the EphA2 antigenic polypeptides of the invention by expressing nucleic acid containing EphA2 antigenic gene sequences are described herein. Methods which are well known to those skilled in the art can be used to construct expression vectors containing, e.g., EphA2 antigenic polypeptide coding sequences (including but not limited to nucleic acids encoding all or an antigenic portion of a polypeptide of SEQ ID NO:2) and appropriate transcriptional and translational control signals. These methods include, for example, *in vitro* recombinant DNA techniques, synthetic techniques, and *in vivo* genetic recombination. See, for example, the techniques described in Sambrook *et al.*, 1989, *supra*, and Ausubel *et al.*, 1989, *supra*. Alternatively, RNA capable of encoding EphA2 antigenic polypeptide sequences may be chemically synthesized using, for example, synthesizers (see, e.g., the techniques described in *Oligonucleotide Synthesis*, 1984, Gait, M.J. ed., IRL Press, Oxford).

[00168] In certain embodiments, the EphA2 antigenic polypeptide is functionally coupled to an internalization signal peptide, also referred to as a "protein transduction domain," that would allow its uptake into the cell nucleus. In certain specific embodiments, the internalization signal is that of Antennapedia (reviewed by Prochiantz, 1996, Curr.

- 5 Opin. Neurobiol. 6:629 634, Hox A5 (Chatelin *et al.*, 1996, *Mech. Dev.* 55:111 117), HIV TAT protein (Vives *et al.*, 1997, *J. Biol. Chem.* 272:16010 16017) or VP22 (Phelan *et al.*, 1998, *Nat. Biotechnol.* 16:440 443).

#### 5.4 Polynucleotides Encoding An EphA2 Antigenic Peptide

[00169] The present invention also encompasses compositions and methods that  
10 employ an EphA2 antigenic peptide expression vehicle.

[00170] In certain embodiments, the expression vehicles comprise or contain polynucleotides that hybridize under high stringency, intermediate or lower stringency hybridization conditions, *e.g.*, as defined *infra*, to polynucleotides that encode an EphA2 of the invention.

15 [00171] By way of example and not limitation, procedures using such conditions of low stringency for regions of hybridization of over 90 nucleotides are as follows (*see also* Shilo and Weinberg, 1981, *Proc. Natl. Acad. Sci. USA* 78:6789-6792). Filters containing DNA are pretreated for 6 hours at 40°C in a solution containing 35% formamide, 5X SSC, 50 mM Tris-HCl (pH 7.5), 5 mM EDTA, 0.1% PVP, 0.1% Ficoll, 1% BSA, and 500 µg/ml  
20 denatured salmon sperm DNA. Hybridizations are carried out in the same solution with the following modifications: 0.02% PVP, 0.02% Ficoll, 0.2% BSA, 100 µg/ml salmon sperm DNA, 10% (wt/vol) dextran sulfate, and 5-20 X 10<sup>6</sup> cpm <sup>32</sup>P-labeled probe is used. Filters are incubated in hybridization mixture for 18-20 h at 40°C, and then washed for 1.5 h at 55°C in a solution containing 2X SSC, 25 mM Tris-HCl (pH 7.4), 5 mM EDTA, and 0.1%  
25 SDS. The wash solution is replaced with fresh solution and incubated an additional 1.5 h at 60°C. Filters are blotted dry and exposed for autoradiography. If necessary, filters are washed for a third time at 65-68°C and re-exposed to film. Other conditions of low stringency which may be used are well known in the art (*e.g.*, as employed for cross-species hybridizations).

30 [00172] Also, by way of example and not limitation, procedures using such conditions of high stringency for regions of hybridization of over 90 nucleotides are as follows. Prehybridization of filters containing DNA is carried out for 8 h to overnight at 65°C in buffer composed of 6X SSC, 50 mM Tris-HCl (pH 7.5), 1 mM EDTA, 0.02% PVP, 0.02% Ficoll, 0.02% BSA, and 500 µg/ml denatured salmon sperm DNA. Filters are  
35 hybridized for 48 h at 65°C in prehybridization mixture containing 100 µg/ml denatured

salmon sperm DNA and  $5\text{-}20 \times 10^6$  cpm of  $^{32}\text{P}$ -labeled probe. Washing of filters is done at  $37^\circ\text{C}$  for 1 h in a solution containing 2X SSC, 0.01% PVP, 0.01% Ficoll, and 0.01% BSA. This is followed by a wash in 0.1X SSC at  $50^\circ\text{C}$  for 45 min before autoradiography.

[00173] Other conditions of high stringency which may be used depend on the nature  
5 of the nucleic acid (*e.g.*, length, GC content, *etc.*) and the purpose of the hybridization (detection, amplification, *etc.*) and are well known in the art. For example, stringent hybridization of a nucleic acid of approximately 15-40 bases to a complementary sequence in the polymerase chain reaction (PCR) is done under the following conditions: a salt concentration of 50 mM KCl, a buffer concentration of 10 mM Tris-HCl, a  $\text{Mg}^{2+}$   
10 concentration of 1.5 mM, a pH of 7-7.5 and an annealing temperature of  $55\text{-}60^\circ\text{C}$ .

[00174] Selection of appropriate conditions for moderate stringencies is also well known in the art (*see, e.g.*, Sambrook *et al.*, 1989, Molecular Cloning, A Laboratory Manual, 2d Ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York; *see also*, Ausubel *et al.*, eds., in the Current Protocols in Molecular Biology series of laboratory  
15 technique manuals, © 1987-1997, Current Protocols, © 1994-1997 John Wiley and Sons, Inc.).

[00175] The nucleic acids useful in the present methods may be made by any method known in the art. For example, if the nucleotide sequence of the EphA2 antigenic peptide is known, a nucleic acid encoding the peptide may be assembled from chemically synthesized  
20 oligonucleotides (*e.g.*, as described in Kutmeier *et al.*, 1994, *BioTechniques* 17:242), which, briefly, involves the synthesis of overlapping oligonucleotides containing portions of the sequence encoding the peptide, annealing and ligating of those oligonucleotides, and then amplification of the ligated oligonucleotides by PCR.



[00177] Further, a nucleic acid that is useful in the present methods may be manipulated using methods well known in the art for the manipulation of nucleotide sequences, e.g., recombinant DNA techniques, site directed mutagenesis, PCR, *etc.* (see, for example, the techniques described in Sambrook *et al.*, 1990, Molecular Cloning, A Laboratory Manual, 2d Ed., Cold Spring Harbor Laboratory, Cold Spring Harbor, NY and Ausubel *et al.*, eds., 1998, Current Protocols in Molecular Biology, John Wiley & Sons, NY, which are both incorporated by reference herein in their entireties), to generate EphA2 antigenic peptides having a different amino acid sequence from the amino acid sequence depicted in SEQ ID NO:2, for example to create amino acid substitutions, deletions, and/or insertions.

#### **5.5 Recombinant Expression Of An EphA2 antigenic peptide**

[00178] Recombinant expression of an EphA2 antigenic peptide of the invention, or fragment or derivative thereof, requires construction of an expression vector containing a polynucleotide that encodes the EphA2 antigenic peptide. Once a polynucleotide encoding an EphA2 antigenic peptide of the invention has been obtained, the vector for the production of the EphA2 antigenic peptide may be produced by recombinant DNA technology using techniques well known in the art. Thus, methods for preparing a protein by expressing a polynucleotide containing an EphA2 antigenic peptide-encoding nucleotide sequence are described herein. Methods which are well known to those skilled in the art can be used to construct expression vectors containing peptide coding sequences and appropriate transcriptional and translational control signals. These methods include, for example, *in vitro* recombinant DNA techniques, synthetic techniques, and *in vivo* genetic recombination. The invention, thus, provides replicable vectors comprising a nucleotide sequence encoding an EphA2 antigenic peptide of the invention.

[00179] The expression vector is transferred to a host cell by conventional techniques and the transfected cells are then cultured by conventional techniques to produce an EphA2 antigenic peptide of the invention. Thus, the invention includes host cells containing a polynucleotide encoding an EphA2 antigenic peptide of the invention or fragments thereof, operably linked to a heterologous promoter.

microorganisms such as bacteria (*e.g.*, *E. coli* and *B. subtilis*) transformed with recombinant bacteriophage DNA, plasmid DNA or cosmid DNA expression vectors containing peptide coding sequences; yeast (*e.g.*, *Saccharomyces Pichia*) transformed with recombinant yeast expression vectors containing peptide coding sequences; insect cell systems infected with recombinant virus expression vectors (*e.g.*, baculovirus) containing peptide coding sequences; plant cell systems infected with recombinant virus expression vectors (*e.g.*, cauliflower mosaic virus, CaMV; tobacco mosaic virus, TMV) or transformed with recombinant plasmid expression vectors (*e.g.*, Ti plasmid) containing peptide coding sequences; or mammalian cell systems (*e.g.*, COS, CHO, BHK, 293, NS0, and 3T3 cells) harboring recombinant expression constructs containing promoters derived from the genome of mammalian cells (*e.g.*, metallothionein promoter) or from mammalian viruses (*e.g.*, the adenovirus late promoter; the vaccinia virus 7.5K promoter). Preferably, bacterial cells such as *E. coli*, and more preferably, eukaryotic cells, especially for the expression of whole recombinant EphA2 antigenic peptide, are used for the expression of a recombinant EphA2 antigenic peptide. For example, mammalian cells such as Chinese hamster ovary cells (CHO), in conjunction with a vector such as the major intermediate early gene promoter element from human cytomegalovirus is an effective expression system for peptides (Foecking *et al.*, 1986, *Gene* 45:101; and Cockett *et al.*, 1990, *BioTechnology* 8:2). In a specific embodiment, the expression of nucleotide sequences encoding an EphA2 antigenic peptide of the invention is regulated by a constitutive promoter, inducible promoter or tissue specific promoter.

**[00181]** In bacterial systems, a number of expression vectors may be advantageously selected depending upon the use intended for the EphA2 antigenic peptide being expressed. For example, when a large quantity of such a protein is to be produced, for the generation of an EphA2 antigenic peptide vaccine, vectors which direct the expression of high levels of protein products that are readily purified may be desirable. Such vectors include, but are not limited to, the *E. coli* expression vector pUR278 (Ruther *et al.*, 1983, *EMBO* 12:1791), in which the peptide coding sequence may be ligated individually into the vector in frame with the lac Z coding region so that a fusion protein is produced; pIN vectors (Inouye & Inouye, 1985, *Nucleic Acids Res.* 13:3101-3109; Van Heeke & Schuster, 1989, *J. Biol. Chem.* 264:5503-5509); and the like. pGEX vectors may also be used to express foreign polypeptides as fusion proteins with glutathione S-transferase (GST). In general, such fusion proteins are soluble and can easily be purified from lysed cells by adsorption and binding to matrix glutathione-agarose beads followed by elution in the presence of free

glutathione. The pGEX vectors are designed to include thrombin or factor Xa protease cleavage sites so that the cloned target gene product can be released from the GST moiety.

[00182] In an insect system, Autographa californica nuclear polyhedrosis virus (AcNPV) is used as a vector to express foreign genes. The virus grows in *Spodoptera frugiperda* cells. The peptide coding sequence may be cloned individually into non-essential regions (e.g., the polyhedrin gene) of the virus and placed under control of an AcNPV promoter (e.g., the polyhedrin promoter).

[00183] In mammalian host cells, a number of viral-based expression systems may be utilized. In cases where an adenovirus is used as an expression vector, the peptide coding sequence of interest may be ligated to an adenovirus transcription/translation control complex, e.g., the late promoter and tripartite leader sequence. This chimeric gene may then be inserted in the adenovirus genome by *in vitro* or *in vivo* recombination. Insertion in a non-essential region of the viral genome (e.g., region EI or E3) will result in a recombinant virus that is viable and capable of expressing the EphA2 antigenic peptide in infected hosts (e.g., see Logan & Shenk, 1984, *PNAS* 81:355-359). Specific initiation signals may also be required for efficient translation of inserted peptide coding sequences. These signals include the ATG initiation codon and adjacent sequences. Furthermore, the initiation codon must be in phase with the reading frame of the desired coding sequence to ensure translation of the entire insert. These exogenous translational control signals and initiation codons can be of a variety of origins, both natural and synthetic. The efficiency of expression may be enhanced by the inclusion of appropriate transcription enhancer elements, transcription terminators, etc. (see, e.g., Bittner *et al.*, 1987, *Methods in Enzymol.* 153:516-544).

[00184] In addition, a host cell strain may be chosen which modulates the expression of the inserted sequences, or modifies and processes the gene product in the specific fashion desired. Such modifications (e.g., glycosylation) and processing (e.g., cleavage) of protein products may be important for the function of the protein. Different host cells have characteristic and specific mechanisms for the post-translational processing and modification of proteins and gene products. Appropriate cell lines or host systems can be chosen to ensure the correct modification and processing of the foreign protein expressed. To this end, eukaryotic host cells which possess the cellular machinery for proper processing of the primary transcript, glycosylation, and phosphorylation of the gene product may be used. Such mammalian host cells include but are not limited to CHO, VERO, BHK, HeLa, COS, MDCK, 293, 3T3, W138, BT483, Hs578T, HTB2, BT20, NS1 and

T47D, NS0 (a murine myeloma cell line that does not endogenously produce any peptide chains), CRL7030 and HsS78Bst cells.

**[00185]** For long-term, high-yield production of recombinant proteins, stable expression is preferred. For example, cell lines which stably express the EphA2 antigenic peptide may be engineered. Rather than using expression vectors which contain viral origins of replication, host cells can be transformed with DNA controlled by appropriate expression control elements (e.g., promoter, enhancer, sequences, transcription terminators, polyadenylation sites, etc.), and a selectable marker. Following the introduction of the foreign DNA, engineered cells may be allowed to grow for 1-2 days in an enriched media, and then are switched to a selective media. The selectable marker in the recombinant plasmid confers resistance to the selection and allows cells to stably integrate the plasmid into their chromosomes and grow to form foci which in turn can be cloned and expanded into cell lines. This method may advantageously be used to engineer cell lines which express the EphA2 antigenic peptide. Such engineered cell lines may be particularly useful in screening and evaluation of compositions that interact directly or indirectly with the EphA2 antigenic peptide.

**[00186]** A number of selection systems may be used, including but not limited to, the herpes simplex virus thymidine kinase (Wigler *et al.*, 1977, *Cell* 11:223), glutamine synthetase, hypoxanthine guanine phosphoribosyltransferase (Szybalski & Szybalski, 1992, *Proc. Natl. Acad. Sci. USA* 48:202), and adenine phosphoribosyltransferase (Lowy *et al.*, 1980, *Cell* 22:8-17) genes can be employed in tk-, gs-, hgprt- or aprt- cells, respectively. Also, antimetabolite resistance can be used as the basis of selection for the following genes: *dhfr*, which confers resistance to methotrexate (Wigler *et al.*, 1980, *PNAS* 77:357; O'Hare *et al.*, 1981, *PNAS* 78:1527); *gpt*, which confers resistance to mycophenolic acid (Mulligan & Berg, 1981, *PNAS* 78:2072); neo, which confers resistance to the aminoglycoside G-418 (Wu and Wu, 1991, *Biotherapy* 3:87; Tolstoshev, 1993, *Ann. Rev. Pharmacol. Toxicol.* 32:573; Mulligan, 1993, *Science* 260:926; and Morgan and Anderson, 1993, *Ann. Rev. Biochem.* 62: 191; May, 1993, *TIB TECH* 11:155-); and *hygro*, which confers resistance to hygromycin (Santerre *et al.*, 1984, *Gene* 30:147). Methods commonly known in the art of recombinant DNA technology may be routinely applied to select the desired recombinant clone, and such methods are described, for example, in Ausubel *et al.* (eds.), *Current Protocols in Molecular Biology*, John Wiley & Sons, NY (1993); Krieger, *Gene Transfer and Expression, A Laboratory Manual*, Stockton Press, NY (1990); and in Chapters 12 and 13, Dracopoli *et al.* (eds.), *Current Protocols in Human Genetics*, John Wiley & Sons, NY

(1994); Colberre-Garapin *et al.*, 1981, *J. Mol. Biol.* 150:1, which are incorporated by reference herein in their entireties.

**[00187]** The expression levels of an EphA2 antigenic peptide can be increased by vector amplification (for a review, see Bebbington and Hentschel, The use of vectors based on gene amplification for the expression of cloned genes in mammalian cells in DNA cloning, Vol.3. (Academic Press, New York, 1987)). When a marker in the vector system expressing peptide is amplifiable, increase in the level of inhibitor present in culture of host cell will increase the number of copies of the marker gene. Since the amplified region is associated with the peptide gene, production of the peptide will also increase (Crouse *et al.*, 1983, *Mol. Cell. Biol.* 3:257).

**[00188]** Once an EphA2 antigenic peptide of the invention has been produced by recombinant expression, it may be purified by any method known in the art for purification of an EphA2 antigenic peptide, for example, by chromatography (*e.g.*, ion exchange, affinity, particularly by affinity for the specific antigen after Protein A, and sizing column chromatography), centrifugation, differential solubility, or by any other standard technique for the purification of proteins. Further, the EphA2 peptides of the present invention or fragments thereof may be fused to heterologous polypeptide sequences described herein or otherwise known in the art to facilitate purification.

## **5.6 Gene Therapy**

**[00189]** As discussed above, the present invention encompasses compositions and methods employ an EphA2 antigenic peptide expression vehicles. In certain embodiments, the expression vehicle is any gene therapy vector available in the art can be used. Exemplary gene therapy vectors that may be used as EphA2 antigenic peptide expression vehicles are described below.

**[00190]** For general reviews of the methods of gene therapy, *see*, Goldspiel *et al.*, 1993, *Clinical Pharmacy* 12:488-505; Wu and Wu, 1991, *Biotherapy* 3:87-95; Tolstoshev, 1993, *Ann. Rev. Pharmacol. Toxicol.* 32:573-596; Mulligan, 1993, *Science* 260:926-932; Morgan and Anderson, 1993, *Ann. Rev. Biochem.* 62:191-217; May, 1993, *TIBTECH* 1, 1(5):155-215. Methods commonly known in the art of recombinant DNA technology which can be used are described in Ausubel *et al.* (eds.), *Current Protocols in Molecular Biology*, John Wiley & Sons, NY (1993); and Kriegler, *Gene Transfer and Expression, A Laboratory Manual*, Stockton Press, NY (1990).

**[00191]** In a preferred aspect, the expression vehicle comprises nucleic acid sequences encoding an EphA2 antigenic peptide, said nucleic acid sequences being part of expression vectors that express the EphA2 antigenic peptide in a suitable host. In

particular, such nucleic acid sequences have promoters operably linked to the EphA2 antigenic peptide, said promoter being inducible or constitutive, and, optionally, tissue-specific. In another particular embodiment, nucleic acid molecules are used in which the EphA2 antigenic peptide coding sequences and any other desired sequences are flanked by regions that promote homologous recombination at a desired site in the genome, thus providing for intrachromosomal expression of the EphA2 antigenic peptide (Koller and Smithies, 1989, Proc. Natl. Acad. Sci. USA 86:8932-8935; Zijlstra *et al.*, 1989, Nature 342:435-438).

**[00192]** Delivery of the nucleic acids into a subject may be either direct, in which case the patient is directly exposed to the nucleic acid or nucleic acid-carrying vectors, or indirect, in which case, cells are first transformed with the nucleic acids *in vitro*, then transplanted into the patient. These two approaches are known, respectively, as *in vivo* or *ex vivo* gene therapy.

**[00193]** In a specific embodiment, the nucleic acid sequences are directly administered *in vivo*, where it is expressed to produce the encoded EphA2 antigenic peptide. This can be accomplished by any of numerous methods known in the art, for example by constructing them as part of an appropriate nucleic acid expression vector and administering the vector so that the nucleic acid sequences become intracellular. Gene therapy vectors can be administered by infection using defective or attenuated retrovirals or other viral vectors (*see, e.g.*, U.S. Patent No. 4,980,286); direct injection of naked DNA; use of microparticle bombardment (*e.g.*, a gene gun; Biolistic, Dupont); coating with lipids or cell-surface receptors or transfecting agents; encapsulation in liposomes, microparticles, or microcapsules; administration in linkage to a peptide which is known to enter the nucleus; administration in linkage to a ligand subject to receptor-mediated endocytosis (*see, e.g.*, Wu and Wu, 1987, J. Biol. Chem. 262:4429-4432) (which can be used to target cell types specifically expressing the receptors); *etc.* In another embodiment, nucleic acid-ligand complexes can be formed in which the ligand comprises a fusogenic viral peptide to disrupt endosomes, allowing the nucleic acid to avoid lysosomal degradation. In yet another embodiment, the nucleic acid can be targeted *in vivo* for cell specific uptake and expression, by targeting a specific receptor (*see, e.g.*, PCT Publications WO 92/06 180; WO 92/22635; W092/20316; W093/14188, and WO 93/20221). Alternatively, the nucleic acid can be introduced intracellularly and incorporated within host cell DNA for expression by homologous recombination (Koller and Smithies, 1989, Proc. Natl. Acad. Sci. USA 86:8932-8935; Zijlstra *et al.*, 1989, Nature 342:435-438).

[00194] In a specific embodiment, viral vectors that contain nucleic acid sequences encoding an EphA2 antigenic peptide. For example, a retroviral vector can be used (see Miller *et al.*, 1993, Meth. Enzymol. 217:581-599). These retroviral vectors contain the components necessary for the correct packaging of the viral genome and integration into the host cell DNA. The nucleic acid sequences encoding the EphA2 antigenic peptide to be used in gene therapy are cloned into one or more vectors, thereby facilitating delivery of the gene into a patient. More detail about retroviral vectors can be found in Boesen *et al.*, 1994, Biotherapy 6:291-302, which describes the use of a retroviral vector to deliver the *mdr 1* gene to hematopoietic stem cells in order to make the stem cells more resistant to chemotherapy. Other references illustrating the use of retroviral vectors in gene therapy are: Clowes *et al.*, 1994, J. Clin. Invest. 93:644-651; Klein *et al.*, 1994, Blood 83:1467-1473; Salmons and Gunzberg, 1993, Human Gene Therapy 4:129-141; and Grossman and Wilson, 1993, Curr. Opin. in Genetics and Devel. 3:110-114.

[00195] One approach to gene therapy encompassed by the present methods and compositions involves transferring a gene, *e.g.*, a nucleic acid encoding an EphA2 antigenic peptide, to cells in tissue culture by such methods as electroporation, lipofection, calcium phosphate mediated transfection, or viral infection. Usually, the method of transfer includes the transfer of a selectable marker to the cells. The cells are then placed under selection to isolate those cells that have taken up and are expressing the transferred gene. Those cells are then delivered to a subject.

[00196] In this embodiment, the nucleic acid is introduced into a cell prior to administration *in vivo* of the resulting recombinant cell. Such introduction can be carried out by any method known in the art, including but not limited to transfection, electroporation, microinjection, infection with a viral or bacteriophage vector containing the nucleic acid sequences, cell fusion, chromosome-mediated gene transfer, microcell mediated gene transfer, spheroplast fusion, *etc.* Numerous techniques are known in the art for the introduction of foreign genes into cells (see, *e.g.*, Loeffler and Behr, 1993, Meth. Enzymol. 217:599-618; Cohen *et al.*, 1993, Meth. Enzymol. 217:618-644; Cline, 1985, Pharmac. Ther. 29:69-92) and may be used in accordance with the present invention, provided that the necessary developmental and physiological functions of the recipient cells are not disrupted. The technique should provide for the stable transfer of the nucleic acid to the cell, so that the nucleic acid is expressible by the cell and preferably heritable and expressible by its cell progeny.

[00197] The resulting recombinant cells can be delivered to a patient by various methods known in the art. Recombinant blood cells (*e.g.*, hematopoietic stem or progenitor

cells) are preferably administered intravenously. The amount of cells envisioned for use depends on the desired effect, patient state, *etc.*, and can be determined by one skilled in the art.

[00198] Cells into which a nucleic acid can be introduced for purposes of gene therapy encompass any desired, available cell type, and include but are not limited to fibroblasts; blood cells such as T lymphocytes, B lymphocytes, monocytes, macrophages, neutrophils, eosinophils, megakaryocytes, granulocytes; various stem or progenitor cells, in particular hematopoietic stem or progenitor cells, *e.g.*, as obtained from bone marrow, umbilical cord blood, peripheral blood, fetal liver, *etc.*

[00199] In a preferred embodiment, the cell used for gene therapy is autologous to the patient.

[00200] In an embodiment in which recombinant cells are used in gene therapy, nucleic acid sequences encoding EphA2 antigenic peptide are introduced into the cells such that they are expressible by the cells or their progeny, and the recombinant cells are then administered *in vivo* for therapeutic effect. In a specific embodiment, stem or progenitor cells are used. Any stem and/or progenitor cells which can be isolated and maintained *in vitro* can potentially be used in accordance with this embodiment of the present invention (see *e.g.* PCT Publication WO 94/08598; Stemple and Anderson, 1992, Cell 71:973-985; Rheinwald, 1980, Meth. Cell Bio. 21A:229; and Pittelkow and Scott, 1986, Mayo Clinic Proc. 61:771).

[00201] In a specific embodiment, the nucleic acid to be introduced for purposes of gene therapy comprises an inducible promoter operably linked to the coding region, such that expression of the nucleic acid is controllable by controlling the presence or absence of the appropriate inducer of transcription.

## 5.7 Bacterial Expression Vehicles

[00202] In certain embodiments, the present invention provides EphA2 antigenic peptide expression vehicles in the form of a microorganism, and, in specific embodiments, the microorganism is a bacterium.

[00203] Microorganisms useful for the methods of the present invention include but are not limited to *Borrelia burgdorferi*, *Brucella melitensis*, *Escherichia coli*, enteroinvasive *Escherichia coli*, *Legionella pneumophila*, *Salmonella typhi*, *Salmonella typhimurium*, *Shigella spp.*, *Streptococcus spp.*, *Treponema pallidum*, *Yersinia enterocolitica*, *Listeria monocytogenes*, *Mycobacterium avium*, *Mycobacterium bovis*, *Mycobacterium tuberculosis*, BCG, *Mycoplasma hominis*, *Rickettsiae quintana*, *Cryptococcus neoformans*, *Histoplasma capsulatum*, *Pneumocystis carinii*, *Eimeria*



*acervulina*, *Neospora caninum*, *Plasmodium falciparum*, *Sarcocystis suihominis*, *Toxoplasma gondii*, *Leishmania amazonensis*, *Leishmania major*, *Leishmania mexicana*, *Leptomonas karyophilus*, *Phytomonas* spp., *Trypanosoma cruzi*, *Encephalitozoon cuniculi*, *Nosema helminthorum*, *Unikaryon legeri*.

- 5 [00204] Many of the microorganisms encompassed by the present invention are causative agents of diseases in humans and animals. For example, sepsis from gram negative bacteria is a serious problem because of the high mortality rate associated with the onset of septic shock (R.C. Bone, 1993, Clinical Microbiol. Revs. 6:57-68). Therefore, to allow the safe use of these microorganisms in both diagnostics and treatment of humans and
- 10 animals, the microorganisms are attenuated in their virulence for causing disease. The end result is to reduce the risk of toxic shock or other side effects due to administration of the vector to the patient. Such attenuated microorganisms can be isolated by a number of techniques. Such methods include use of antibiotic-sensitive strains of microorganisms, mutagenesis of the microorganisms, selection for microorganism mutants that lack
- 15 virulence factors, and construction of new strains of microorganisms with altered cell wall lipopolysaccharides.

- [00205] In certain embodiments, the microorganisms can be attenuated by the deletion or disruption of DNA sequences which encode for virulence factors which insure survival of the microorganisms in the host cell, especially macrophages and neutrophils, by,
- 20 for example, homologous recombination techniques and chemical or transposon mutagenesis. Many, but not all, of these studied virulence factors are associated with survival in macrophages such that these factors are specifically expressed within macrophages due to stress, for example, acidification, or are used to induced specific host cell responses, for example, macropinocytosis, Fields et al., 1986, Proc. Natl. Acad. Sci. USA 83:5189-5193. Bacterial virulence factors include, for example: cytolysin; defensin
- 25 resistance loci; DNA K; fimbriae; GroEL; inv loci; lipoprotein; LPS; lysosomal fusion inhibition; macrophage survival loci; oxidative stress response loci; pho loci (e.g., PhoP and PhoQ); pho activated genes (pag; e.g., pagB and pagC); phoP and phoQ regulated genes (prg); porins; serum resistance peptide; virulence plasmids (such as spvB, traT and ty2).

- 30 [00206] Yet another method for the attenuation of the microorganisms is to modify substituents of the microorganism which are responsible for the toxicity of that microorganism. For example, lipopolysaccharide (LPS) or endotoxin is primarily responsible for the pathological effects of bacterial sepsis. The component of LPS which results in this response is lipid A (LA). Elimination or mitigation of the toxic effects of LA
- 35 results in an attenuated bacteria since 1) the risk of septic shock in the patient would be

reduced and 2) higher levels of the bacterial EphA2 antigenic peptide expression vehicle could be tolerated.

[00207] *Rhodobacter* (*Rhodopseudomonas*) *sphaeroides* and *Rhodobacter capsulatus* each possess a monophosphoryl lipid A (MLA) which does not elicit a septic shock response in experimental animals and, further, is an endotoxin antagonist. Loppnow et al., 1990, Infect. Immun. 58:3743-3750; Takayma et al., 1989, Infect. Immun. 57:1336-1338. Gram negative bacteria other than *Rhodobacter* can be genetically altered to produce MLA, thereby reducing its potential of inducing septic shock.

[00208] Yet another example for altering the LPS of bacteria involves the introduction of mutations in the LPS biosynthetic pathway. Several enzymatic steps in LPS biosynthesis and the genetic loci controlling them in a number of bacteria have been identified, and several mutant bacterial strains have been isolated with genetic and enzymatic lesions in the LPS pathway. In certain embodiments, the LPS pathway mutant is a *firA* mutant. *firA* is the gene that encodes the enzyme UDP-3-O-(R-30 hydroxymyristoyl)-glycocytamine N-acyltransferase, which regulates the third step in endotoxin biosynthesis (Kelley et al., 1993, J. Biol. Chem. 268:19866-19874).

[00209] As a method of insuring the attenuated phenotype and to avoid reversion to the non-attenuated phenotype, the bacteria may be engineered such that it is attenuated in more than one manner, e.g., a mutation in the pathway for lipid A production and one or more mutations to auxotrophy for one or more nutrients or metabolites, such as uracil biosynthesis, purine biosynthesis, and arginine biosynthesis.

[00210] In certain embodiments of the present invention, the bacterial EphA2 antigenic peptide expression vehicles are engineered to deliver suicide genes to the target EphA2-expressing cells. These suicide genes include pro-drug converting enzymes, such as Herpes simplex thymidine kinase (TK) and bacterial cytosine deaminase (CD). TK phosphorylates the non-toxic substrates acyclovir and ganciclovir, rendering them toxic via their incorporation into genomic DNA. CD converts the non-toxic 5-fluorocytosine (5-FC) into 5-fluorouracil (5-FU), which is toxic via its incorporation into RNA. Additional examples of pro-drug converting enzymes encompassed by the present invention include cytochrome p450 NADPH oxidoreductase which acts upon mitomycin C and porfiromycin (Murray et al., 1994, J. Pharmacol. Exp. Therapeut. 270:645-649). Other exemplary pro-drug converting enzymes that may be used in the methods and compositions of the present invention include: carboxypeptidase; beta-glucuronidase; penicillin-V-amidase; penicillin-G-amidase; beta-lactamase; beta.-glucosidase; nitroreductase; and carboxypeptidase A.

[00211] Where the EphA2 vaccine comprises a microorganism that expresses an EphA2 antigenic peptide and, optionally, a pro-drug converting enzyme, the expression constructs are preferably designed such that the microorganism-produced peptides and enzymes are secreted by the microorganism. A number of bacterial secretion signals are

well known in the art and may be used in the compositions and methods of the present invention. Exemplary secretion signals that can be used with gram-positive microorganisms include SecA (Sadaie et al., Gene 98:101-105, 1991), SecY (Suh et al., Mol. Microbiol. 4:305-314, 1990), SecE (Jeong et al., Mol. Microbiol. 10:133-142, 1993), FtsY an FfH (PCT/NL 96/00278), and PrsA (WO 94/19471). Exemplary secretion signals that may be used with gram-negative microorganisms include those of soluble cytoplasmic proteins such as SecB and heat shock proteins; that of the peripheral membrane-associated protein SecA; and those of the integral membrane proteins SecY, SecE, SecD and SecF.

[00212] The promoters driving the expression of the EphA2 antigenic peptides and, optionally, pro-drug converting enzymes, may be either constitutive, in which the peptides or enzymes are continually expressed, inducible, in which the peptides or enzymes are expressed only upon the presence of an inducer molecule(s), or cell-type specific control, in which the peptides or enzymes are expressed only in certain cell types. For example, a suitable inducible promoter can be a promoter responsible for the bacterial "SOS" response (Friedberg et al., In: DNA Repair and Mutagenesis, pp. 407-455, Am. Soc. Microbiol. Press, 1995). Such a promoter is inducible by numerous agents including chemotherapeutic alkylating agents such as mitomycin (Oda et al., 1985, Mutation Research 147:219-229; Nakamura et al., 1987, Mutation Res. 192:239-246; Shimda et al., 1994, Carcinogenesis 15:2523-2529) which is approved for use in humans. Promoter elements which belong to this group include umuC, *suA* and others (Shinagawa et al., 1983, Gene 23:167-174; Schnarr et al., 1991, Biochemie 73:423-431). The *suA* promoter includes the ATG of the *suA* gene and the following 27 nucleotides as well as 70 nucleotides upstream of the ATG (Cole, 1983, Mol. Gen. Genet. 189:400-404). Therefore, it is useful both in expressing foreign genes and in creating gene fusions for sequences lacking initiating codons.

#### 5.7.1 Exemplary Embodiment: *Listeria monocytogenes* As An Expression Vehicle

[00213] *Listeria monocytogenes* (*Listeria*) is a Gram-positive facultative intracellular bacterium that is being developed for use in antigen-specific vaccines due to its ability to prime a potent CD4+/CD8+ T-cell mediated response via both MHC class I and class II antigen presentation pathways, and as such it has been tested recently as a vaccine vector in a human clinical trial among normal healthy volunteers.

**[00214]** *Listeria* has been studied for many years as a model for stimulating both innate and adaptive T cell-dependent antibacterial immunity. The ability of *Listeria* to effectively stimulate cellular immunity is based on its intracellular lifecycle. Upon infecting the host, the bacterium is rapidly taken up by phagocytes including macrophages and dendritic cells into a phagolysosomal compartment. The majority of the bacteria are subsequently degraded. Peptides resulting from proteolytic degradation of pathogens within phagosomes of infected APCs are loaded directly onto MHC class II molecules, and these MHC II-peptide complexes activate CD4+ "helper" T cells that stimulate the production of antibodies, and the processed antigens are expressed on the surface of the antigen presenting cell via the class II endosomal pathway. Within the acidic compartment, certain bacterial genes are activated including the cholesterol-dependent cytolysin, LLO, which can degrade the phagolysosome, releasing the bacterium into the cytosolic compartment of the host cell, where the surviving *Listeria* propagate. Efficient presentation of heterologous antigens via the MHC class I pathway requires de novo endogenous protein expression by *Listeria*. Within antigen presenting cells (APC), proteins synthesized and secreted by *Listeria* are sampled and degraded by the proteasome. The resulting peptides are shuttled into the endoplasmic reticulum by TAP proteins and loaded onto MHC class I molecules. The MHC I-peptide complex is delivered to the cell surface, which in combination with sufficient co-stimulation (signal 2) activates and stimulates cytotoxic T lymphocytes (CTLs) having the cognate T cell receptor to expand and subsequently recognize the MHC I-peptide complex.

**[00215]** The EphA2 antigenic peptides are preferably expressed in *Listeria* using a heterologous gene expression cassette. A heterologous gene expression cassette is typically comprised of the following ordered elements: (1) prokaryotic promoter; (2) Shine-Dalgarno sequence; (3) secretion signal (signal peptide); and, (4) heterologous gene. Optionally, the heterologous gene expression cassette may also contain a transcription termination sequence, in constructs for stable integration within the bacterial chromosome. While not required, inclusion of a transcription termination sequence as the final ordered element in a heterologous gene expression cassette may prevent polar effects on the regulation of expression of adjacent genes, due to read-through transcription.

**[00216]** The expression vectors introduced into the *Listeria*-based EphA2 vaccine are preferably designed such that the *Listeria*-produced EphA2 peptides and, optionally, prodrug converting enzymes, are secreted by the *Listeria*. A number of bacterial secretion signals are well known in the art and may be used in the compositions and methods of the present invention. Exemplary secretion signals that can be used with gram-positive

microorganisms include SecA (Sadaie *et al.*, 1991, *Gene* 98:101-105), SecY (Suh *et al.*, 1990, *Mol. Microbiol.* 4:305-314), SecE (Jeong *et al.*, 1993, *Mol. Microbiol.* 10:133-142), FtsY and Ffh (PCT/NL 96/00278), and PrsA (WO 94/19471).

[00217] The promoters driving the expression of the EphA2 antigenic peptides and, optionally, pro-drug converting enzymes, may be either constitutive, in which the peptides or enzymes are continually expressed, inducible, in which the peptides or enzymes are expressed only upon the presence of an inducer molecule(s), or cell-type specific control, in which the peptides or enzymes are expressed only in certain cell types. For example, a suitable inducible promoter can be a promoter responsible for the bacterial "SOS" response (Friedberg *et al.*, In: DNA Repair and Mutagenesis, pp. 407-455, Am. Soc. Microbiol. Press, 1995). Such a promoter is inducible by numerous agents including chemotherapeutic alkylating agents such as mitomycin (Oda *et al.*, 1985, *Mutation Research* 147:219-229; Nakamura *et al.*, 1987, *Mutation Res.* 192:239-246; Shimda *et al.*, 1994, *Carcinogenesis* 15:2523-2529) which is approved for use in humans. Promoter elements which belong to this group include umuC, sulA and others (Shinagawa *et al.*, 1983, *Gene* 23:167-174; Schnarr *et al.*, 1991, *Biochemie* 73:423-431). The sulA promoter includes the ATG of the sulA gene and the following 27 nucleotides as well as 70 nucleotides upstream of the ATG (Cole, 1983, *Mol. Gen. Genet.* 189:400-404). Therefore, it is useful both in expressing foreign genes and in creating gene fusions for sequences lacking initiating codons.

[00218] Preferred embodiments of components of the EphA2 antigenic peptide expression system, to be used in conjunction with nucleic acids encoding EphA2 antigenic peptides described in Section 5.2, are provided below.

#### 5.7.1.1. Construct Backbone

[00219] One of ordinary skill in the art will recognize that a variety of plasmid construct backbones are available which are suitable for use in the assembly of a heterologous gene expression cassette. A particular plasmid construct backbone is selected based on whether expression of the heterologous gene from the bacterial chromosome or from an extra-chromosomal episome is desired.

[00220] Given as non-limiting examples, incorporation of the heterologous gene expression cassette into the bacterial chromosome of *Listeria monocytogenes* (*Listeria*) is accomplished with an integration vector that contains an expression cassette for a listeriophage integrase that catalyzes sequence-specific integration of the vector into the *Listeria* chromosome. For example, the integration vectors known as pPL1 and pPL2 program stable single-copy integration of a heterologous protein (e.g., EphA2-antigenic peptide) expression cassette within an innocuous region of the bacterial genome, and have

been described in the literature (Lauer et. al.2002 J. Bacteriol. 184:4177-4178). The integration vectors are stable as plasmids in *E. coli* and are introduced via conjugation into the desired *Listeria* background. Each vector lacks a *Listeria*-specific origin of replication and encodes a phage integrase, such that the vectors are stable only upon integration into a chromosomal phage attachment site. Starting with a desired plasmid construct, the process of generating a recombinant *Listeria* strain expressing a desired protein(s) takes approximately one week. The pPL1 and pPL2 integration vectors are based, respectively, on the U153 and PSA listeriphages. The pPL1 vector integrates within the open reading frame of the *comK* gene, while pPL2 integrates within the *tRNAArg* gene in such a manner that the native sequence of the gene is restored upon successful integration, thus keeping its native expressed function intact. The pPL1 and pPL2 integration vectors contain a multiple cloning site sequence in order to facilitate construction of plasmids containing the heterologous protein (e.g., EphA2-antigenic peptide) expression cassette. Alternatively, incorporation of the EphA2-antigenic peptide expression cassette into the *Listeria* chromosome can be accomplished through allelic exchange methods, known to those skilled in the art. In particular, compositions in which it is desired to not incorporate a gene encoding an antibiotic resistance protein as part of the construct containing the heterologous gene expression cassette, methods of allelic exchange are desirable. For example, the pKSV7 vector (Camilli et. al. Mol. Microbiol. 1993 8,143-157), contains a temperature-sensitive *Listeria* Gram-positive replication origin which is exploited to select for recombinant clones at the non-permissive temperature that represent the pKSV7 plasmid recombined into the *Listeria* chromosome. The pKSV7 allelic exchange plasmid vector contains a multiple cloning site sequence in order to facilitate construction of plasmids containing the heterologous protein (e.g., EphA2-antigenic peptide) expression cassette, and also a chloramphenicol resistance gene. For insertion into the *Listeria* chromosome, the heterologous EphA2-antigenic peptide expression cassette construct is optimally flanked by approximately 1 kb of chromosomal DNA sequence that corresponds to the precise location of desired integration. The pKSV7-heterologous protein (e.g., EphA2-antigenic peptide) expression cassette plasmid is introduced optimally into a desired bacterial strain by electroporation, according to standard methods for electroporation of Gram positive bacteria. Briefly, bacteria electroporated with the pKSV7-heterologous protein (e.g., EphA2-antigenic peptide) expression cassette plasmid are selected by plating on BHI agar media containing chloramphenicol (10 µg/ml), and incubated at the permissive temperature of 30°C. Single cross-over integration into the bacterial chromosome is selected by passing several individual colonies for multiple generations at the non-permissive

temperature of 41°C in media containing chloramphenicol. Finally, plasmid excision and curing (double cross-over) is achieved by passaging several individual colonies for multiple generations at the permissive temperature of 30°C in BHI media not containing chloramphenicol. Verification of integration of the heterologous protein (e.g., EphA2-antigenic peptide) expression cassette into the bacteria chromosome can be accomplished by PCR, utilizing a primer pair that amplifies a region defined from within the heterologous protein (e.g., EphA2-antigenic peptide) expression cassette to the bacterial chromosome targeting sequence not contained in the pKSV7 plasmid vector construct.

[00221] In other compositions, it may be desired to express the heterologous protein (e.g., EphA2-antigenic peptide) from a stable plasmid episome. Maintenance of the plasmid episome through passaging for multiple generations requires the co-expression of a protein that confers a selective advantage for the plasmid-containing bacterium. As non-limiting examples, the protein co-expressed from the plasmid in combination with the heterologous protein (e.g., EphA2-antigenic peptide) may be an antibiotic resistance protein, for example chloramphenicol, or may be a bacterial protein (that is expressed from the chromosome in wild-type bacteria), that can also confer a selective advantage. Non-limiting examples of bacterial proteins include enzyme required for purine or amino acid biosynthesis (selection under defined media lacking relevant amino acids or other necessary precursor macromolecules), or a transcription factor required for the expression of genes that confer a selective advantage in vitro or in vivo (Gunn et. al. 2001 J. Immunol. 167:6471-6479). As a non-limiting example, pAM401 is a suitable plasmid for episomal expression of a selected heterologous protein (e.g., EphA2-antigenic peptide) in diverse Gram-positive bacterial genera (Wirth et. al. 1986 J. Bacteriol 165:831-836).

#### 5.7.1.2. Shine-Dalgarno Sequence

[00222] At the 3' end of the promoter is contained a poly-purine Shine-Dalgarno sequence, the element required for engagement of the 30S ribosomal subunit (via 16S rRNA) to the heterologous gene RNA transcript and initiation of translation. The Shine-Dalgarno sequence has typically the following consensus sequence: (SEQ ID NO:66): 5'-NAGGAGGU-N5-10-AUG (start codon)-3'. There are variations of the poly-purine Shine-Dalgarno sequence. Notably, the *Listeria* hly gene that encodes listerolysin O (LLO) has the following Shine-Dalgarno sequence (SEQ ID NO:67): AAGGAGAGTGAAACCCATG (Shine-Dalgarno sequence is underlined, and the translation start codon is bolded).

#### 5.7.1.3. Codon Optimization

[00223] It is well known to those skilled in the art that for optimal translation efficiency of a selected heterologous protein, it is desirable to utilize codons favored by the bacterium. The preferred codon usage for bacterial expression can be found at the following link: <http://www.kazusa.or.jp/codon/>. In some embodiments, expression of heterologous proteins from *Listeria monocytogenes* is desired. The preferred *Listeria* monocytogenes codon usage can be found at the following link:

[http://www.kazusa.or.jp/codon/cgi-bin/showcodon.cgi?species=Listeria+monocytogenes+\[gbtct\]](http://www.kazusa.or.jp/codon/cgi-bin/showcodon.cgi?species=Listeria+monocytogenes+[gbtct])

[00224] The optimal codons utilized by *Listeria monocytogenes* for each amino acid are shown in the table below.

**Listeria Codon Bias: Codons to be used for optimizing expression**

Amino Acid	One Letter Code	Optimal <i>Listeria</i> Codon
Alanine	A	GCA
Arginine	R	CGU
Asparagine	N	AAU
Aspartate	D	GAU
Cysteine	C	UGU
Glutamine	Q	CAA
Glutamate	E	GAA
Glycine	G	GGU
Histidine	H	CAU
Isoleucine	I	AUU
Leucine	L	UUA
Lysine	K	AAA
Methionine	M	AUG
Phenylalanine	F	UUU
Proline	P	CCA
Serine	S	AGU
Threonine	T	ACA
Tryptophan	W	UGG
Tyrosine	Y	UAU
Valine	V	GUU

#### 5.7.1.4. Signal Peptides

[00225] Bacteria utilize diverse pathways for protein secretion, including secA1 and Twin-Arg Translocation (Tat), which are located at the N-terminal end of the pre-protein.

The majority of secreted proteins utilize the Sec pathway, in which the protein translocates through the bacterial membrane-embedded proteinaceous Sec pore in an unfolded



conformation. In contrast, the proteins utilizing the Tat pathway are secreted in a folded conformation. Nucleotide sequence encoding signal peptides corresponding to either of these protein secretion pathways can be fused genetically in-frame to a desired heterologous protein coding sequence. The signal peptides optimally contain a signal peptidase at their carboxyl terminus for release of the authentic desired protein into the extra-cellular environment (Sharkov and Cai. 2002 J. Biol. Chem. 277:5796-5803; Nielsen et. al. 1997 Protein Engineering 10:1-6; and, <http://www.cbs.dtu.dk/services/SignalP/>). The signal peptides can be derived not only from diverse secretion pathways, but also from diverse bacterial genera. Signal peptides have a common structural organization, having a charged N-terminus (N-domain), a hydrophobic core region (H-domain) and a more polar C-terminal region (C-domain), however, they do not show sequence conservation. The C-domain of the signal peptide carries a type I signal peptidase (SPase I) cleavage site, having the consensus sequence A-X-A, at positions -1 and -3 relative to the cleavage site. Proteins secreted via the sec pathway have signal peptides that average 28 residues. Signal peptides related to proteins secreted by the Tat pathway have a tripartite organization similar to Sec signal peptides, but are characterized by having an RR-motif (R-R-X-##, where # is a hydrophobic residue), located at the N-domain / H-domain boundary. Bacterial Tat signal peptides average 14 amino acids longer than sec signal peptides. The *Bacillus subtilis* secretome may contain as many as 69 putative proteins that utilize the Tat secretion pathway, 14 of which contain a SPase I cleavage site (Jongbloed et. al. 2002 J. Biol. Chem. 277:44068-44078; Thalsma et. al., 2000 Microbiol. Mol. Biol. Rev. 64:515-547). Shown in the table below are non-limiting examples of signal peptides that can be used in fusion compositions with a selected heterologous gene, resulting in secretion from the bacterium of the encoded protein.

Secretion Pathway	Signal Peptide Amino Acid Sequence (NH <sub>2</sub> -CO <sub>2</sub> )	Signal peptidase Site (cleavage site represented by *)	Gene	Genus/species	SEQ ID NO:
secA1	MKKIMLVFITLILVSLPI AQTEAKD	TEA*KD (SEQ ID NO:70)	<i>hly</i> (LLO)	<i>Listeria monocytogenes</i>	44
	MTDKKSENQTEKTETK ENKGMTRREMLKLSAV AGTGIAVGATGLGTLIN VVDQVDKALT	DKA*LT (SEQ ID NO:71)	lmo0367	<i>Listeria monocytogenes</i>	45
	MAYDSRFDEWVQKLLK EESFQNNTFDRRKFIQG AGKIAGLSLGLTIAQSV GAFG	VGA*FG (SEQ ID NO:72)	PhoD (alkaline phosphatase)	<i>Bacillus subtilis</i>	46

[00226] It should be appreciated by those skilled in the art that there exists a variety of proteins secreted via the Tat pathway among diverse bacterial genera, and that selected

Tat signal peptides corresponding to these proteins can be fused genetically in-frame to a desired sequence encoding a heterologous protein, to facilitate secretion of the functionally linked Tat signal peptide-heterologous protein chimera via the Tat pathway. Provided below are non-limiting examples of proteins from *Bacillus subtilis* and *Listeria (innocua and monocytogenes)* that are predicted to utilize Tat pathway secretion.

Putative *Bacillus subtilis* Proteins Secreted by Tat  
Pathway<http://www.sas.upenn.edu/~pohlschr/TABLE1.html>

- [00227] >gi|2635523|emb|CAB15017.1| similar to two (component sensor histidine kinase (YtsA) (*Bacillus subtilis*)
- 10 [00228] >gi|2632548|emb|CAB12056.1| phosphodiesterase/alkaline phosphatase D (*Bacillus subtilis*)
- [00229] >gi|2632573|emb|CAB12081.1| similar to hypothetical proteins (*Bacillus subtilis*)
- [00230] >gi|2633776|emb|CAB13278.1| similar to hypothetical proteins (*Bacillus subtilis*)
- 15 [00231] >gi|2634674|emb|CAB14172.1| menaquinol:cytochrome c oxidoreductase (iron (sulfur subunit) (*Bacillus subtilis*)
- [00232] >gi|2635595|emb|CAB15089.1| yubF (*Bacillus subtilis*)
- [00233] >gi|2636361|emb|CAB15852.1| alternate gene name: ipa (29d~similar to
- 20 hypothetical proteins (*Bacillus subtilis*)

Putative *Listeria* Proteins Secreted by Tat  
Pathway<http://www.sas.upenn.edu/~pohlschr/TABLE1.html>

- [00234] *Listeria innocua*<http://www.sas.upenn.edu/~pohlschr/TABLE1.html>
- [00235] >gi|16799463|ref|NP\_469731.1| conserved hypothetical protein similar to B. subtilis YwbN protein (*Listeria innocua*)
- 25 [00236] >gi|16801368|ref|NP\_471636.1| similar to 3 (oxoacyl (acyl (carrier protein synthase (*Listeria innocua*)
- [00237] *Listeria monocytogenes* EGD (e)  
<http://www.sas.upenn.edu/~pohlschr/TABLE1.html>
- 30 [00238] >gi|16802412|ref|NP\_463897.1| conserved hypothetical protein similar to B. subtilis YwbN protein (*Listeria monocytogenes* EGD (e)
- [00239] Organisms utilize codon bias to regulate expression of particular endogenous genes. Thus, signal peptides utilized for secretion of selected heterologous proteins may not contain codons that utilize preferred codons, resulting in non-optimal levels of protein
- 35 synthesis. In some some embodiments, the signal peptide sequence fused in frame with a

gene encoding a selected heterologous protein is codon-optimized for codon usage in a selected bacterium. In some embodiments for expression and secretion from recombinant *Listeria monocytogenes*, a nucleotide sequence of a selected signal peptide is codon optimized for expression in *Listeria monocytogenes*, according to the table (*ibid.*)

#### 5 **5.7.1.5. Transcription Termination Sequence**

[00240] In some embodiments, a transcription termination sequence can be inserted into the heterologous protein expression cassette, downstream from the C-terminus of the translational stop codon related to the heterologous protein. Appropriate sequence elements known to those who are skilled in the art that promote either rho-dependent or rho-independent transcription termination can be placed in the heterologous protein expression cassette.

### **5.8 Anti-Idiotypic Antibodies**

[00241] The present invention relates to methods and compositions utilizing EphA2 vaccines for eliciting immune responses against EphA2-expressing cells and treatment and prevention of disorders involving EphA2-expressing cells. In certain embodiments, the EphA2 vaccines of the invention comprises an anti-idiotypic of an anti-EphA2 antibody.

[00242] The idiotopes on a single antibody molecule are thought to mimic and be the "internal image" of any foreign or self epitope at the molecular level. By means of Mab technology, an antibody against an EphA2 epitope is produced, purified and subsequently used as an immunogen to elicit an anti-idiotypic antibody which may be an internal image of the original EphA2 epitope. Thus, as predicted by the Jerne "network" theory (Jerne, 1974, *Ann. Inst. Pasteur. Immun.* 125C:373-389), immunization with an anti-idiotypic antibody that is directed against antigen combining sites of an anti-EphA2 epitope antibody would elicit a humoral immune response specific for the nominal antigen. The resulting anti-anti-idiotypic antibody should react with the original primary EphA2 epitope.

express EphA2 and thus be useful in treating, preventing or managing hyperproliferative diseases involving EphA2-overexpressing cells.

### **5.9 Prophylactic/Therapeutic Methods**

[00244] The present invention encompasses methods for treating, preventing, or  
5 managing a disease or disorder associated with overexpression of EphA2 and/or cell  
hyperproliferative disorders, preferably cancer, in a subject comprising administering one or  
more EphA2 vaccines of the invention.

[00245] The present invention further encompasses methods for eliciting an immune  
response against an EphA2-expressing cell associated with a hyperproliferative disorder,  
10 comprising administering to a subject one or more EphA2 vaccines of the invention in an  
amount effective for eliciting an immune response against the EphA2-expressing cell.

[00246] An EphA2 vaccine may comprise one more EphA2 antigenic peptides, one  
ore more EphA2 antigenic peptide expression vehicles, or antigen presenting cells  
sensitized with one ore more EphA2 antigenic peptides.

[00247] In another specific embodiment, the disorder to be treated, prevented, or  
15 managed is a pre-cancerous condition associated with cells that overexpress EphA2. In  
more specific embodiments, the pre-cancerous condition is high-grade prostatic  
intraepithelial neoplasia (PIN), fibroadenoma of the breast, fibrocystic disease, or  
compound nevi.

[00248] In one embodiment, the peptides of the invention can be administered in  
20 combination with one or more other therapeutic agents useful in the treatment, prevention  
or management of diseases or disorders associated with EphA2 overexpression,  
hyperproliferative disorders, and/or cancer. In certain embodiments, one or more EphA2  
antigenic peptides of the invention are administered to a mammal, preferably a human,  
25 concurrently with one or more other therapeutic agents useful for the treatment of cancer.  
The term "concurrently" is not limited to the administration of prophylactic or therapeutic  
agents at exactly the same time, but rather it is meant that the EphA2 antigenic peptides of  
the invention and the other agent are administered to a subject in a sequence and within a  
time interval such that the peptides of the invention can act together with the other agent to  
30 provide an increased benefit than if they were administered otherwise. For example, each  
prophylactic or therapeutic agent may be administered at the same time or sequentially in  
any order at different points in time; however, if not administered at the same time, they  
should be administered sufficiently close in time so as to provide the desired therapeutic or  
prophylactic effect. Each therapeutic agent can be administered separately, in any  
35 appropriate form and by any suitable route. In other embodiments, the EphA2 antigenic

peptides of the invention are administered before, concurrently or after surgery. Preferably the surgery completely removes localized tumors or reduces the size of large tumors. Surgery can also be done as a preventive measure or to relieve pain.

[00249] In various embodiments, the prophylactic or therapeutic agents are administered less than 1 hour apart, at about 1 hour apart, at about 1 hour to about 2 hours apart, at about 2 hours to about 3 hours apart, at about 3 hours to about 4 hours apart, at about 4 hours to about 5 hours apart, at about 5 hours to about 6 hours apart, at about 6 hours to about 7 hours apart, at about 7 hours to about 8 hours apart, at about 8 hours to about 9 hours apart, at about 9 hours to about 10 hours apart, at about 10 hours to about 11 hours apart, at about 11 hours to about 12 hours apart, no more than 24 hours apart or no more than 48 hours apart. In preferred embodiments, two or more components are administered within the same patient visit.

[00250] The dosage amounts and frequencies of administration provided herein are encompassed by the terms therapeutically effective and prophylactically effective. The dosage and frequency further will typically vary according to factors specific for each patient depending on the specific therapeutic or prophylactic agents administered, the severity and type of cancer, the route of administration, as well as age, body weight, response, and the past medical history of the patient. Suitable regimens can be selected by one skilled in the art by considering such factors and by following, for example, dosages reported in the literature and recommended in the *Physician's Desk Reference* (56<sup>th</sup> ed., 2002).

### **5.9.1 Patient Population**

[00251] The invention provides methods for treating, preventing, and managing a disease or disorder associated with EphA2 overexpression and/or hyperproliferative cell disease, particularly cancer, by administering to a subject in need thereof one or more EphA2 vaccines of the invention in a therapeutically or prophylactically effective amount or an amount effective to elicit an immune response against EphA2-expressing cells associated with the hyperproliferative disorder. In another embodiment, the EphA2 vaccines of the invention can be administered in combination with one or more other therapeutic or prophylactic agents. The subject is preferably a mammal such as non-primate (e.g., cows, pigs, horses, cats, dogs, rats, etc.) and a primate (e.g., monkey, such as a cynomolgous monkey and a human). In a preferred embodiment, the subject is a human.

[00252] Specific examples of cancers that can be treated by the methods encompassed by the invention include, but are not limited to, cancers that overexpress EphA2. In a further embodiment, the cancer is of an epithelial origin. Examples of such

cancers are cancer of the lung, colon, ovary, esophagus, prostate, breast, and skin. Other cancers include cancer of the bladder and pancreas and renal cell carcinoma and melanoma. In yet a further embodiment, the cancer is of a T cell origin. Examples of such cancers are leukemias and lymphomas. Additional cancers are listed by example and not by limitation in the following section 5.9.1.1. In particular embodiments, methods of the invention can be used to treat and/or prevent metastasis from primary tumors.

**[00253]** The methods and compositions of the invention comprise the administration of one or more EphA2 vaccines of the invention to subjects/patients suffering from or expected to suffer from cancer, *e.g.*, have a genetic predisposition for a particular type of cancer, have been exposed to a carcinogen, or are in remission from a particular cancer. As used herein, “cancer” refers to primary or metastatic cancers. Such patients may or may not have been previously treated for cancer. The methods and compositions of the invention may be used as a first line or second line cancer treatment. Included in the invention is also the treatment of patients undergoing other cancer therapies and the methods and compositions of the invention can be used before any adverse effects or intolerance of these other cancer therapies occurs. The invention also encompasses methods for administering one or more EphA2 vaccines of the invention to treat or ameliorate symptoms in refractory patients. In a certain embodiment, that a cancer is refractory to a therapy means that at least some significant portion of the cancer cells are not killed or their cell division arrested. The determination of whether the cancer cells are refractory can be made either *in vivo* or *in vitro* by any method known in the art for assaying the effectiveness of treatment on cancer cells, using the art-accepted meanings of “refractory” in such a context. In various embodiments, a cancer is refractory where the number of cancer cells has not been significantly reduced, or has increased. The invention also encompasses methods for administering one or more EphA2 vaccines to prevent the onset or recurrence of cancer in patients predisposed to having cancer.

**[00254]** In particular embodiments, the EphA2 vaccines of the invention are administered to reverse resistance or reduced sensitivity of cancer cells to certain hormonal, radiation and chemotherapeutic agents thereby resensitizing the cancer cells to one or more of these agents, which can then be administered (or continue to be administered) to treat or manage cancer, including to prevent metastasis. In a specific embodiment, the EphA2 vaccines of the invention are administered to patients with increased levels of the cytokine IL-6, which has been associated with the development of cancer cell resistance to different treatment regimens, such as chemotherapy and hormonal therapy. In another specific embodiment, the EphA2 vaccines of the invention are administered to patients suffering

from breast cancer that have a decreased responsiveness or are refractory to tamoxifen treatment. In another specific embodiment, the EphA2 vaccines of the invention are administered to patients with increased levels of the cytokine IL-6, which has been associated with the development of cancer cell resistance to different treatment regimens, such as chemotherapy and hormonal therapy.

[00255] In alternate embodiments, the invention provides methods for treating patients' cancer by administering one or more EphA2 vaccines of the invention in combination with any other treatment or to patients who have proven refractory to other treatments but are no longer on these treatments. In certain embodiments, the patients being treated by the methods of the invention are patients already being treated with chemotherapy, radiation therapy, hormonal therapy, or biological therapy/immunotherapy. Among these patients are refractory patients and those with cancer despite treatment with existing cancer therapies. In other embodiments, the patients have been treated and have no disease activity and one or more agonistic peptides of the invention are administered to prevent the recurrence of cancer.

[00256] In preferred embodiments, the existing treatment is chemotherapy. In particular embodiments, the existing treatment includes administration of chemotherapies including, but not limited to, methotrexate, taxol, mercaptopurine, thioguanine, hydroxyurea, cytarabine, cyclophosphamide, ifosfamide, nitrosoureas, cisplatin, carboplatin, mitomycin, dacarbazine, procarbazine, etoposides, campathecins, bleomycin, doxorubicin, idarubicin, daunorubicin, dactinomycin, plicamycin, mitoxantrone, asparaginase, vinblastine, vincristine, vinorelbine, paclitaxel, docetaxel, etc. Among these patients are patients treated with radiation therapy, hormonal therapy and/or biological therapy/immunotherapy. Also among these patients are those who have undergone surgery for the treatment of cancer.

[00257] Alternatively, the invention also encompasses methods for treating patients undergoing or having undergone radiation therapy. Among these are patients being treated or previously treated with chemotherapy, hormonal therapy and/or biological therapy/immunotherapy. Also among these patients are those who have undergone surgery for the treatment of cancer.

[00258] In other embodiments, the invention encompasses methods for treating patients undergoing or having undergone hormonal therapy and/or biological therapy/immunotherapy. Among these are patients being treated or having been treated with chemotherapy and/or radiation therapy. Also among these patients are those who have undergone surgery for the treatment of cancer.

[00259] Additionally, the invention also provides methods of treatment of cancer as an alternative to chemotherapy, radiation therapy, hormonal therapy, and/or biological therapy/immunotherapy where the therapy has proven or may prove too toxic, *i.e.*, results in unacceptable or unbearable side effects, for the subject being treated. The subject being  
5 treated with the methods of the invention may, optionally, be treated with other cancer treatments such as surgery, chemotherapy, radiation therapy, hormonal therapy or biological therapy, depending on which treatment was found to be unacceptable or unbearable.

[00260] In other embodiments, the invention provides administration of one or more EphA2 vaccines of the invention without any other cancer therapies for the treatment of  
10 cancer, but who have proved refractory to such treatments. In specific embodiments, patients refractory to other cancer therapies are administered one or more EphA2 vaccines in the absence of cancer therapies.

[00261] In other embodiments, patients with a pre-cancerous condition associated with cells that overexpress EphA2 can be administered vaccines of the invention to treat the  
15 disorder and decrease the likelihood that it will progress to malignant cancer. In a specific embodiments, the pre-cancerous condition is high-grade prostatic intraepithelial neoplasia (PIN), fibroadenoma of the breast, fibrocystic disease, or compound nevi.

[00262] In yet other embodiments, the invention provides methods of treating, preventing and managing non-cancer hyperproliferative cell disorders, particularly those  
20 associated with overexpression of EphA2, including but not limited to, asthma, chronic obstructive pulmonary disorder (COPD), restenosis (smooth muscle and/or endothelial), psoriasis, etc. These methods include methods analogous to those described above for treating, preventing and managing cancer, for example, by administering the EphA2 vaccines of the invention, combination therapy, administration to patients refractory to  
25 particular treatments, etc.

#### **5.9.1.1. Cancers**

[00263] Cancers and related disorders that can be treated, prevented, or managed by methods and compositions of the present invention include but are not limited to cancers of an epithelial cell origin. Examples of such cancers include the following: leukemias, such  
30 as but not limited to, acute leukemia, acute lymphocytic leukemia, acute myelocytic leukemias, such as, myeloblastic, promyelocytic, myelomonocytic, monocytic, and erythroleukemia leukemias and myelodysplastic syndrome; chronic leukemias, such as but not limited to, chronic myelocytic (granulocytic) leukemia, chronic lymphocytic leukemia, hairy cell leukemia; polycythemia vera; lymphomas such as but not limited to Hodgkin's  
35 disease, non-Hodgkin's disease; multiple myelomas such as but not limited to smoldering



multiple myeloma, nonsecretory myeloma, osteosclerotic myeloma, plasma cell leukemia, solitary plasmacytoma and extramedullary plasmacytoma; Waldenström's macroglobulinemia; monoclonal gammopathy of undetermined significance; benign monoclonal gammopathy; heavy chain disease; bone and connective tissue sarcomas such as but not limited to bone sarcoma, osteosarcoma, chondrosarcoma, Ewing's sarcoma, malignant giant cell tumor, fibrosarcoma of bone, chordoma, periosteal sarcoma, soft-tissue sarcomas, angiosarcoma (hemangiosarcoma), fibrosarcoma, Kaposi's sarcoma, leiomyosarcoma, liposarcoma, lymphangiosarcoma, neurilemmoma, rhabdomyosarcoma, synovial sarcoma; brain tumors such as but not limited to, glioma, astrocytoma, brain stem glioma, ependymoma, oligodendroglioma, nonglial tumor, acoustic neurinoma, craniopharyngioma, medulloblastoma, meningioma, pineocytoma, pineoblastoma, primary brain lymphoma; breast cancer including but not limited to ductal carcinoma, adenocarcinoma, lobular (small cell) carcinoma, intraductal carcinoma, medullary breast cancer, mucinous breast cancer, tubular breast cancer, papillary breast cancer, Paget's disease, and inflammatory breast cancer; adrenal cancer such as but not limited to pheochromocytoma and adrenocortical carcinoma; thyroid cancer such as but not limited to papillary or follicular thyroid cancer, medullary thyroid cancer and anaplastic thyroid cancer; pancreatic cancer such as but not limited to, insulinoma, gastrinoma, glucagonoma, vipoma, somatostatin-secreting tumor, and carcinoid or islet cell tumor; pituitary cancers such as but limited to Cushing's disease, prolactin-secreting tumor, acromegaly, and diabetes insipidus; eye cancers such as but not limited to ocular melanoma such as iris melanoma, choroidal melanoma, and ciliary body melanoma, and retinoblastoma; vaginal cancers such as squamous cell carcinoma, adenocarcinoma, and melanoma; vulvar cancer such as squamous cell carcinoma, melanoma, adenocarcinoma, basal cell carcinoma, sarcoma, and Paget's disease; cervical cancers such as but not limited to, squamous cell carcinoma, and adenocarcinoma; uterine cancers such as but not limited to endometrial carcinoma and uterine sarcoma; ovarian cancers such as but not limited to, ovarian epithelial carcinoma, borderline tumor, germ cell tumor, and stromal tumor; esophageal cancers such as but not limited to, squamous cancer, adenocarcinoma, adenoid cystic carcinoma, mucoepidermoid carcinoma, adenosquamous carcinoma, sarcoma, melanoma, plasmacytoma, verrucous carcinoma, and oat cell (small cell) carcinoma; stomach cancers such as but not limited to, adenocarcinoma, fungating (polypoid), ulcerating, superficial spreading, diffusely spreading, malignant lymphoma, liposarcoma, fibrosarcoma, and carcinosarcoma; colon cancers; rectal cancers; liver cancers such as but not limited to hepatocellular carcinoma and hepatoblastoma; gallbladder cancers such as adenocarcinoma;

- cholangiocarcinomas such as but not limited to pappillary, nodular, and diffuse; lung cancers such as non-small cell lung cancer, squamous cell carcinoma (epidermoid carcinoma), adenocarcinoma, large-cell carcinoma and small-cell lung cancer; testicular cancers such as but not limited to germinal tumor, seminoma, anaplastic, classic (typical),
- 5 spermatocytic, nonseminoma, embryonal carcinoma, teratoma carcinoma, choriocarcinoma (yolk-sac tumor), prostate cancers such as but not limited to, prostatic intraepithelial neoplasia, adenocarcinoma, leiomyosarcoma, and rhabdomyosarcoma; penial cancers; oral cancers such as but not limited to squamous cell carcinoma; basal cancers; salivary gland cancers such as but not limited to adenocarcinoma, mucoepidermoid carcinoma, and
- 10 adenoidcystic carcinoma; pharynx cancers such as but not limited to squamous cell cancer, and verrucous; skin cancers such as but not limited to, basal cell carcinoma, squamous cell carcinoma and melanoma, superficial spreading melanoma, nodular melanoma, lentigo malignant melanoma, acral lentiginous melanoma; kidney cancers such as but not limited to renal cell carcinoma, adenocarcinoma, hypernephroma, fibrosarcoma, transitional cell
- 15 cancer (renal pelvis and/ or uterus); Wilms' tumor; bladder cancers such as but not limited to transitional cell carcinoma, squamous cell cancer, adenocarcinoma, carcinosarcoma. In addition, cancers include myxosarcoma, osteogenic sarcoma, endotheliosarcoma, lymphangioendotheliosarcoma, mesothelioma, synovioma, hemangioblastoma, epithelial carcinoma, cystadenocarcinoma, bronchogenic carcinoma, sweat gland carcinoma,
- 20 sebaceous gland carcinoma, papillary carcinoma and papillary adenocarcinomas (for a review of such disorders, see Fishman et al., 1985, *Medicine*, 2d Ed., J.B. Lippincott Co., Philadelphia and Murphy et al., 1997, *Informed Decisions: The Complete Book of Cancer Diagnosis, Treatment, and Recovery*, Viking Penguin, Penguin Books U.S.A., Inc., United States of America)
- 25 **[00264]** Accordingly, the methods and compositions of the invention are also useful in the treatment or prevention of a variety of cancers or other abnormal proliferative diseases, including (but not limited to) the following: carcinoma, including that of the bladder, breast, ovary, oesophagus, colon, kidney, liver, lung, ovary, pancreas, stomach, cervix, thyroid and skin; including squamous cell carcinoma; hematopoietic tumors of
- 30 lymphoid lineage, including leukemia, acute lymphocytic leukemia, acute lymphoblastic leukemia, B-cell lymphoma, T-cell lymphoma, Burkitt's lymphoma; hematopoietic tumors of myeloid lineage, including acute and chronic myelogenous leukemias and promyelocytic leukemia; tumors of mesenchymal origin, including fibrosarcoma and rhabdomyosarcoma; other tumors, including melanoma, seminoma, tetratocarcinoma, neuroblastoma and
- 35 glioma; tumors of the central and peripheral nervous system, including astrocytoma,

neuroblastoma, glioma, and schwannomas; tumors of mesenchymal origin, including fibrosarcoma, rhabdomyosarcoma, and osteosarcoma; and other tumors, including melanoma, xeroderma pigmentosum, keratoactanthoma, seminoma, thyroid follicular cancer and teratocarcinoma. It is also contemplated that cancers caused by aberrations in apoptosis would also be treated by the methods and compositions of the invention. Such cancers may include but not be limited to follicular lymphomas, carcinomas with p53 mutations, hormone dependent tumors of the breast, prostate and ovary, and precancerous lesions such as familial adenomatous polyposis, and myelodysplastic syndromes. In specific embodiments, malignancy or dysproliferative changes (such as metaplasias and dysplasias), or hyperproliferative disorders, are treated or prevented in the skin, lung, colon, breast, prostate, bladder, kidney, pancreas, ovary, or uterus. In other specific embodiments, sarcoma, melanoma, or leukemia is treated or prevented.

[00265] In some embodiments, the cancer is malignant and overexpresses EphA2. In other embodiments, the disorder to be treated is a pre-cancerous condition associated with cells that overexpress EphA2. In a specific embodiment, the pre-cancerous condition is high-grade prostatic intraepithelial neoplasia (PIN), fibroadenoma of the breast, fibrocystic disease, or compound nevi.

[00266] In preferred embodiments, the methods and compositions of the invention are used for the treatment and/or prevention of breast, colon, ovarian, oesophageal, lung, and prostate cancers and melanoma and are provided below by example rather than by limitation.

[00267] In another preferred embodiment, the methods and compositions of the invention are used for the treatment and/or prevention of cancers of T cell origin, including, but not limited to, leukemias and lymphomas.

#### 5.9.1.2. Treatment of Breast Cancer

[00268] In specific embodiments, patients with breast cancer are administered an effective amount of one or more EphA2 vaccines of the invention. In another embodiment, the peptides of the invention can be administered in combination with an effective amount of one or more other agents useful for breast cancer therapy including but not limited to: doxorubicin, epirubicin, the combination of doxorubicin and cyclophosphamide (AC), the combination of cyclophosphamide, doxorubicin and 5-fluorouracil (CAF), the combination of cyclophosphamide, epirubicin and 5-fluorouracil (CEF), herceptin, tamoxifen, the combination of tamoxifen and cytotoxic chemotherapy, taxanes (such as docetaxel and paclitaxel). In a further embodiment, peptides of the invention can be administered with

taxanes plus standard doxorubicin and cyclophosphamide for adjuvant treatment of node-positive, localized breast cancer.

[00269] In a specific embodiment, patients with pre-cancerous fibroadenoma of the breast or fibrocystic disease are administered an EphA2 vaccine of the invention to treat the disorder and decrease the likelihood that it will progress to malignant breast cancer. In another specific embodiment, patients refractory to treatment, particularly hormonal therapy, more particularly tamoxifen therapy, are administered an EphA2 vaccine of the invention to treat the cancer and/or render the patient non-refractory or responsive.

#### **5.9.1.3. Treatment of Colon Cancer**

[00270] In specific embodiments, patients with colon cancer are administered an effective amount of one or more EphA2 vaccines of the invention. In another embodiment, the peptides of the invention can be administered in combination with an effective amount of one or more other agents useful for colon cancer therapy including but not limited to: the combination of 5-FU and leucovorin, the combination of 5-FU and levamisole, irinotecan (CPT-11) or the combination of irinotecan, 5-FU and leucovorin (IFL).

#### **5.9.1.4. Treatment of Prostate Cancer**

[00271] In specific embodiments, patients with prostate cancer are administered an effective amount of one or more EphA2 vaccines of the invention. In another embodiment, the peptides of the invention can be administered in combination with an effective amount of one or more other agents useful for prostate cancer therapy including but not limited to: external-beam radiation therapy, interstitial implantation of radioisotopes (*i.e.*,  $I^{125}$ , palladium, iridium), leuprolide or other LHRH agonists, non-steroidal antiandrogens (flutamide, nilutamide, bicalutamide), steroidal antiandrogens (cyproterone acetate), the combination of leuprolide and flutamide, estrogens such as DES, chlorotrianisene, ethinyl estradiol, conjugated estrogens U.S.P., DES-diphosphate, radioisotopes, such as strontium-89, the combination of external-beam radiation therapy and strontium-89, second-line hormonal therapies such as aminoglutethimide, hydrocortisone, flutamide withdrawal, progesterone, and ketoconazole, low-dose prednisone, or other chemotherapy regimens reported to produce subjective improvement in symptoms and reduction in PSA level including docetaxel, paclitaxel, estramustine/docetaxel, estramustine/etoposide, estramustine/vinblastine, and estramustine/paclitaxel.

[00272] In a specific embodiment, patients with pre-cancerous high-grade prostatic intraepithelial neoplasia (PIN) are administered an EphA2 vaccine of the invention to treat the disorder and decrease the likelihood that it will progress to malignant prostate cancer.

#### **5.9.1.5. Treatment of Melanoma**

[00273] In specific embodiments, patients with melanoma are administered an effective amount of one or more EphA2 vaccines of the invention. In another embodiment, the peptides of the invention can be administered in combination with an effective amount of one or more other agents useful for melanoma cancer therapy including but not limited to: dacarbazine (DTIC), nitrosoureas such as carmustine (BCNU) and lomustine (CCNU), agents with modest single agent activity including vinca alkaloids, platinum compounds, and taxanes, the Dartmouth regimen (cisplatin, BCNU, and DTIC), interferon alpha (IFN-A), and interleukin-2 (IL-2). In a specific embodiment, an effective amount of one or more EphA2 vaccines of the invention can be administered in combination with isolated hyperthermic limb perfusion (ILP) with melphalan (L-PAM), with or without tumor necrosis factor-alpha (TNF-alpha) to patients with multiple brain metastases, bone metastases, and spinal cord compression to achieve symptom relief and some shrinkage of the tumor with radiation therapy.

[00274] In a specific embodiment, patients with pre-cancerous compound nevi are administered an EphA2 vaccine of the invention to treat the disorder and decrease the likelihood that it will progress to malignant melanoma.

#### **5.9.1.6. Treatment of Ovarian Cancer**

[00275] In specific embodiments, patients with ovarian cancer are administered an effective amount of one or more EphA2 vaccines of the invention. In another embodiment, the peptides of the invention can be administered in combination with an effective amount of one or more other agents useful for ovarian cancer therapy including but not limited to: intraperitoneal radiation therapy, such as P<sup>32</sup> therapy, total abdominal and pelvic radiation therapy, cisplatin, the combination of paclitaxel (Taxol) or docetaxel (Taxotere) and cisplatin or carboplatin, the combination of cyclophosphamide and cisplatin, the combination of cyclophosphamide and carboplatin, the combination of 5-FU and leucovorin, etoposide, liposomal doxorubicin, gemcitabine or topotecan. It is contemplated that an effective amount of one or more EphA2 vaccines of the invention is administered in combination with the administration Taxol for patients with platinum-refractory disease.

Included is the treatment of patients with refractory ovarian cancer including administration of: ifosfamide in patients with disease that is platinum-refractory, hexamethylmelamine (HMM) as salvage chemotherapy after failure of cisplatin-based combination regimens, and tamoxifen in patients with detectable levels of cytoplasmic estrogen receptor on their tumors.

#### **5.9.1.7. Treatment of Lung Cancers**

[00276] In specific embodiments, patients with small lung cell cancer are administered an effective amount of one or more EphA2 vaccines of the invention. In another embodiment, the peptides of the invention can be administered in combination with an effective amount of one or more other agents useful for lung cancer therapy including but not limited to: thoracic radiation therapy, cisplatin, vincristine, doxorubicin, and etoposide, alone or in combination, the combination of cyclophosphamide, doxorubicin, vincristine/etoposide, and cisplatin (CAV/EP), local palliation with endobronchial laser therapy, endobronchial stents, and/or brachytherapy.

[00277] In other specific embodiments, patients with non-small lung cell cancer are administered an effective amount of one or more EphA2 vaccines of the invention in combination with an effective amount of one or more other agents useful for lung cancer therapy including but not limited to: palliative radiation therapy, the combination of cisplatin, vinblastine and mitomycin, the combination of cisplatin and vinorelbine, paclitaxel, docetaxel or gemcitabine, the combination of carboplatin and paclitaxel, interstitial radiation therapy for endobronchial lesions or stereotactic radiosurgery.

#### **5.9.1.8. Treatment of T Cell Malignancies**

[00278] In specific embodiments, patients with T cell malignancies, such as leukemias and lymphomas (see, e.g., section 5.9.1.1), are administered an effective amount of one or more EphA2 vaccines of the invention. In another embodiment, the EphA2 vaccines of the invention can be administered in combination with an effective amount of one or more other agents useful for the prevention, treatment or amelioration of cancer, particularly T cell malignancies or one or more symptoms thereof, said combination therapies comprising administering to a subject in need thereof a prophylactically or therapeutically effective amount of one or more EphA2 vaccines of the invention and a prophylactically or therapeutically effective amount of one or more cancer therapies, including chemotherapies, hormonal therapies, biological therapies, immunotherapies, or radiation therapies.

[00279] In another specific embodiment, patients with T cell malignancies are administered an effective amount of one or more EphA2 vaccines of the invention in combination with one or more cancer chemotherapeutic agents, such as but not limited to: doxorubicin, epirubicin, cyclophosphamide, 5-fluorouracil, taxanes such as docetaxel and paclitaxel, leucovorin, levamisole, irinotecan, estramustine, etoposide, vinblastine, dacarbazine, nitrosoureas such as carmustine and lomustine, vinca alkaloids, platinum compounds, cisplatin, mitomycin, vinorelbine, gemcitabine, carboplatin,

hexamethylmelamine and/or topotecan. Such methods can optionally further comprise the administration of other cancer therapies, such as but not limited to radiation therapy, biological therapies, hormonal therapies and/or surgery.

**[00280]** In yet another specific embodiment, patients with T cell malignancies are administered an effective amount of one or more EphA2 vaccines of the invention in combination with one or more types of radiation therapy, such as external-beam radiation therapy, interstitial implantation of radioisotopes (I-125, palladium, iridium), radioisotopes such as strontium-89, thoracic radiation therapy, intraperitoneal P-32 radiation therapy, and/or total abdominal and pelvic radiation therapy. Such methods can optionally further comprise the administration of other cancer therapies, such as but not limited to chemotherapies, biological therapies/immunotherapies, hormonal therapies and/or surgery.

**[00281]** In yet another specific embodiment, patients with T cell malignancies are administered an effective amount of one or more EphA2 vaccines of the invention in combination with one or more biological therapies/immunotherapies or hormonal therapies, such as tamoxifen, leuprolide or other LHRH agonists, non-steroidal antiandrogens (flutamide, nilutamide, bicalutamide), steroidal antiandrogens (cyproterone acetate), estrogens (DES, chlorotrianisene, ethinyl estradiol, conjugated estrogens U.S.P., DES-diphosphate), aminoglutethimide, hydrocortisone, flutamide withdrawal, progesterone, ketoconazole, prednisone, interferon- $\alpha$ , interleukin-2, tumor necrosis factor- $\alpha$ , and/or melphalan. Biological therapies also included are cytokines such as but not limited to TNF ligand family members such as TRAIL anti-cancer agonists that induce apoptosis, TRAIL antibodies that bind to TRAIL receptors 1 and 2 otherwise known as DR4 and DR5 (Death Domain Containing Receptors 4 and 5), as well as DR4 and DR5. TRAIL and TRAIL antibodies, ligands and receptors are known in the art and described in U.S. Patent Nos. 6,342,363, 6,284,236, 6,072,047 and 5,763,223. Such methods can optionally further comprise the administration of other cancer therapies, such as but not limited to radiation therapy, chemotherapies, and/or surgery.

**[00282]** In yet another specific embodiment, patients with T cell malignancies are administered an effective amount of one or more EphA2 vaccines of the invention in combination with standard and experimental therapies of T cell malignancies. Standard and experimental therapies of T cell malignancies that can be used in the methods and compositions of the invention include, but are not limited to, antibody therapy (e.g., Campath®, anti-Tac, HuM291 (humanized murine IgG2 monoclonal antibody against CD3), antibody drug conjugates (e.g., Mylotarg), radiolabeled monoclonal antibodies (e.g., Bexxar, Zevalin, Lym-1)), cytokine therapy, aggressive combination chemotherapy with or

without cytotoxic agents, purine analogs, hematopoietic stem cell transplantation, and T cell mediated therapy (e.g., CD8+ T cells with anti-leukemic activity against target antigens including but not limited to leukemia specific proteins (e.g., bcr/abl, PML/RAR $\alpha$ , EMV/AML-1), leukemia-associated proteins (e.g., proteinase 3, WT-1, h-TERT, hdm-2)).

- 5 (See Riddell et al., 2002, Cancer Control, 9(2): 114-122; Dearden et al. , 2002, Medical Oncology, 19, Suppl. S27-32; Waldmann et al. 2000, Hematology (Am Soc Hematol Educ Program):394-408).

### **5.9.2 Treatment or Prevention of Disorders Associated with Aberrant Angiogenesis**

- 10 [00283] EphA2 is as a marker of angiogenic blood vessels and plays a critical role in angiogenesis or neovascularization (see, e.g., Ogawa *et al.*, 2000, Oncogene. 19(52):6043-52; Hess *et al.*, 2001, Cancer Res. 61(8):3250-5). Angiogenesis is characterized by the invasion, migration and proliferation of smooth muscle and endothelial cells. The growth of new blood vessels, or angiogenesis, contributes to pathological conditions such as
- 15 diabetic retinopathy (Adonis *et al.*, Amer. J. Ophthal., Vol. 118, (1994) 445), rheumatoid arthritis (Peacock *et al.*, 1992, J. Exp. Med., 175:1135) and osteoarthritis (Ondrick *et al.*, 1992, Clin. Podiatr. Med. Surg. 9:185).
- [00284] The EphA2 vaccines of the invention may therefore be administered to a subject in need thereof to prevent, manage, treat or ameliorate a disorder associated with
- 20 aberrant angiogenesis or one or more symptoms thereof.
- [00285] Diseases or disorders that are associated with or characterized by aberrant angiogenesis and may be prevented, treated, managed, or ameliorated with the EphA2 vaccines of the invention include, but not limited to, neoplastic diseases (non-limiting examples are metastases of tumors and leukemia); diseases of ocular neovascularization
- 25 (non-limiting examples are age-related macular degeneration, diabetic retinopathy, and retinopathy of prematurity, vascular restenosis); skin diseases (non-limiting examples are infantile hemangiomas, verruca vulgaris, psoriasis, basal cell and squamous cell carcinomas, cutaneous melanoma, Kaposi's sarcoma, neurofibromatosis, recessive dystrophic epidermolysis bullosa); arthritis (non-limiting examples are rheumatoid arthritis,
- 30 ankylosing spondylitis, systemic lupus, psoriatic arthropathy, Reiter's syndrome, and Sjogren's syndrome); gynecologic diseases (non-limiting examples are endometriosis, preeclampsia during pregnancy, carcinoma of the ovary, endometrium and cervix); and cardiovascular diseases (non-limiting examples are formation of atherosclerotic plaques, atherosclerosis and coronary artery disease).



[00286] In specific embodiments, the diseases or disorders that are associated with or characterized by aberrant angiogenesis and that may be prevented, treated, managed, or ameliorated with the EphA2 vaccines of the invention include chronic articular rheumatism, psoriasis, diabetic retinopathy, neovascular glaucoma, macular degeneration, capillary proliferation in atherosclerotic plaques as well as cancers in which EphA2 is expressed in the vasculature. Such cancer disorders can include, for example, solid tumors, tumor metastasis, angiofibromas, retrolental, fibroplasia, hemangiomas, Kaposi's sarcoma.

[00287] In certain embodiments, the EphA2 vaccines are employed in combination therapy regimens involving other therapies. Non-limiting examples of such therapies include analgesics, angiogenesis inhibitors, anti-cancer therapies and anti-inflammatory agents, in particular analgesics and angiogenesis inhibitors.

### 5.9.3 Other Prophylactic/Therapeutic Agents

[00288] In some embodiments, therapy by administration of one or more EphA2 vaccines is combined with the administration of one or more therapies such as, but not limited to, chemotherapies, radiation therapies, hormonal therapies, and/or biological therapies/immunotherapies. Prophylactic/therapeutic agents include, but are not limited to, proteinaceous molecules, including, but not limited to, peptides, polypeptides, proteins, including post-translationally modified proteins, peptides etc.; or small molecules (less than 1000 daltons), inorganic or organic compounds; or nucleic acid molecules including, but not limited to, double-stranded or single-stranded DNA, or double-stranded or single-stranded RNA, as well as triple helix nucleic acid molecules. Prophylactic/therapeutic agents can be derived from any known organism (including, but not limited to, animals, plants, bacteria, fungi, and protista, or viruses) or from a library of synthetic molecules.

[00289] In a specific embodiment, the methods of the invention encompass administration of an EphA2 vaccine of the invention in combination with the administration of one or more prophylactic/therapeutic agents that are inhibitors of kinases such as, but not limited to, ABL, ACK, AFK, AKT (*e.g.*, AKT-1, AKT-2, and AKT-3), ALK, AMP-PK, ATM, Auroral, Aurora2, bARK1, bArk2, BLK, BMX, BTK, CAK, CaM kinase, CDC2, CDK, CK, COT, CTD, DNA-PK, EGF-R, ErbB-1, ErbB-2, ErbB-3, ErbB-4, ERK (*e.g.*, ERK1, ERK2, ERK3, ERK4, ERK5, ERK6, ERK7), ERT-PK, FAK, FGR (*e.g.*, FGF1R, FGF2R), FLT (*e.g.*, FLT-1, FLT-2, FLT-3, FLT-4), FRK, FYN, GSK (*e.g.*, GSK1, GSK2, GSK3-alpha, GSK3-beta, GSK4, GSK5), G-protein coupled receptor kinases (GRKs), HCK, HER2, HKII, JAK (*e.g.*, JAK1, JAK2, JAK3, JAK4), JNK (*e.g.*, JNK1, JNK2, JNK3), KDR, KIT, IGF-1 receptor, IKK-1, IKK-2, INSR (insulin receptor), IRAK1, IRAK2, IRK, ITK, LCK, LOK, LYN, MAPK, MAPKAPK-1, MAPKAPK-2, MEK, MET,

MFPK, MHCK, MLCK, MLK3, NEU, NIK, PDGF receptor alpha, PDGF receptor beta, PHK, PI-3 kinase, PKA, PKB, PKC, PKG, PRK1, PYK2, p38 kinases, p135tyk2, p34cdc2, p42cdc2, p42mapk, p44mpk, RAF, RET, RIP, RIP-2, RK, RON, RS kinase, SRC, SYK, S6K, TAK1, TEC, TIE1, TIE2, TRKA, TXK, TYK2, UL13, VEGFR1, VEGFR2, YES,

5 YRK, ZAP-70, and all subtypes of these kinases (see *e.g.*, Hardie and Hanks (1995) The Protein Kinase Facts Book, I and II, Academic Press, San Diego, Calif.). In preferred embodiments, an EphA2 vaccine of the invention is administered in combination with the administration of one or more prophylactic/therapeutic agents that are inhibitors of Eph receptor kinases (*e.g.*, EphA2, EphA4). In a most preferred embodiment, an EphA2 vaccine

10 of the invention is administered in combination with the administration of one or more prophylactic/therapeutic agents that are inhibitors of EphA2.

[00290] In another specific embodiment, the methods of the invention encompass administration of an EphA2 vaccine of the invention in combination with the administration of one or more prophylactic/therapeutic agents that are angiogenesis inhibitors such as, but

15 not limited to: Angiostatin (plasminogen fragment); antiangiogenic antithrombin III; Angiozyme; ABT-627; Bay 12-9566; Benefin; Bevacizumab; BMS-275291; cartilage-derived inhibitor (CDI); CAI; CD59 complement fragment; CEP-7055; Col 3; Combretastatin A-4; Endostatin (collagen XVIII fragment); fibronectin fragment; Gro-beta; Halofuginone; Heparinases; Heparin hexasaccharide fragment; HMV833; Human chorionic gonadotropin (hCG); IM-862; Interferon alpha/beta/gamma; Interferon inducible protein

20 (IP-10); Interleukin-12; Kringle 5 (plasminogen fragment); Marimastat; Metalloproteinase inhibitors (TIMPs); 2-Methoxyestradiol; MMI 270 (CGS 27023A); MoAb IMC-1C11; Neovastat; NM-3; Panzem; PI-88; Placental ribonuclease inhibitor; Plasminogen activator inhibitor; Platelet factor-4 (PF4); Prinomastat; Prolactin 16kD fragment; Proliferin-related

25 protein (PRP); PTK 787/ZK 222594; Retinoids; Solimastat; Squalamine; SS 3304; SU 5416; SU6668; SU11248; Tetrahydrocortisol-S; tetrathiomolybdate; thalidomide; Thrombospondin-1 (TSP-1); TNP-470; Transforming growth factor-beta (TGF-β); Vasculostatin; Vasostatin (calreticulin fragment); ZD6126; ZD6474; farnesyl transferase inhibitors (FTI); and bisphosphonates.

30 [00291] In another specific embodiment, the methods of the invention encompass administration of an EphA2 vaccine of the invention in combination with the administration of one or more prophylactic/therapeutic agents that are anti-cancer agents such as, but not limited to: acivicin, aclarubicin, acodazole hydrochloride, acronine, adozelesin, aldesleukin, altretamine, ambomycin, ametantrone acetate, aminoglutethimide, amsacrine,

35 anastrozole, anthramycin, asparaginase, asperlin, azacitidine, azetepa, azotomycin,

- batimastat, benzodepa, bicalutamide, bisantrene hydrochloride, bisnafide dimesylate, bizelesin, bleomycin sulfate, brequinar sodium, broprimine, busulfan, cactinomycin, calusterone, caracemide, carbetimer, carboplatin, carmustine, carubicin hydrochloride, carzelesin, cedefingol, chlorambucil, cirolemycin, cisplatin, cladribine, crisnatol mesylate,
- 5 cyclophosphamide, cytarabine, dacarbazine, dactinomycin, daunorubicin hydrochloride, decarbazine, decitabine, dexormaplatin, dezaguanine, dezaguanine mesylate, diaziquone, docetaxel, doxorubicin, doxorubicin hydrochloride, droloxifene, droloxifene citrate, dromostanolone propionate, duazomycin, edatrexate, eflornithine hydrochloride, elsamitucin, enloplatin, enpromate, epipropidine, epirubicin hydrochloride, erbulozole,
- 10 esorubicin hydrochloride, estramustine, estramustine phosphate sodium, etanidazole, etoposide, etoposide phosphate, etoprine, fadrozole hydrochloride, fazarabine, fenretinide, floxuridine, fludarabine phosphate, fluorouracil, flurocitabine, fosquidone, fostriecin sodium, gemcitabine, gemcitabine hydrochloride, hydroxyurea, idarubicin hydrochloride, ifosfamide, ilmofofene, interleukin 2 (including recombinant interleukin 2, or rIL2),
- 15 interferon alpha-2a, interferon alpha-2b, interferon alpha-n1, interferon alpha-n3, interferon beta-1 a, interferon gamma-1 b, iproplatin, irinotecan hydrochloride, lanreotide acetate, letrozole, leuprolide acetate, liarazole hydrochloride, lometrexol sodium, lomustine, losoxantrone hydrochloride, masoprocol, maytansine, mechlorethamine hydrochloride, megestrol acetate, melengestrol acetate, melphalan, menogaril, mercaptopurine,
- 20 methotrexate, methotrexate sodium, metoprine, meturedopa, mitindomide, mitocarcin, mitocromin, mitogillin, mitomalcin, mitomycin, mitosper, mitotane, mitoxantrone hydrochloride, mycophenolic acid, nitrosoureas, nocodazole, nogalamycin, ormaplatin, oxisuran, paclitaxel, pegaspargase, peliomycin, pentamustine, peplomycin sulfate, perfosfamide, pipobroman, piposulfan, piroxantrone hydrochloride, plicamycin,
- 25 plomestane, porfimer sodium, porfiromycin, prednimustine, procarbazine hydrochloride, puromycin, puromycin hydrochloride, pyrazofurin, riboprine, rogletimide, safingol, safingol hydrochloride, semustine, simtrazene, sparfosate sodium, sparsomycin, spirogermanium hydrochloride, spiromustine, spiroplatin, streptonigrin, streptozocin, sulofenur, talisomycin, tecogalan sodium, tegafur, teloxantrone hydrochloride, temoporfin, teniposide, teroxirone,
- 30 testolactone, thiamiprine, thioguanine, thiotepa, tiazaofurin, tirapazamine, toremifene citrate, trestolone acetate, triceribine phosphate, trimetrexate, trimetrexate glucuronate, triptorelin, tubulozole hydrochloride, uracil mustard, uredepa, vapreotide, verteporfin, vinblastine sulfate, vincristine sulfate, vindesine, vindesine sulfate, vinepidine sulfate, vinglycinat sulfate, vinleurosine sulfate, vinorelbine tartrate, vinrosidine sulfate, vinzolidine sulfate,
- 35 vorozole, zeniplatin, zinostatin, zorubicin hydrochloride. Other anti-cancer drugs include,

- but are not limited to: 20-epi-1,25 dihydroxyvitamin D3, 5-ethynyluracil, abiraterone, aclarubicin, acylfulvene, adecypenol, adozelesin, aldesleukin, ALL-TK antagonists, altretamine, ambamustine, amidox, amifostine, aminolevulinic acid, amrubicin, amsacrine, anagrelide, anastrozole, andrographolide, angiogenesis inhibitors, antagonist D, antagonist G, antarelix, anti-dorsalizing morphogenetic protein-1, antiandrogens, antiestrogens, antineoplaston, aphidicolin glycinate, apoptosis gene modulators, apoptosis regulators, apurinic acid, ara-CDP-DL-PTBA, arginine deaminase, asulacrine, atamestane, atrimustine, axinastatin 1, axinastatin 2, axinastatin 3, azasetron, azatoxin, azatyrosine, baccatin III derivatives, balanol, batimastat, BCR/ABL antagonists, benzochlorins,
- 10 benzoylstaurosporine, beta lactam derivatives, beta-alethine, betaclamycin B, betulinic acid, bFGF inhibitor, bicalutamide, bisantrene, bisaziridinylspermine, bisnafide, bistratene A, bizelesin, breflate, bropirimine, budotitane, buthionine sulfoximine, calcipotriol, calphostin C, camptothecin derivatives, canarypox IL-2, capecitabine, carboxamide-amino-triazole, carboxamidotriazole, CaRest M3, CARN 700, cartilage derived inhibitor, carzelesin,
- 15 casein kinase inhibitors (ICOS), castanospermine, cecropin B, cetorelix, chloroquinoxaline sulfonamide, cicaprost, cis-porphyrin, cladribine, clomifene analogues, clotrimazole, collismycin A, collismycin B, combretastatin A4, combretastatin analogue, conagenin, crambescidin 816, crisnatol, cryptophycin 8, cryptophycin A derivatives, curacin A, cyclopentantraquinones, cycloplatam, cypemycin, cytarabine ocfosfate, cytolytic factor,
- 20 cytostatin, daclicimab, decitabine, dehydrididemnin B, deslorelin, dexamethasone, dexifosfamide, dextrazoxane, dexverapamil, diaziquone, didemnin B, didox, diethylnorspermine, dihydro-5-azacytidine, dihydrotaxol, dioxamycin, diphenyl spiromustine, docetaxel, docosanol, dolasetron, doxifluridine, droloxifene, dronabinol, duocarmycin SA, ebselen, ecomustine, edelfosine, edrecolomab, eflornithine, elemene,
- 25 emitefur, epirubicin, epristeride, estramustine analogue, estrogen agonists, estrogen antagonists, etanidazole, etoposide phosphate, exemestane, fadrozole, fazarabine, fenretinide, filgrastim, finasteride, flavopiridol, flezelastine, fluasterone, fludarabine, fluorodaunorubicin hydrochloride, forfenimex, formestane, fostriecin, fotemustine, gadolinium texaphyrin, gallium nitrate, galocitabine, ganirelix, gelatinase inhibitors,
- 30 gemcitabine, glutathione inhibitors, hepsulfam, heregulin, hexamethylene bisacetamide, hypericin, ibandronic acid, idarubicin, idoxifene, idramantone, ilmofofene, ilomastat, imidazoacridones, imiquimod, immunostimulant peptides, insulin-like growth factor-1 receptor inhibitor, interferon agonists, interferons, interleukins, iobenguane, iododoxorubicin, ipomeanol, iroplact, irsogladine, isobengazole, isohomohalicondrin B,
- 35 itasetron, jasplakinolide, kahalalide F, lamellarin-N triacetate, lanreotide, leinamycin,

- lenograstim, lentinan sulfate, leptolstatin, letrozole, leukemia inhibiting factor, leukocyte alpha interferon, leuprolide+estrogen+progesterone, leuprorelin, levamisole, liarozole, linear polyamine analogue, lipophilic disaccharide peptide, lipophilic platinum compounds, lissoclinamide 7, lobaplatin, lombricine, lometrexol, lonidamine, losoxantrone, lovastatin,
- 5 loxoribine, lurtotecan, lutetium texaphyrin, lysofylline, lytic peptides, maitansine, mannostatin A, marimastat, masoprocol, maspin, matrilysin inhibitors, matrix metalloproteinase inhibitors, menogaril, merbarone, meterelin, methioninase, metoclopramide, MIF inhibitor, mifepristone, miltefosine, mirimostim, mismatched double stranded RNA, mitoguazone, mitolactol, mitomycin analogues, mitonafide, mitotoxin
- 10 fibroblast growth factor-saporin, mitoxantrone, mofarotene, molgramostim, EphA2 vaccine, human chorionic gonadotrophin, monophosphoryl lipid A+mycobacterium cell wall sk, mopidamol, multiple drug resistance gene inhibitor, multiple tumor suppressor 1-based therapy, mustard anticancer agent, mycaperoxide B, mycobacterial cell wall extract, myriaporone, N-acetyldinaline, N-substituted benzamides, nafarelin, nagrestip,
- 15 naloxone+pentazocine, napavin, naphterpin, nartograstim, nedaplatin, nemorubicin, neridronic acid, neutral endopeptidase, nilutamide, nisamycin, nitric oxide modulators, nitroxide antioxidant, nitrullyn, O6-benzylguanine, octreotide, okicenone, oligonucleotides, onapristone, ondansetron, ondansetron, oracin, oral cytokine inducer, ormaplatin, osaterone, oxaliplatin, oxanumycin, paclitaxel, paclitaxel analogues, paclitaxel derivatives,
- 20 palauamine, palmitoylrhizoxin, pamidronic acid, panaxytriol, panomifene, parabactin, pazelliptine, pegaspargase, peldesine, pentosan polysulfate sodium, pentostatin, pentozole, perflubron, perfosfamide, perillyl alcohol, phenazinomycin, phenylacetate, phosphatase inhibitors, picibanil, pilocarpine hydrochloride, pirarubicin, piritrexim, placetin A, placetin B, plasminogen activator inhibitor, platinum complex, platinum compounds, platinum-triamine complex, porfimer sodium, porfiromycin, prednisone, propyl bis-acridone, prostaglandin J2, proteasome inhibitors, protein A-based immune modulator, protein kinase C inhibitor, protein kinase C inhibitors, microalgal, protein tyrosine phosphatase inhibitors, purine nucleoside phosphorylase inhibitors, purpurins, pyrazoloacridine, pyridoxylated hemoglobin polyoxyethylene conjugate, raf antagonists, raltitrexed, ramosetron, ras farnesyl
- 30 protein transferase inhibitors, ras inhibitors, ras-GAP inhibitor, retelliptine demethylated, rhenium Re 186 etidronate, rhizoxin, ribozymes, RII retinamide, rogletimide, rohitukine, romurtide, roquinimex, rubiginone B1, ruboxyl, safingol, saintopin, SarCNU, sarcophytol A, sargramostim, Sdi 1 mimetics, semustine, senescence derived inhibitor 1, sense oligonucleotides, signal transduction inhibitors, signal transduction modulators, single chain
- 35 antigen binding protein, sizofiran, sobuzoxane, sodium borocaptate, sodium phenylacetate,

solverol, somatomedin binding protein, sonermin, sparfosic acid, spicamycin D, spiromustine, splenopentin, spongistatin 1, squalamine, stem cell inhibitor, stem-cell division inhibitors, stipiamide, stromelysin inhibitors, sulfinosine, superactive vasoactive intestinal peptide antagonist, suradista, suramin, swainsonine, synthetic

- 5 glycosaminoglycans, tallimustine, tamoxifen methiodide, tauromustine, taxol, tazarotene, tecogalan sodium, tegafur, tellurapyrylium, telomerase inhibitors, temoporfin, temozolomide, teniposide, tetrachlorodecaoxide, tetrazomine, thaliblastine, thalidomide, thiocoraline, thioguanine, thrombopoietin, thrombopoietin mimetic, thymalfasin, thymopoietin receptor agonist, thymotrinan, thyroid stimulating hormone, tin ethyl
- 10 etiopurpurin, tirapazamine, titanocene bichloride, topsentin, toremifene, totipotent stem cell factor, translation inhibitors, tretinoin, triacetyluridine, triciribine, trimetrexate, triptorelin, tropisetron, turosteride, tyrosine kinase inhibitors, tyrphostins, UBC inhibitors, ubenimex, urogenital sinus-derived growth inhibitory factor, urokinase receptor antagonists, vapreotide, variolin B, vector system, erythrocyte gene therapy, velaresol, veramine,
- 15 verdins, verteporfin, vinorelbine, vinxaltine, vitaxin, vorozole, zanoterone, zeniplatin, zilascorb, and zinostatin stimalamer. Preferred additional anti-cancer drugs are 5-fluorouracil and leucovorin.

**[00292]** In more particular embodiments, the present invention also comprises the administration of one or more EphA2 vaccines of the invention in combination with the administration of one or more therapies such as, but not limited to anti-cancer agents such as those disclosed in Table 2, preferably for the treatment of breast, ovary, melanoma, prostate, colon and lung cancers as described above.

20

**TABLE 2**

Therapeutic Agent	Administration	Dose	Intervals
doxorubicin hydrochloride (Adriamycin RDF® and Adriamycin PFS®)	Intravenous	60-75 mg/m <sup>2</sup> on Day 1	21 day intervals
epirubicin hydrochloride (Ellence™)	Intravenous	100-120 mg/m <sup>2</sup> on Day 1 of each cycle or divided equally and given on Days 1-8 of the cycle	3-4 week cycles
flourouracil	Intravenous	How supplied: 5 ml and 10 ml vials (containing 250 and 500 mg flourouracil respectively)	
docetaxel (Taxotere®)	Intravenous	60- 100 mg/m <sup>2</sup> over 1 hour	Once every 3 weeks
paclitaxel (Taxol®)	Intravenous	175 mg/m <sup>2</sup> over 3 hours	Every 3 weeks for 4 courses (administered sequentially to

Therapeutic Agent	Administration	Dose	Intervals
			doxorubicin-containing combination chemotherapy)
tamoxifen citrate (Nolvadex®)	Oral (tablet)	20-40 mg Dosages greater than 20 mg should be given in divided doses (morning and evening)	Daily
leucovorin calcium for injection	Intravenous or intramuscular injection	How supplied: 350 mg vial	Dosage is unclear from text. PDR 3610
luprolide acetate (Lupron®)	Single subcutaneous injection	1 mg (0.2 ml or 20 unit mark)	Once a day
flutamide (Eulexin®)	Oral (capsule)	250 mg (capsules contain 125 mg flutamide each)	3 times a day at 8 hour intervals (total daily dosage 750 mg)
nilutamide (Nilandron®)	Oral (tablet)	300 mg or 150 mg (tablets contain 50 or 150 mg nilutamide each)	300 mg once a day for 30 days followed by 150 mg once a day
bicalutamide (Casodex®)	Oral (tablet)	50 mg (tablets contain 50 mg bicalutamide each)	Once a day
progesterone	Injection	USP in sesame oil 50 mg/ml	
ketoconazole (Nizoral®)	Cream	2% cream applied once or twice daily depending on symptoms	
prednisone	Oral (tablet)	Initial dosage may vary from 5 mg to 60 mg per day depending on the specific disease entity being treated.	
estramustine phosphate sodium (Emcyt®)	Oral (capsule)	14 mg/ kg of body weight (i.e. one 140 mg capsule for each 10 kg or 22 lb of body weight)	Daily given in 3 or 4 divided doses
etoposide or VP-16	Intravenous	5 ml of 20 mg/ ml solution (100 mg)	
dacarbazine (DTIC-Dome®)	Intravenous	2-4.5 mg/knowning	Once a day for 10 days. May be repeated at 4 week intervals
polifeprosan 20 with carmustine implant (BCNU) (nitrosourea) (Gliadel®)	wafer placed in resection cavity	8 wafers, each containing 7.7 mg of carmustine, for a total of 61.6 mg, if size and shape of resection cavity allows	
cisplatin	Injection	How supplied: solution of 1 mg/ml in multi-dose vials of 50mL and 100mL	
mitomycin	Injection	supplied in 5 mg and 20 mg vials (containing 5 mg and 20 mg mitomycin)	
gemcitabine HCl (Gemzar®)	Intravenous	For NSCLC- 2 schedules have been investigated and the optimum schedule has not	4 week schedule- Days 1,8 and 15 of each 28-day cycle. Cisplatin

Therapeutic Agent	Administration	Dose	Intervals
		been determined 4 week schedule- administration intravenously at 1000 mg/m <sup>2</sup> over 30 minutes on 3 week schedule- Gemzar administered intravenously at 1250 mg/m <sup>2</sup> over 30 minutes	intravenously at 100 mg/m <sup>2</sup> on day 1 after the infusion of Gemzar. 3 week schedule- Days 1 and 8 of each 21 day cycle. Cisplatin at dosage of 100 mg/m <sup>2</sup> administered intravenously after administration of Gemzar on day 1.
carboplatin (Paraplatin®)	Intravenous	Single agent therapy: 360 mg/m <sup>2</sup> I.V. on day 1 (infusion lasting 15 minutes or longer)  Other dosage calculations: Combination therapy with cyclophosphamide, Dose adjustment recommendations, Formula dosing, etc.	Every 4 weeks
ifosamide (Ifex®)	Intravenous	1.2 g/m <sup>2</sup> daily	5 consecutive days Repeat every 3 weeks or after recovery from hematologic toxicity
topotecan hydrochloride (Hycamtin®)	Intravenous	1.5 mg/m <sup>2</sup> by intravenous infusion over 30 minutes daily	5 consecutive days, starting on day 1 of 21 day course

[00293] The invention also encompasses administration of the EphA2 vaccines of the invention in combination with radiation therapy comprising the use of x-rays, gamma rays and other sources of radiation to destroy the cancer cells. In preferred embodiments, the radiation treatment is administered as external beam radiation or teletherapy wherein the radiation is directed from a remote source. In other preferred embodiments, the radiation treatment is administered as internal therapy or brachytherapy wherein a radioactive source is placed inside the body close to cancer cells or a tumor mass.

[00294] Cancer therapies and their dosages, routes of administration and recommended usage are known in the art and have been described in such literature as the *Physician's Desk Reference* (56th ed., 2002).

#### **5.10 Characterization And Demonstration Of Therapeutic Or Prophylactic Utility**

[00295] Toxicity and efficacy of the prophylactic and/or therapeutic protocols of the instant invention can be determined by standard pharmaceutical procedures in experimental animals, e.g., for determining the LD<sub>50</sub> (the dose lethal to 50% of the population) and the ED<sub>50</sub> (the dose therapeutically effective in 50% of the population). The dose ratio between toxic and therapeutic effects is the therapeutic index and it can be expressed as the ratio



LD<sub>50</sub>/ED<sub>50</sub>. Prophylactic and/or therapeutic agents that exhibit large therapeutic indices are preferred. While prophylactic and/or therapeutic agents that exhibit toxic side effects may be used, care should be taken to design a delivery system that targets such agents to the site of affected tissue in order to minimize potential damage to uninfected cells and, thereby, reduce side effects.

[00296] The data obtained from the animal studies can be used in formulating a range of dosage of the prophylactic and/or therapeutic agents for use in humans. The dosage of such agents lies preferably within a range of circulating concentrations that include the ED<sub>50</sub> with little or no toxicity. The dosage may vary within this range depending upon the dosage form employed and the route of administration utilized. For any agent used in the method of the invention, the therapeutically effective dose can be estimated initially from cell culture assays. A dose may be formulated in animal models to achieve a circulating plasma concentration range that includes the IC<sub>50</sub> (*i.e.*, the concentration of the vaccine or test compound that achieves a half-maximal inhibition of symptoms) as determined in animal studies. Such information can be used to more accurately determine useful doses in humans. Levels in plasma may be measured, for example, by high performance liquid chromatography.

[00297] The anti-cancer activity of the therapies used in accordance with the present invention also can be determined by using various experimental animal models for the study of cancer, such as an immunocompetent mouse model, *e.g.*, Balb/c or C57/Bl/6, or transgenic mice where a mouse EphA2 is replaced with the human EphA2, mouse models to which murine tumor cell lines engineered to express human EphA2 are administered, animal models described in Section 6 *infra*, or any animal model (including hamsters, rabbits, etc.) known in the art and described in *Relevance of Tumor Models for Anticancer Drug Development* (1999, eds. Fiebig and Burger); *Contributions to Oncology* (1999, Karger); *The Nude Mouse in Oncology Research* (1991, eds. Boven and Winograd); and *Anticancer Drug Development Guide* (1997 ed. Teicher), herein incorporated by reference in their entireties.

[00298] Compounds for use in therapy can also be tested in other suitable animal model systems prior to testing in humans, including but not limited to in rats, mice, chicken, cows, monkeys, rabbits, hamsters, etc., for example, the animal models described above. The compounds can then be used in the appropriate clinical trials.

[00299] Further, any assays known to those skilled in the art can be used to evaluate the prophylactic and/or therapeutic utility of the vaccines and combinatorial therapies disclosed herein for treatment or prevention of hyperproliferative disorders such as cancer.

### 5.11 Vaccine Compositions

**[00300]** The compositions of the invention include bulk drug compositions useful in the manufacture of non-pharmaceutical compositions (*e.g.*, impure or non-sterile compositions) and pharmaceutical compositions (*i.e.*, compositions that are suitable for administration to a subject or patient) which can be used in the preparation of unit dosage forms. Such compositions comprise a prophylactically or therapeutically effective amount of a prophylactic and/or therapeutic agent disclosed herein or a combination of those agents and a pharmaceutically acceptable carrier. Preferably, compositions of the invention comprise a prophylactically or therapeutically effective amount of one or more EphA2 vaccines of the invention. The EphA2 vaccines of the invention may comprise one or more EphA2 antigenic peptides of the invention and a pharmaceutically acceptable carrier, one or more EphA2 antigenic peptide expression vehicles of the invention and a pharmaceutically acceptable carrier, or one or more antigen presenting cells sensitized with an EphA2 antigenic peptide and a pharmaceutically acceptable carrier.

**[00301]** Where an EphA2 vaccine of the invention comprises an EphA2 antigenic peptides, the EphA2 antigenic peptide of the invention can be modified. For example, in certain embodiments, the EphA2 antigenic peptide may be formulated with lipid as a lipopeptide or linked to a carrier molecule (and/or polymerized).

**[00302]** In a further embodiment, the composition of the invention further comprises an additional prophylactic or therapeutic, *e.g.*, anti-cancer, agent.

**[00303]** In a specific embodiment, the term “pharmaceutically acceptable” means approved by a regulatory agency of the Federal or a state government or listed in the U.S. Pharmacopeia or other generally recognized pharmacopeia for use in animals, and more particularly in humans. The term “carrier” refers to a diluent, adjuvant (*e.g.*, Freund’s adjuvant (complete and incomplete) or, more preferably, MF59C.1 adjuvant available from Chiron, Emeryville, CA), excipient, or vehicle with which the therapeutic is administered. Such pharmaceutical carriers can be sterile liquids, such as water and oils, including those of petroleum, animal, vegetable or synthetic origin, such as peanut oil, soybean oil, mineral oil, sesame oil and the like. Water is a preferred carrier when the pharmaceutical composition is administered intravenously. Saline solutions and aqueous dextrose and glycerol solutions can also be employed as liquid carriers, particularly for injectable solutions. Suitable pharmaceutical excipients include starch, glucose, lactose, sucrose, gelatin, malt, rice, flour, chalk, silica gel, sodium stearate, glycerol monostearate, talc, sodium chloride, dried skim milk, glycerol, propylene, glycol, water, ethanol and the like.

The composition, if desired, can also contain minor amounts of wetting or emulsifying

agents, or pH buffering agents. These compositions can take the form of solutions, suspensions, emulsion, tablets, pills, capsules, powders, sustained-release formulations and the like.

**[00304]** Generally, the ingredients of compositions of the invention are supplied either separately or mixed together in unit dosage form, for example, as a dry lyophilized powder or water free concentrate in a hermetically sealed container such as an ampoule or sachette indicating the quantity of active agent. Where the composition is to be administered by infusion, it can be dispensed with an infusion bottle containing sterile pharmaceutical grade water or saline. Where the composition is administered by injection, an ampoule of sterile water for injection or saline can be provided so that the ingredients may be mixed prior to administration.

**[00305]** The compositions of the invention can be formulated as neutral or salt forms. Pharmaceutically acceptable salts include those formed with anions such as those derived from hydrochloric, phosphoric, acetic, oxalic, tartaric acids, etc., and those formed with cations such as those derived from sodium, potassium, ammonium, calcium, ferric hydroxides, isopropylamine, triethylamine, 2-ethylamino ethanol, histidine, procaine, etc.

**[00306]** Various delivery systems are known and can be used to administer an EphA2 vaccine of the invention or the combination of an EphA2 vaccine of the invention and a prophylactic agent or therapeutic agent useful for preventing or treating cancer, e.g., encapsulation in liposomes, microparticles, microcapsules, recombinant cells capable of expressing the EphA2 antigenic peptide, receptor-mediated endocytosis (see, e.g., Wu and Wu, 1987, *J. Biol. Chem.* 262:4429-4432), construction of a nucleic acid as part of a retroviral or other vector, etc. Methods of administering an EphA2 vaccine or the combination of an EphA2 vaccine of the invention and prophylactic or therapeutic agent, but are not limited to, parenteral administration (e.g., intradermal, intramuscular, intraperitoneal, intravenous and subcutaneous), epidural, and mucosal (e.g., intranasal, inhaled, and oral routes). In a specific embodiment, an EphA2 vaccine of the invention or the combination of an EphA2 vaccine of the invention and prophylactic or therapeutic agent are administered intramuscularly, intravenously, or subcutaneously. The EphA2 vaccine of the invention or the combination of an EphA2 vaccine of the invention and prophylactic or therapeutic agent may be administered by any convenient route, for example by infusion or bolus injection, by absorption through epithelial or mucocutaneous linings (e.g., oral mucosa, rectal and intestinal mucosa, etc.) and may be administered together with other biologically active agents. Administration can be systemic or local.

- [00307]** In a specific embodiment, it may be desirable to administer the EphA2 vaccine of the invention or the combination of an EphA2 vaccine of the invention and prophylactic or therapeutic agents of the invention locally to the area in need of treatment; this may be achieved by, for example, and not by way of limitation, local infusion, by
- 5 injection, or by means of an implant, said implant being of a porous, non-porous, or gelatinous material, including membranes, such as sialastic membranes, or fibers.
- [00308]** In yet another embodiment, the EphA2 vaccine of the invention or the combination of an EphA2 vaccine of the invention and prophylactic or therapeutic agent can be delivered in a controlled release or sustained release system. In one embodiment, a
- 10 pump may be used to achieve controlled or sustained release (see Langer, *supra*; Sefton, 1987, *CRC Crit. Ref. Biomed. Eng.* 14:20; Buchwald et al., 1980, *Surgery* 88:507; Saudek et al., 1989, *N. Engl. J. Med.* 321:574). In another embodiment, polymeric materials can be used to achieve controlled or sustained release of the EphA2 antigenic peptides of the invention (see *e.g.*, Medical Applications of Controlled Release, Langer and Wise (eds.),
- 15 CRC Pres., Boca Raton, Florida (1974); Controlled Drug Bioavailability, Drug Product Design and Performance, Smolen and Ball (eds.), Wiley, New York (1984); Ranger and Peppas, 1983, *J. Macromol. Sci. Rev. Macromol. Chem.* 23:61; see also Levy et al., 1985, *Science* 228:190; During et al., 1989, *Ann. Neurol.* 25:351; Howard et al., 1989, *J. Neurosurg.* 7 1:105); U.S. Patent Nos. 5,679,377; 5,916,597; 5,912,015; 5,989,463;
- 20 5,128,326; International Publication Nos. WO 99/15154 and WO 99/20253. Examples of polymers used in sustained release formulations include, but are not limited to, poly(2-hydroxy ethyl methacrylate), poly(methyl methacrylate), poly(acrylic acid), poly(ethylene-co-vinyl acetate), poly(methacrylic acid), polyglycolides (PLG), polyanhydrides, poly(N-vinyl pyrrolidone), poly(vinyl alcohol), polyacrylamide, poly(ethylene glycol), polylactides
- 25 (PLA), poly(lactide-co-glycolides) (PLGA), and polyorthoesters. In a preferred embodiment, the polymer used in a sustained release formulation is inert, free of leachable impurities, stable on storage, sterile, and biodegradable. In yet another embodiment, a controlled or sustained release system can be placed in proximity of the prophylactic or therapeutic target, thus requiring only a fraction of the systemic dose (see, *e.g.*, Goodson, in
- 30 Medical Applications of Controlled Release, *supra*, vol. 2, pp. 115-138 (1984)).
- [00309]** Controlled release systems are discussed in the review by Langer (1990, *Science* 249:1527-1533). Any technique known to one of skill in the art can be used to produce sustained release formulations comprising one or more therapeutic agents of the invention. See, *e.g.*, U.S. Patent No. 4,526,938; International Publication Nos. WO
- 35 91/05548 and WO 96/20698; Ning et al., 1996, *Radiotherapy & Oncology* 39:179-189;

Song et al., 1995, *PDA Journal of Pharmaceutical Science & Technology* 50:372-397; Cleek et al., 1997, *Pro. Int'l. Symp. Control. Rel. Bioact. Mater.* 24:853-854; and Lam et al., 1997, *Proc. Int'l. Symp. Control Rel. Bioact. Mater.* 24:759-760, each of which is incorporated herein by reference in its entirety.

5                                    **5.11.1 Formulations**

[00310]            Pharmaceutical compositions for use in accordance with the present invention may be formulated in conventional manner using one or more physiologically acceptable carriers or excipients.

10            [00311]            Thus, the EphA2 antigenic peptides of the invention and their physiologically acceptable salts and solvates (or EphA2 antigenic peptide expression vehicles or antigen presenting cells sensitized with an EphA2 antigenic peptide) may be formulated for administration by inhalation or insufflation (either through the mouth or the nose) or oral, parenteral or mucosal (such as buccal, vaginal, rectal, sublingual) administration. In a preferred embodiment, local or systemic parenteral administration is  
15            used.

[00312]            For oral administration, the vaccine may take the form of, for example, tablets or capsules prepared by conventional means with pharmaceutically acceptable excipients such as binding agents (*e.g.*, pregelatinised maize starch, polyvinylpyrrolidone or hydroxypropyl methylcellulose); fillers (*e.g.*, lactose, microcrystalline cellulose or calcium  
20            hydrogen phosphate); lubricants (*e.g.*, magnesium stearate, talc or silica); disintegrants (*e.g.*, potato starch or sodium starch glycolate); or wetting agents (*e.g.*, sodium lauryl sulphate). The tablets may be coated by methods well known in the art. Liquid preparations for oral administration may take the form of, for example, solutions, syrups or suspensions, or they may be presented as a dry product for constitution with water or other suitable vehicle  
25            before use. Such liquid preparations may be prepared by conventional means with pharmaceutically acceptable additives such as suspending agents (*e.g.*, sorbitol syrup, cellulose derivatives or hydrogenated edible fats); emulsifying agents (*e.g.*, lecithin or acacia); non-aqueous vehicles (*e.g.*, almond oil, oily esters, ethyl alcohol or fractionated vegetable oils); and preservatives (*e.g.*, methyl or propyl-p-hydroxybenzoates or sorbic  
30            acid). The preparations may also contain buffer salts, flavoring, coloring and sweetening agents as appropriate.

[00313]            Preparations for oral administration may be suitably formulated to give controlled release of the active compound.

35            [00314]            For buccal administration the compositions may take the form of tablets or lozenges formulated in conventional manner.

**[00315]** For administration by inhalation, the prophylactic or therapeutic agents for use according to the present invention are conveniently delivered in the form of an aerosol spray presentation from pressurized packs or a nebulizer, with the use of a suitable propellant, *e.g.*, dichlorodifluoromethane, trichlorofluoromethane,

- 5 dichlorotetrafluoroethane, carbon dioxide or other suitable gas. In the case of a pressurized aerosol the dosage unit may be determined by providing a valve to deliver a metered amount. Capsules and cartridges of *e.g.*, gelatin for use in an inhaler or insufflator may be formulated containing a powder mix of the compound and a suitable powder base such as lactose or starch.

- 10 **[00316]** The EphA2 vaccine may be formulated for parenteral administration by injection, *e.g.*, by bolus injection or continuous infusion. Formulations for injection may be presented in unit dosage form, *e.g.*, in ampoules or in multi-dose containers, with an added preservative. The compositions may take such forms as suspensions, solutions or emulsions in oily or aqueous vehicles, and may contain formulatory agents such as suspending,
- 15 stabilizing and/or dispersing agents. Alternatively, the active ingredient may be in powder form for constitution with a suitable vehicle, *e.g.*, sterile pyrogen-free water, before use.

**[00317]** The vaccines of the invention may also be formulated in rectal compositions such as suppositories or retention enemas, *e.g.*, containing conventional suppository bases such as cocoa butter or other glycerides.

- 20 **[00318]** In addition to the formulations described previously, the prophylactic or therapeutic agents may also be formulated as a depot preparation. Such long acting formulations may be administered by implantation (for example subcutaneously or intramuscularly) or by intramuscular injection. Thus, for example, the prophylactic or therapeutic agents may be formulated with suitable polymeric or hydrophobic materials (for
- 25 example as an emulsion in an acceptable oil) or ion exchange resins, or as sparingly soluble derivatives, for example, as a sparingly soluble salt.

**[00319]** The invention also provides that an EphA2 vaccine of the invention is packaged in a hermetically sealed container such as an ampoule or sachette indicating the quantity. In one embodiment, the vaccine is supplied as a dry sterilized lyophilized powder

30 or water free concentrate in a hermetically sealed container and can be reconstituted, *e.g.*, with water or saline to the appropriate concentration for administration to a subject.

**[00320]** In a preferred embodiment of the invention, the formulation and administration of various chemotherapeutic, biological/immunotherapeutic and hormonal therapeutic agents for use in combination with the vaccine of the invention are known in the

35 art and often described in the *Physician's Desk Reference*, 56<sup>th</sup> ed. (2002). For instance, in

certain specific embodiments of the invention, the agents can be formulated and supplied as provided in Table 2.

[00321] In other embodiments of the invention, radiation therapy agents such as radioactive isotopes can be given orally as liquids in capsules or as a drink. Radioactive isotopes can also be formulated for intravenous injections. The skilled oncologist can determine the preferred formulation and route of administration.

[00322] In certain embodiments the EphA2 antigenic peptides and anti-idiotypic antibodies of the invention are formulated at 1 mg/ml, 5 mg/ml, 10 mg/ml, and 25 mg/ml for intravenous injections and at 5 mg/ml, 10 mg/ml, and 80 mg/ml for repeated subcutaneous administration and intramuscular injection.

[00323] Where the EphA2 vaccine is a bacterial vaccine, the vaccine can be formulated at amounts ranging between approximately  $1 \times 10^2$  CFU/ml to approximately  $1 \times 10^{12}$  CFU/ml, for example at  $1 \times 10^2$  CFU/ml,  $5 \times 10^2$  CFU/ml,  $1 \times 10^3$  CFU/ml,  $5 \times 10^3$  CFU/ml,  $1 \times 10^4$  CFU/ml,  $5 \times 10^4$  CFU/ml,  $1 \times 10^5$  CFU/ml,  $5 \times 10^5$  CFU/ml,  $1 \times 10^6$  CFU/ml,  $5 \times 10^6$  CFU/ml,  $1 \times 10^7$  CFU/ml,  $5 \times 10^7$  CFU/ml,  $1 \times 10^8$  CFU/ml,  $5 \times 10^8$  CFU/ml,  $1 \times 10^9$  CFU/ml,  $5 \times 10^9$  CFU/ml,  $1 \times 10^{10}$  CFU/ml,  $5 \times 10^{10}$  CFU/ml,  $1 \times 10^{11}$  CFU/ml,  $5 \times 10^{11}$  CFU/ml, or  $1 \times 10^{12}$  CFU/ml.

[00324] The compositions may, if desired, be presented in a pack or dispenser device that may contain one or more unit dosage forms containing the active ingredient. The pack may for example comprise metal or plastic foil, such as a blister pack. The pack or dispenser device may be accompanied by instructions for administration.

### **5.11.2 Dosages**

[00325] The amount of the composition of the invention which will be effective in the treatment, prevention or management of cancer can be determined by standard research techniques. For example, the dosage of the EphA2 vaccine of the invention which will be effective in the treatment, prevention or management of cancer can be determined by administering the composition to an animal model such as, e.g., the animal models disclosed herein or known to those skilled in the art. In addition, *in vitro* assays may optionally be employed to help identify optimal dosage ranges.

[00326] Selection of the preferred effective dose can be determined (e.g., via clinical trials) by a skilled artisan based upon the consideration of several factors which will be known to one of ordinary skill in the art. Such factors include the disease to be treated or prevented, the symptoms involved, the patient's body mass, the patient's immune status and other factors known by the skilled artisan to reflect the accuracy of administered pharmaceutical compositions.

[00327] The precise dose to be employed in the formulation will also depend on the route of administration, and the seriousness of the cancer, and should be decided according to the judgment of the practitioner and each patient's circumstances. Effective doses may be extrapolated from dose-response curves derived from *in vitro* or animal model test systems.

[00328] For EphA2 antigenic peptides or anti-idiotypic antibodies, the dosage administered to a patient is typically 0.1 mg/kg to 100 mg/kg of the patient's body weight. Preferably, the dosage administered to a patient is between 0.1 mg/kg and 20 mg/kg of the patient's body weight, more preferably 1 mg/kg to 10 mg/kg of the patient's body weight.

[00329] With respect to the dosage of bacterial EphA2 vaccines of the invention, the dosage is based on the amount colony forming units (c.f.u.). Generally, in various embodiments, the dosage ranges are from about 1.0 c.f.u./kg to about  $1 \times 10^{10}$  c.f.u./kg; from about 1.0 c.f.u./kg to about  $1 \times 10^8$  c.f.u./kg; from about  $1 \times 10^2$  c.f.u./kg to about  $1 \times 10^8$  c.f.u./kg; and from about  $1 \times 10^4$  c.f.u./kg to about  $1 \times 10^8$  c.f.u./kg. Effective doses may be extrapolated from dose-response curves derived animal model test systems. In certain exemplary embodiments, the dosage ranges are 0.001-fold to 10,000-fold of the murine LD<sub>50</sub>, 0.01-fold to 1,000-fold of the murine LD<sub>50</sub>, 0.1-fold to 500-fold of the murine LD<sub>50</sub>, 0.5-fold to 250-fold of the murine LD<sub>50</sub>, 1-fold to 100-fold of the murine LD<sub>50</sub>, and 5-fold to 50-fold of the murine LD<sub>50</sub>. In certain specific embodiments, the dosage ranges are 0.001-fold, 0.01-fold, 0.1-fold, 0.5-fold, 1-fold, 5-fold, 10-fold, 50-fold, 100-fold, 200-fold, 500-fold, 1,000-fold, 5,000-fold or 10,000-fold of the murine LD<sub>50</sub>.

[00330] For other cancer therapeutic agents administered to a patient, the typical doses of various cancer therapeutics known in the art are provided in Table 2. Given the invention, certain preferred embodiments will encompass the administration of lower dosages in combination treatment regimens than dosages recommended for the administration of single agents.

[00331] The invention provides for any method of administering lower doses of known prophylactic or therapeutic agents than previously thought to be effective for the prevention, treatment, management or amelioration of cancer. Preferably, lower doses of known anti-cancer therapies are administered in combination with lower doses of EphA2 vaccines of the invention.

#### 5.12 Kits

[00332] The invention provides a pack or kit comprising one or more containers filled with an EphA2 vaccine of the invention or a component of an EphA2 vaccine of the invention. Additionally, one or more other prophylactic or therapeutic agents useful for the



treatment of a cancer or other hyperproliferative disorder can also be included in the pack or kit. Optionally associated with such container(s) can be a notice in the form prescribed by a governmental agency regulating the manufacture, use or sale of pharmaceuticals or biological products, which notice reflects approval by the agency of manufacture, use or sale for human administration.

[00333] The present invention provides kits that can be used in the above methods. In one embodiment, a kit comprises one or more a EphA2 vaccines of the invention. In another embodiment, a kit further comprises one or more other prophylactic or therapeutic agents useful for the treatment of cancer or another hyperproliferative disorder, in one or more containers. In other embodiments, the prophylactic or therapeutic agent is a biological or hormonal therapeutic.

**6. EXAMPLES: LISTERIA-BASED EPHA2 VACCINES PROVIDE THERAPEUTIC AND PROPHYLACTIC BENEFITS AGAINST EPHA2-EXPRESSING CANCERS**

[00334] The receptor tyrosine kinase EphA2 is selectively over-expressed in a variety of malignant cell types and tumors. Additionally, recent studies have identified patient-derived T lymphocytes that recognize EphA2. As such, EphA2 provides a much-needed target for active immunotherapy. Here, we show that ectopic expression of human EphA2 in the Gram-positive facultative intracellular bacterium *Listeria monocytogenes* (*Listeria*) can provide antigen-specific anti-tumor responses in vaccinated animals. *Listeria* infects critical antigen presenting cells and thereby provides efficacy as a cancer therapy based its ability to induce potent and robust CD4+ and CD8+ T cell responses against encoded antigens. Attenuated *Listeria* mutant strains, which retain the antigen delivery potency of wild-type bacteria, yet are nearly 10,000-fold less pathogenic in mice, were employed. To demonstrate the efficacy of a *Listeria*-based EphA2 vaccine, *Listeria actA*<sup>-</sup> strains were engineered to express the extracellular (ECD) or intracellular (ICD) domain of human EphA2 (actA-hEphA2-ECD or actA-hEphA2-ICD). Expression and secretion of hEphA2-EX and -CO from *Listeria* was confirmed by Western blot analysis. Protective immunization with actA-hEphA2EX significantly inhibited the subcutaneous growth of CT26 cells that express full-length hEphA2 (p=0.0037). As controls, mice vaccinated with the parental actA strain developed tumors that were comparable to vehicle-treated control mice. Protective immunization with actA-hEphA2CO significantly increased the survival rate in mice challenged with RenCA-hEphA2. Subsequently, the therapeutic efficacy of actA-hEphA2-ECD or actA-hEphA2-ICD was evaluated using the experimental CT26-hEphA2 lung tumor model. Following intravenous implantation of tumor cells, Balb/c mice

were immunized with actA, actA-hEphA2EX or actA-hEphA2-ICD. Immunization with either actA-hEphA2-ECD or actA-hEphA2-ICD significantly prolonged survival (median survival >43 days,  $p=0.0035$ ), as compared to matched controls (vehicle or actA median survival time was 19 and 20 days, respectively). Importantly, 80% of the huEphA2

5 immunized mice survived until Day 43 following tumor implantation. Together, these data demonstrate that *Listeria*-mediated vaccination targeting the EphA2 tumor antigen can provide both preventative and therapeutic efficacy against a variety of malignancies.

## 6.1 EXAMPLE 1: *LISTERIA* LIFE CYCLE

[00335] The life cycle of *Listeria monocytogenes*, encompassing the steps of  
10 endocytosis, phagolysosomal lysis, and cell to cell spread, are shown in Figure 1A-1B.

## 6.2 EXAMPLE 2: CONSTRUCTION OF EphA2-EXPRESSING AND CONTROL *LISTERIA* STRAINS

### 6.2.1 Background

15 [00336] Given the mechanisms by which *Listeria* programs the presentation of heterologous antigens via the MHC class I pathway, the efficiency of both expression of heterologous genes and secretion of the newly synthesized protein from the bacterium into the cytoplasm of the infected (antigen presenting) cell is related directly to the potency of CD8+ T cell priming and/or activation. As the level of Ag-specific T cell priming is related  
20 directly to vaccine efficacy, the efficiency of heterologous protein expression and secretion is linked directly to vaccine potency. Thus, the efficiency of EphA2 expression and secretion was optimized to maximize the potency of *Listeria*-based vaccines, in terms of priming and/or activating CD8+ T cell responses specific for the encoded EphA2 protein.

### 6.2.2 Preparation of mutant *Listeria* strains.

[00338] In some vaccines, mutant strains of *Listeria* that are deficient with respect to internalin B (Genbank accession number AL591975 (*Listeria monocytogenes* strain EGD, complete genome, segment 3/12; *inlB* gene region: nts. 97008-98963), incorporated by reference herein in its entirety, and/or the sequence listed as Genbank accession number NC\_003210 (*Listeria monocytogenes* strain EGD, complete genome, *inlB* gene region: nts. 457008-458963), incorporated by reference herein in its entirety) are used. One particular *actA<sup>+</sup>inlB<sup>-</sup>* strain (DP-L4029*inlB*) was deposited with the American Type Culture Collection (ATCC) on October 3, 2003, and designated with accession number PTA-5562).

### 6.2.3 Cloning vectors

[00339] Selected heterologous antigen expression cassette molecular constructs were inserted into pPL2 (Lauer et. al. J. Bacteriol. 2002), or pAM401 (Wirth et. al., J. Bacteriol. 165:831-836), modified to contain the multiple cloning sequence of pPL2 (Aat II small fragment, 171 bps), inserted between blunted *Xba* I and *Nru* I recognition sites, within the tetracycline resistance gene (pAM401-MCS). In general, the *hly* promoter and (selected) signal peptide sequence was inserted between the unique *Kpn* I and *Bam* HI sites in the pPL2 or pAM401-MCS plasmid vectors. Selected *EphA2* genes (sometimes modified to contain N-terminal and C-terminal epitope tags; see description below) were cloned subsequently into these constructs between unique *Bam* HI and *Sac* I sites. Molecular constructs based on the pAM401-MCS plasmid vector were introduced by electroporation into selected *Listeria monocytogenes* strains also treated with lysozyme, utilizing methods common to those skilled in the art. The expected plasmid structure in *Listeria*-transfectants was verified by isolating DNA from colonies that formed on chloramphenicol-containing BHI agar plates (10 µg/ml) by restriction enzyme analysis. Recombinant *Listeria* transformed with various pAM401-MCS based heterologous protein expression cassette constructs were utilized to measure heterologous protein expression and secretion, as described below.

[00340] The pPL2 based heterologous protein expression cassette constructs were incorporated into the *tRNA<sup>Arg</sup>* gene in the genome of selected *Listeria* strains, according to the methods as described previously (Lauer et al., 2002, J. Bacteriol. 184:4177-4186). Briefly, the pPL2 heterologous protein expression cassette constructs plasmid was first introduced into the *E. coli* host strain SM10 (Simon et al., 1983, BioTechnology 1:784-791) by electroporation or by chemical means. Subsequently, the pPL2-based plasmid was transferred from transformed SM10 to the selected *Listeria* strains by conjugation. Following incubation on drug-selective BHI agar plates containing 7.5 µg of chloramphenicol per ml and 200 µg of streptomycin per ml as described, selected colonies

are purified by passing 3 times on plates with the same composition. To verify integration of the pPL2 vector at the phage attachment site, individual colonies are picked and screened by PCR using the primer pair of forward primer NC16 (5'-gtcaaacatacgtcttcttc-3') (SEQ ID NO:47) and reverse primer PL95 (5'-acataatcagtcacaaagtagatgc-3') (SEQ ID NO:48). Selected colonies having the pPL2-based plasmid incorporated into the tRNA<sup>Arg</sup> gene in the genome of selected *Listeria* strains yielded a diagnostic DNA amplicon of 499 bps.

#### 6.2.4 Promoter

[00341] Heterologous protein expression cassettes contained the *prfA*-dependent *hly* promoter, which drives the transcription of the gene encoding Listeriolysin O (LLO), and is activated within the microenvironment of the infected cell. Nucleotides 205586-206000 (414 bps) were amplified by PCR from *Listeria monocytogenes*, strain DP-L4056, using the primer pair shown below. The region amplified includes the *hly* promoter and also the first 28 amino acids of LLO, comprising the *secA1* signal peptide (ibid) and PEST domain. The expected sequence of this region for *Listeria monocytogenes*, strain EGD can be found in GenBank (Accession number: gi|16802048|ref|NC\_003210.1[16802048]).

[00342] Primer Pair

[00343] *Forward* (KpnI-LLO nts. 1257-1276):

[00344] 5'-CTCTGGTACCTCCTTTGATTAGTATATTC (SEQ ID NO:49)

[00345] *Reverse* (Bam HI-LLO nts. X-x):

[00346] 5'-CTCTGGATCCATCCGCGTGTTCCTTTTCG (SEQ ID NO:50)

[00347] (Restriction endonuclease recognition sites are underlined)

[00348] The 422 bp PCR amplicon was cloned into the plasmid vector pCR-XL-

TOPO (Invitrogen, Carlsbad, CA), according to the manufacturer's specifications. The

nucleotide sequences of *Listeria*-specific bases in the pCR-XL-TOPO-*hly* promoter plasmid clone was determined. *Listeria monocytogenes* strain DP-L4056 contained eight nucleotide base changes flanking the *prfA* box in the *hly* promoter, as compared to the EGD strain. The *hly* promoter alignment for the *Listeria monocytogenes* DP-L4056 and EGD strains is shown in the Figure below (SEQ ID NOs: 68 and 69, respectively).

## Listeria hly DP-L4056 and EGD Alignment

Query: Listeria EGD  
Subject: DP-L4056 (wild-type, Portnoy strain)

```

Query: 1      ggtacctctctttgattagtagtattcctatctttaaagtgacttttatgttgaggcattaac 60
               |||
Sbjct: 1      ggtacctctctttgattagtagtattcctatctttaaagttacttttatgttgaggcattaac 60

Query: 61      atttggtaaagcagcagataaaagggacagcaggactagaataaagctataaaagcaagcatata 120
               |||
Sbjct: 61      atttggtaaatgacgtcaaaaggatagcaagactagaataaagctataaaagcaagcatata 120

Query: 121     atattgcgtttcactcttttagaagcgaatttcgccaatattataaattatcaaaagagaggg 180
               |||
Sbjct: 121     atattgcgtttcactcttttagaagcgaatttcgccaatattataaattatcaaaagagaggg 180

Query: 181     gtggcaaacggtatttggcattattaggttaaaaaatgtagaaggagagtgaaacccatg 240
               |||
Sbjct: 181     gtggcaaacggtatttggcattattaggttaaaaaatgtagaaggagagtgaaacccatg 240
               Shine-Dalgarno LLO start

```

[00349] The 422 bp DNA corresponding to the hly promoter and secA1 LLO signal peptide were liberated from the pCR-XL-TOPO-hly promoter plasmid clone by digestion with *Kpn I* and *Bam HI*, and cloned into the pPL2 plasmid vector (Lauer *et al.*, 2002, *J. Bact.*), according to conventional methods well-known to those skilled in the art. This plasmid is known as pPL2-hlyP (native).

### 6.2.5 Cloning and Insertion of EphA2 into pPL2 vectors for expression in selected recombinant *Listeria monocytogenes* strains

[00350] The external (EX2) and cytoplasmic (CO) domains of EphA2 which flank the EphA2 transmembrane helix were cloned separately for insertion into various pPL2-signal peptide expression constructs. Genes corresponding to the native mammalian sequence or codon-optimized for expression in *Listeria monocytogenes* of EphA2 EX2 and CO domains were used. The optimal codons in *Listeria* (see table, *ibid*) for each of the 20 amino acids were utilized for codon-optimized EphA2 EX2 and EphA2 CO. The codon-optimized EphA2 EX2 and CO domains were synthesized by extension of overlapping oligonucleotides, using techniques common to those skilled in the art. The expected sequence of all synthesized EphA2 constructs was verified by nucleotide sequencing.

[00351] SEQ ID NOS:23, 21 and 22 represent the primary amino acid sequences, together with the native and codon-optimized nucleotide sequences, respectively, for the EX2 domain of EphA2.

[00352] SEQ ID NOS: 34, 32 and 33 represent the primary amino acid sequences, together with the native and codon-optimized nucleotide sequences, respectively, for the CO domain of EphA2.

[00353] Additionally, FLAG (Stratagene, La Jolla, CA) and myc epitope tags were inserted, respectively, in-frame at the amino and carboxy termini of synthesized EphA2

EX2 and CO genes for detection of expressed and secreted EphA2 by Western blot analysis using antibodies specific for the FLAG or proteins. Thus, the expressed protein had the following ordered elements: NH<sub>2</sub>-Signal Peptide-FLAG-EphA2-myc-CO<sub>2</sub>. Shown below are the FLAG and myc epitope tag amino acid and codon-optimized nucleotide sequences.

- 5 [00354] FLAG
- [00355] 5'-GATTATAAAGATGATGATGATAAA (SEQ ID NO:51)
- [00356] NH<sub>2</sub>-DYKDDDDK-CO<sub>2</sub> (SEQ ID NO:52)
- [00357] Myc
- [00358] 5'-GAACAAAAATTAATTAGTGAAGAAGATTTA (SEQ ID NO:53)
- 10 [00359] NH<sub>2</sub>-EQKLISEEDL-CO<sub>2</sub> (SEQ ID NO:54)

#### **6.2.6 Detection of synthesized and secreted heterologous proteins by Western blot analysis**

- [00360] Synthesis of EphA2 protein and secretion from various selected recombinant *Listeria*-EphA2 strains was determined by Western blot analysis of trichloroacetic acid
- 15 (TCA) precipitated bacterial culture fluids. Briefly, mid-log phase cultures of *Listeria* grown in BHI media were collected in a 50 mL conical centrifuge tube, the bacteria were pelleted, and ice-cold TCA was added to a final [6%] concentration to the bacterial culture supernatant and incubated on ice minimally for 90 min or overnight. The TCA-precipitated proteins were collected by centrifugation at 2400 X g for 20 min at 4°C. The pellet was
- 20 then resuspended in 300-600 µl volume of TE, pH 8.0 containing 15 µg/ml phenol red. Sample dissolution was facilitated by vortexing. Sample pH was adjusted by NH<sub>4</sub>OH addition if necessary until color was pink. All samples were prepared for electrophoresis by addition of 100 µl of 4X SDS loading buffer and incubating for 10 min. at 90°C. The samples were then centrifuged from 5 min at 14,000 rpm in a micro-centrifuge, and the
- 25 supernatants collected and stored at -20°C. For Western bolt analysis, 20 µl of prepared fractions (the equivalent of culture fluids from of 1-4 x 10<sup>9</sup> bacteria), were loaded on the 4-12% SDS-PAGE gel, electrophoresed, and the proteins were transferred to PDDF membrane, according to common methods used by those skilled in the art. Transferred membranes were prepared s for incubation with antibody, by incubating in 5% dry milk in
- 30 PBS for 2 hr. at room temperature with agitation. Antibodies were used under the following dilutions in PBST buffer (0.1% Tween 20 in PBS): (1) Rabbit anti-Myc polyclonal antibody (ICL laboratories, Newberg, Oregon) at 1:10,000; (2) murine anti-FLAG monoclonal antibody (Stratagene, *ibid*) at 1:2,000; and, (3) Rabbit anti-EphA2 (carboxy terminus-specific) polyclonal antibody (sc-924, Santa Cruz Biotechnology, Inc.,
- 35 Santa Cruz, CA). Specific binding of antibody to protein targets was evaluated by

secondary incubation with goat anti-rabbit or anti-mouse antibody conjugated with horseradish peroxidase and detection with the ECL chemiluminescence assay kit (Amersham), and exposure to film.

## 6.2.7 Secretion of EphA2 protein by recombinant *Listeria* encoding various forms of EphA2

### 6.2.7.1. *Listeria*: [strains DP-L4029 (*actA*) or DP-L4017 (LLO L461T)]

[00361] Expression cassette construct: LLOss-PEST-CO-EphA2 (SEQ ID NO:35)

[00362] The native sequence of the EphA2 CO domain was genetically fused to the native secA1 LLO sequence, and the heterologous antigen expression cassette under control of the *Listeria hly* promoter was inserted into the pPL2 plasmid between the *Kpn I* and *Sac I* sites as described (*ibid*). The pPL2-EphA2 plasmid constructs were introduced by conjugation into the *Listeria* strains DP-L4029 (*actA*) and DP-L4017 (L461T LLO) as described (*ibid*). Figure 2 shows the results of a Western blot analysis of TCA-precipitated bacterial culture fluids of 4029-EphA2 CO and 4017-EphA2 CO. This analysis demonstrated that recombinant *Listeria* engineered to contain a heterologous protein expression cassette comprised of native sequences corresponding to the secA1 and EphA2 CO fusion protein secreted multiple EphA2-specific fragments that were lower than the 52 kDa expected molecular weight, demonstrating the need for modification of the expression cassette.

### 6.2.7.2. *Listeria*: [DP-L4029 (*actA*)]

[00363] Expression cassette constructs:

Native LLOss-PEST-FLAG-EX2\_EphA2-myc-CodonOp (SEQ ID NO:26)

(CodonOp) LLOss-PEST-(CodonOp)FLAG-EX2\_EphA2-myc (SEQ ID NO:28)

[00364] The native secA1 LLO signal peptide sequence or secA1 LLO signal peptide sequence codon-optimized for expression in *Listeria* was fused genetically with the EphA2 EX2 domain sequence codon-optimized for expression in *Listeria*, and the heterologous antigen expression cassette under control of the *Listeria hly* promoter was inserted into the pPL2 plasmid between the *Kpn I* and *Sac I* sites as described (*ibid*). The pPL2-EphA2 plasmid constructs were introduced by conjugation into the *Listeria* strain DP-L4029 (*actA*) as described (*ibid*). Figure 3 shows the results of a Western blot analysis of TCA-precipitated bacterial culture fluids of *Listeria actA* encoding either the native or codon-optimized secA1 LLO signal peptide fused with the codon-optimized EphA2 EX2 domain.

This analysis demonstrated that the combination of utilizing sequence for both signal peptide and heterologous protein optimized for the preferred codon usage in *Listeria monocytogenes* resulted in expression of the expected full-length EphA2 EX2 domain protein. Expression of full-length EphA2 EX2 domain protein was poor with codon-optimization of the EphA2 coding sequence alone. The level of heterologous protein expression (fragmented or full-length) was highest when utilizing the *Listeria monocytogenes* LLO secA1 signal peptide, codon-optimized for expression in *Listeria monocytogenes*.

#### 6.2.7.3. *Listeria*: [DP-L4029 (*actA*)]

- 10 [00365] Expression cassette constructs:
- Native LLOss-PEST-(CodonOp) FLAG-EphA2\_CO-myc (SEQ ID NO:37)
- CodonOp LLOss-PEST-(CodonOp) FLAG- EphA2\_CO-myc (SEQ ID NO:39)
- 15 CodonOp PhoD-(CodonOp) FLAG- EphA2\_CO-myc (SEQ ID NO:41)
- [00366] The native secA1 LLO signal peptide sequence or the secA1 LLO signal peptide sequence codon-optimized for expression in *Listeria*, or, alternatively, the Tat signal peptide of the phoD gene from *Bacillus subtilis* codon-optimized for expression in *Listeria*, was fused genetically with the EphA2 CO domain sequence codon-optimized for
- 20 expression in *Listeria*, and the heterologous antigen expression cassette under control of the *Listeria hly* promoter was inserted into the pAM401-MCS plasmid between the *Kpn I* and *Sac I* sites as described (*ibid*). The pAM401-EphA2 plasmid constructs were introduced by electroporation into the *Listeria* strain DP-L4029 (*actA*) as described (*ibid*). Figure 4 shows the results of a Western blot analysis of TCA-precipitated bacterial culture fluids of *Listeria*
- 25 *actA* encoding either the native or codon-optimized secA1 LLO signal peptide, or codon-optimized *Bacillus subtilis phoD* Tat signal peptide fused with the codon-optimized EphA2 CO domain. This analysis demonstrated once again that the combination of utilizing sequence for both signal peptide and heterologous protein optimized for the preferred codon usage in *Listeria monocytogenes* resulted in expression of the expected full-length EphA2
- 30 CO domain protein. Furthermore, expression and secretion of the expected full-length EphA2 CO domain protein resulted from recombinant *Listeria* encoding codon-optimized *Bacillus subtilis phoD* Tat signal peptide fused with the codon-optimized EphA2 CO domain. This result demonstrates the novel and unexpected finding that signal peptides from distinct bacterial species can be utilized to program the secretion of heterologous
- 35 proteins from recombinant *Listeria*. Expression of full-length EphA2 CO domain protein



was poor with codon-optimization of just the EphA2 sequence. The level of heterologous protein expression was highest when utilizing signal peptides codon-optimized for expression in *Listeria monocytogenes*.

#### **6.2.8 Construction of *Listeria* strains expressing AH1/OVA or AH1-A5/OVA**

[00367] Mutant *Listeria* strains expressing a truncated form of a model antigen ovalbumin (OVA), the immunodominant epitope from mouse colorectal cancer (CT26) known as AH1 (SPSYVYHQF) (SEQ ID NO:55), and the altered epitope AH1-A5 (SPSYAYHQF (SEQ ID NO:56); Slansky *et al.*, 2000, *Immunity*, 13:529-538) were prepared. The pPL2 integrational vector (Lauer *et al.*, *J. Bacteriol.* 184:4177 (2002); U.S. Patent Publication No. 2003/0203472) was used to derive OVA and AH1-A5/OVA recombinant *Listeria* strains containing a single copy integrated into an innocuous site of the *Listeria* genome.

#### **6.2.9 Construction of OVA-expressing *Listeria* (DP-L4056)**

[00368] An antigen expression cassette consisting of hemolysin-deleted LLO fused with truncated OVA and contained in the pPL2 integration vector (pPL2/LLO-OVA) is first prepared. The *Listeria*-OVA vaccine strain is derived by introducing pPL2/LLO-OVA into the phage-cured *L. monocytogenes* strain DP-L4056 at the PSA (Phage from ScottA) attachment site tRNA<sup>Arg</sup>-attBB<sup>3</sup>.

[00369] PCR is used to amplify the hemolysin-deleted LLO using the following template and primers:

Source: DP-L4056 genomic DNA

Primers:

Forward (*KpnI*-LLO nts. 1257-1276):

5'-CTCTGGTACCTCCTTTGATTAGTATATTC (SEQ ID NO:57)  
(T<sub>m</sub>:LLO-spec: 52°C. Overall: 80°C.)

Reverse (*BamHI*-*XhoI*-LLO nts. 2811-2792):

5'-CAATGGATCCCTCGAGATCATATAATTTACTTCATCCC  
(SEQ ID NO:58)

(T<sub>m</sub>:LLO-spec: 52°C. Overall: 102°C)

[00370] PCR is also used to amplify the truncated OVA using the following template and primers:

Source: pDP3616 plasmid DNA from DP-E3616 *E. coli* (Higgins *et al.*, *Mol. Molbiol.* 31:1631-1641 (1999)).

Primers:

Forward (*XhoI*- *NcoI* OVA cDNA nts. 174-186):

5'-ATTTCTCGAGTCCATGGGGGTTCTCATCATC  
(SEQ ID NO:59)

(T<sub>m</sub>: OVA-spec: 60°C. Overall: 88°C.)

Reverse (*XhoI*-*NotI*-*HindIII*):

5'-GGTGCTCGAGTGCGGCCGCAAGCTT  
(SEQ ID NO:60)

(T<sub>m</sub>: Overall: 82°C.)

[00371] One protocol for completing the construction process involves first cutting the LLO amplicon with *KpnI* and *BamHI* and inserting the *KpnI/BamHI* vector into the pPL2 vector (pPL2-LLO). The OVA amplicon is then cut with *XhoI* and *NotI* and inserted into the pPL2-LLO which has been cut with *XhoI/NotI*. (Note: The pPL2 vector does not contain any *XhoI* sites; pDP-3616 contains one *XhoI* site, that is exploited in the OVA reverse primer design.) The construct pPL2/LLO-OVA is verified by restriction analysis (*KpnI*-LLO-*XhoI*-OVA-*NotI*) and sequencing. The plasmid pPL2/LLO-OVA is introduced into *E. coli* by transformation, followed by introduction and integration into *Listeria* (DP-L4056) by conjugation, exactly as described by Lauer et al. (or into another desired strain of *Listeria*).

#### 6.2.10 Construction of *Listeria* strains expressing AH1/OVA or AH1-A5/OVA

[00372] To prepare *Listeria* expressing either the AH1/OVA or the AH1-A5/OVA antigen sequences, inserts bearing the antigen are first prepared from oligonucleotides and then ligated into the vector pPL2-LLO-OVA (prepared as described above).

[00373] The following oligonucleotides are used in preparation of the AH1 or AH1-A5 insert:

*AH1 epitope insert (ClaI-PstI compatible ends):*

Top strand oligo (AH1 Top):

5'-CGATTCCCCTAGTTATGTTTACCACCAATTGCTGCA  
(SEQ ID NO:61)

Bottom strand oligo (AH1 Bottom):

5'-GCAAAATTGGTGGTAAACATAACTAGGGGAAT  
(SEQ ID NO:62)

*AH1-A5 epitope insert (ClaI-AvaII compatible ends):*

[00374] The sequence of the AH1-A5 epitope is SPSYAYHQF (SEQ ID NO:56) (5'-AGT CCA AGT Tat GCA Tat CAT CAA TTT-3') (SEQ ID NO:63).

Top: 5'-CGATAGTCCAAGTTATGCATATCATCAATTTCG  
(SEQ ID NO:64)

Bottom: 5'-GTCGCAAATTGATGATATGCATAACTTGGACTAT  
(SEQ ID NO:65)

- 5 [00375] The oligonucleotide pair for a given epitope are mixed together at an equimolar ratio, heated at 95 °C for 5 min. The oligonucleotide mixture is then allowed to slowly cool. The annealed oligonucleotide pairs are then ligated at a 200 to 1 molar ratio with pPL2-LLO/OVA plasmid prepared by digestion with the relevant restriction enzymes. The identity of the new construct can be verified by restriction analysis and/or sequencing.
- 10 [00376] The plasmid can then be introduced into *E. coli* by transformation, followed by introduction and integration into *Listeria* (DP-L4056) by conjugation, exactly as described by Lauer et al., or into another desired strain of *Listeria*, such as an *actA*<sup>-</sup> mutant strain (DP-L0429), LLO L461T strain (DP-L4017), or *actA*<sup>-</sup>/*inlB*<sup>-</sup> strain (DP-L4029*inlB*).

### 6.3 EXAMPLE 3: GENERATION OF MURINE TUMOR CELL LINES THAT EXPRESS HUMAN EphA2

#### 6.3.1 Background

[00377] A mouse immunotherapy model was created for testing the *Listeria*-based vaccines of the invention. Three murine tumor cell lines, the CT26 murine colon carcinoma cell line, the B16F10 murine melanoma cell line, and the RenCa murine renal cell carcinoma cell line were created to express high levels of the huEphA2 protein. FACS cell sorting assays were performed to identify CT26, B16F10, and RenCa tumor cells expressing high levels of huEphA2, which were pooled and analyzed by Western blot analysis. Clones were further pooled by FACS cell sorting to generate subclones expressing the highest levels of huEphA2.

#### 6.3.2 Selection of CT26 Murine Colon Carcinoma Cells Expressing High Levels of huEphA2

##### 6.3.2.1. Transfection Assays With Lipofectamine<sup>TM</sup>

[00378] CT26 cells were transfected with constructs containing huEphA2 using standard transfection techniques and commercially available Lipofectamine<sup>TM</sup> according to the manufacturer's instructions.

##### 6.3.2.2. Flow Cytometry (FACS) Analysis

[00379] Single cell FACS sorting assays were performed by standard techniques to identify CT26 murine carcinoma tumor cell expressing high levels of human EphA2.

[00380] Figure 5 illustrates a representative experiment, showing that the EphA2-3 clone expressed the highest levels of human EphA2 protein.

### **6.3.2.3. Western Blot of Pooled Populations Expressing High Levels of huEphA2**

5 [00381] Western blotting was also performed using standard techniques to determine the levels of human EphA2 protein expression in CT26 cells following FACS sorting of pooled populations of cells transfected with various constructs containing the huEphA2 gene. Figure 6 depicts results of a representative experiment. Compared to various clones tested, the huEphA2-3 clone expressed the highest levels of human EphA2 protein and was  
10 selected for the *in vivo* efficacy studies described below. Cells were further pooled to generate subclones expressing the highest levels of huEphA2.

### **6.3.3 Selection of B16F10 Murine Melanoma Cells Expressing High Levels of huEphA2**

#### **6.3.3.1. Retroviral Transduction**

15 [00382] Human EphA2 was introduced into B16F10 murine melanoma cells by a retroviral transduction method to create clones expressing high levels of the protein.

#### **6.3.3.2. Flow Cytometry (FACS) Analysis**

[00383] As was performed on the CT26 cells, single cell FACS sorting assays were performed by standard techniques on B16F10 cells expressing huEphA2 to generate clones  
20 expressing high levels of huEphA2. Clones expressing the highest levels of huEphA2 were pooled and further examined by Western blot analysis. A representative FACS experiment is depicted in Figure 7, showing a B16F10 subclone expressing high levels of huEphA2.

### **6.3.3.3. Western Blot of Pooled Populations Expressing High Levels of huEphA2**

25 [00384] Western blotting was also performed as described above to determine levels of huEphA2 protein expression in B16F10 cells following FACS sorting of pooled populations of cells containing the huEphA2 gene introduced by retroviral transduction. Cells were further pooled to generate subclones expressing the highest levels of huEphA2.

### **6.3.4 Selection of RenCa Murine Renal Cell Carcinoma Cells Expressing High Levels of huEphA2**

#### **6.3.4.1. Transfection Assays With Lipofectamine™**

[00385] RenCa cells were transfected with constructs containing huEphA2 using  
5 standard transfection techniques and commercially available Lipofectamine™ according to the manufacturer's instructions.

#### **6.3.4.2. Flow Cytometry (FACS) Analysis**

[00386] Single cell FACS sorting assays were performed by standard techniques to identify RenCa renal cell carcinoma tumor cells expressing high levels of human EphA2.

#### **6.3.4.3. Western Blot of Pooled Populations Expressing High Levels of huEphA2**

[00387] Western blotting was also performed using standard techniques to determine the levels of human EphA2 protein expression in RenCa cells following FACS sorting of  
10 pooled populations of cells transfected with various constructs containing the huEphA2  
15 gene. Cells were further pooled to generate subclones expressing the highest levels of huEphA2.

### **6.3.5 Transfection of 293 Cells with pCDNA4 plasmids encoding full-length EphA2**

[00388] Expression cassette constructs:

20 [00389] pCDNA4-EphA2

[00390] The native full-length EphA2 gene was cloned into the eukaryotic CMV promoter-based expression plasmid pCDNA4 (Invitrogen, Carlsbad, CA). Figure 8 shows the results of a Western blot analysis of lysates prepared from 293 cells transfected with the pCDNA4-EphA2 plasmid, and demonstrates the abundant expression in mammalian cells  
25 of full-length EphA2 protein.

### **6.4 EXAMPLE 4: Assessment of antigen-specific immune responses after vaccination.**

[00391] The vaccines of the present invention can be assessed using a variety of *in vitro* and *in vivo* methods. Some assays involve the analysis of antigen-specific T cells  
30 from the spleens of mice that have been vaccinated. For example C57Bl/6 or Balb/c are vaccinated by intravenous injection of 0.1 LD<sub>50</sub> of a *Listeria* strain expressing OVA (or other appropriate antigen). Seven days after the vaccination, the spleen cells of the mice are harvested (typically 3 mice per group) by placing the spleens into ice cooled RPMI 1640 medium and preparing a single cell suspension from this. As an alternative, the lymph

nodes of the mice could be similarly harvested, prepared as a single cell suspension and substituted for the spleen cells in the assays described below. Typically, spleen cells are assessed for intravenous or intraperitoneal administration of the vaccine while spleen cells and cells from lymph nodes are assessed for intramuscular, subcutaneous or intradermal administration of the vaccine.

**[00392]** Unless otherwise noted, all antibodies used in these examples can be obtained from Pharmingen, San Diego, CA.

#### **6.4.1 ELISPOT Assay:**

- [00393]** Using a *Listeria* strain having an OVA antigen as an example, the quantitative frequency of antigen-specific T cells generated upon immunization in a mouse model is assessed using an ELISPOT assay. The antigen-specific T cells evaluated are OVA specific CD8+ or LLO specific CD8+ or CD4+ T cells. This OVA antigen model assesses the immune response to a heterologous tumor antigen inserted into the vaccine and could be substituted with any antigen of interest. The LLO antigen is specific to *Listeria*.
- The specific T cells are assessed by detection of cytokine release (e.g. IFN- $\gamma$ ) upon recognition of the specific antigen. PVDF-based 96 well plates (BD Biosciences, San Jose, CA) are coated overnight at 4°C with an anti-murine IFN- $\gamma$  monoclonal antibody (mAb R4; 5  $\mu$ g/ml). The plates are washed and blocked for 2 hours at room temperature with 200  $\mu$ L of complete RPMI. Spleen cells from vaccinated mice (or non vaccinated control mice) are added at  $2 \times 10^5$  cells per well and incubated for 20 to 22 hours at 37°C in the presence of various concentrations of peptides ranging from 0.01 to 10  $\mu$ M. The peptides used for OVA and LLO are either SL8, an MHC class I epitope for OVA, LLO<sub>190</sub> (NEKYAQAYPNVS, Invitrogen) an MHC class II epitope for listeriolysin O (*Listeria* antigen), LLO<sub>296</sub> (VAYGRQVYL), an MHC class I epitope for listeriolysin O, or LLO<sub>91</sub> (GYKDGNEYI), an MHC class I epitope for listeriolysin O. LLO<sub>190</sub> and LLO<sub>296</sub> are used in a C57Bl/6 model, while LLO<sub>91</sub> is used in a Balb/c model. After washing, the plates are incubated with secondary biotinylated antibodies specific for IFN- $\gamma$  (XMG1.2) diluted in PBS to 0.5 $\mu$ g/ml. After incubation at room temperature for 2 hours, the plates are washed and incubated for 1 hour at 37 °C with a 1 nm gold goat anti-biotin conjugate (GAB-1; 1:200 dilution; Ted Pella, Redding, CA) diluted in PBS containing 1 % BSA. After thorough washing, the plates are incubated at room temperature for 2 to 10 minutes with substrate (Silver Enhancing Kit; 30 ml/well; Ted Pella) for spot development. The plates are then rinsed with distilled water to stop the substrate reaction. After the plates have been air-dried, spots in each well are counted using an automated ELISPOT plate reader (CTL, Cleveland, OH). The cytokine response is expressed as the number of IFN- $\gamma$  spot-forming

cells (SFCs) per  $2 \times 10^5$  spleen cells for either the OVA specific T cells or the *Listeria* specific T cells.

#### **6.4.2 Intracellular Cytokine Staining Assay (ICS):**

[00394] In order to further assess the number of antigen-specific CD8+ or CD4+ T  
5 cells and correlate the results with those obtained from ELISPOT assays, ICS is performed  
and the cells evaluated by flow cytometry analysis. Spleen cells from vaccinated and  
control groups of mice are incubated with SL8 (stimulates OVA specific CD8+ cells) or  
LLO<sub>190</sub> (stimulates LLO specific CD4+ cells) for 5 hours in the presence of Brefeldin A  
(Pharmingen). The Brefeldin A inhibits secretion of the cytokines produced upon  
10 stimulation of the T cells. Spleen cells incubated with an irrelevant MHC class I peptide are  
used as controls. PMA (phorbol-12-myristate-13-acetate, Sigma) 20 ng/ml and ionomycin  
(Sigma) 2 µg/ml stimulated spleen cells are used as a positive control for IFN-γ and TNF-α  
intracellular cytokine staining. For detection of cytoplasmic cytokine expression, cells are  
15 stained with FITC-anti-CD4 mAb (RM 4-5) and PerCP-anti-CD8 mAb (53-6.7), fixed and  
permeabilized with Cytofix/CytoPerm solution (Pharmingen), and stained with PE-  
conjugated anti-TNF-α mAb (MP6-XT22) and APC-conjugated anti-IFN-γ mAb (XMG1.2)  
for 30 minutes on ice. The percentage of cells expressing intracellular IFN-γ and/or TNF-α  
was determined by flow cytometry (FACScalibur, Becton Dickinson, Mountain View, CA)  
and data analyzed using CELLQuest software (Becton Dickinson Immunocytometry  
20 System). As the fluorescent labels on the various antibodies can all be distinguished by the  
FACScalibur, the appropriate cells are identified by gating for those CD8+ and CD4+ that  
are stained with either or both of the anti-IFN-γ or anti-TNF-α.

#### **6.4.3 Cytokine Expression of Stimulated Spleen Cells:**

[00395] The level of cytokine secretion by the spleen cells of mice can also be  
25 assessed for control and vaccinated C57Bl/6 mice. Spleen cells are stimulated for 24 hours  
with SL8 or LLO<sub>190</sub>. Stimulation with irrelevant peptide HSV-gB<sup>2</sup> (Invitrogen,  
SSIEFARL) is used as a control. The supernatants of the stimulated cells are collected and  
the levels of T helper-1 and T helper 2 cytokines are determined using an ELISA assay  
(eBiosciences, CO) or a Cytometric Bead Array Kit (Pharmingen).

#### **6.4.4 Assessment of Cytotoxic T cell Activity:**

[00396] The OVA specific CD8+ T cells can be further evaluated by assessing their  
cytotoxic activity, either *in vitro* or directly in C57Bl/6 mouse *in vivo*. The CD8+ T cells  
recognize and lyse their respective target cells in an antigen-specific manner. *In vitro*  
cytotoxicity is determined using a chromium release assay. Spleen cells of naïve and

*Listeria*-OVA (internal) vaccinated mice are stimulated at a 10:1 ratio with either irradiated EG7.OVA cells (EL-4 tumor cell line transfected to express OVA, ATCC, Manassas, VA) or with 100 nM SL8, in order to expand the OVA specific T cells in the spleen cell population. After 7 days of culture, the cytotoxic activity of the effector cells is determined in a standard 4-hour <sup>51</sup>Cr-release assay using EG7.OVA or SL8 pulsed EL-4 cells (ATCC, Manassas, VA) as target cells and EL-4 cells alone as negative control. The YAC-1 cell line (ATCC, Manassas, VA) is used as targets to determine NK cell activity, in order to distinguish the activity due to T cells from that due to NK cells. The percentage of specific cytotoxicity is calculated as 100 x (experimental release – spontaneous release) / (maximal release – spontaneous release). Spontaneous release is determined by incubation of target cells without effector cells. Maximal release is determined by lysing cells with 0.1% Triton X-100. Experiments are considered valid for analysis if spontaneous release is < 20% of maximal release.

**[00397]** For the assessment of cytotoxic activity of OVA-specific CD8+ T cells *in vivo*, spleen cells from naïve C57Bl/6 mice are split into two equivalent aliquots. Each group is pulsed with a specific peptide, either target (SL8) or control (HSV-gB<sup>2</sup>), at 0.5 µg/ml for 90 minutes at 37 °C. Cells are then washed 3 times in medium, and twice in PBS + 0.1% BSA. Cells are resuspended at 1 x 10<sup>7</sup> per ml in warm PBS + 0.1% BSA (10 ml or less) for labeling with carboxyfluorescein diacetate succinimidyl ester (CFSE, Molecular Probes, Eugene, OR). To the target cell suspension, 1.25 µL of a 5mM stock of CFSE is added and the sample mixed by vortexing. To the control cell suspension, a ten-fold dilution of the CFSE stock is added and the sample mixed by vortexing. The cells are incubated at 37 °C for 10 minutes. Staining is stopped by addition of a large volume (>40 ml) of ice-cold PBS. The cells are washed twice at room temperature with PBS, then resuspended and counted. Each cell suspension is diluted to 50 x 10<sup>6</sup> per ml, and 100 µL of each population is mixed and injected via the tail vein of either naïve or vaccinated mice. After 12-24 hours, the spleens are harvested and a total of 5 x 10<sup>6</sup> cells are analyzed by flow cytometry. The high (target) and low (control) fluorescent peaks are enumerated, and the ratio of the two is used to establish the percentage of target cell lysis. The *in vivo* cytotoxicity assay permits the assessment of lytic activity of antigen-specific T cells without the need of *in vitro* re-stimulation. Furthermore, this assay assesses the T cell function in their native environment.



## 6.5 EXAMPLE 5: IN VIVO EphA2 EFFICACY STUDIES

### 6.5.1 Background

[00398] Efficacy studies were performed in mice inoculated with CT26 tumor cells expressing the extracellular domain (ED) of human EphA2 in order to characterize the anti-tumor effect of huEphA2. Endpoints measured were tumor volume and percent survival of the mice after tumor inoculation. The routes of inoculation were subcutaneous (s.c.) and intravenous (i.v.). HBSS and *Listeria* were administered as controls.

### 6.5.2 Control Vaccinations With AH1-A5-Expressing *Listeria*

[00399] Balb/c mice (n=5) were immunized with 0.1 LD<sub>50</sub> *Listeria* 3 days post- i.v. inoculation of 1 x 10<sup>5</sup> CT26 cells. Figure 9A demonstrates that therapeutic immunization with *Listeria* expressing AH1-A5 increases survival of the inoculated animals. Figure 9B shows the result of a separate but otherwise equivalent experiment in which lungs of the experimental mice were harvested on Day 19 following cell inoculation and fixed. Gross assessment of lung nodules was also performed, demonstrating the absence of tumors in the lungs of test animals receiving *Listeria*-AH1/A5 as compared to control animals receiving a *Listeria* control.

### 6.5.3 Prophylactic EphA2 Vaccinations

#### 6.5.3.1. Effect of Immunization with *Listeria* Expressing ECD of huEphA2 on CT26-hEphA2 Tumor Growth and Survival

[00400] Preventive studies were performed utilizing a pool of CT26 cells expressing huEphA2 generated by the single cell FACS assays described above. Groups of ten Balb/c mice per group were inoculated s.c. and groups of five mice per group were inoculated i.v. with CT26 colon carcinoma cells transfected with human EphA2 ("CT26-hEphA2"). The mice were immunized with 0.1 LD<sub>50</sub> *Listeria* control or *Listeria* expressing the ECD of hEphA2 in a 200µl bolus. For the studies entailing s.c. inoculations with CD26, AH1/A5 *Listeria* were used as a positive control. The immunizations were performed 14 and 4 days prior to CT26-hEphA2 tumor challenge. Tumor volume measurements were obtained twice weekly for the course of the study to determine an anti-tumor effect of the

Vaccination Group	Tumor Volume (mm <sup>3</sup> ± s.e.m.) (Day 42)	P vs. HBSS	P vs. <i>Listeria</i> Control
HBSS	1202.9 (± 321)	-	0.5528
<i>Listeria</i> Control	945.5 (± 338)	0.5528	-
<i>Listeria</i> -AH1/A5	392.5 (± 225)	0.0471	0.1895
<i>Listeria</i> -hEphA2-ECD	0.0 (± 0.0)	0.0012	0.0118

**TABLE 4**

[00402] Figure 10B demonstrates the anti-tumor efficacy of *Listeria* expressing the ECD of hEphA2 against i.v. inoculations of huEphA2-expressing CT26 cells as compared to the negative controls \*p=0.0017). The data are summarized in Table 5 below:

Vaccination Group	Median Survival (Days)	P vs. HBSS	# Survivors (Day 65)
HBSS	18	-	0
<i>Listeria</i> Control	18	0.754	0
<i>Listeria</i> -AH1/A5	>65	0.0017	5
<i>Listeria</i> -hEphA2-ECD	>65	0.0017	3

**TABLE 5**

[00403] Preventive studies were performed according to the schedule described below. These studies utilized a pool of CT26 cells expressing huEphA2 generated by the single cell FACS assays described above.

[00404] Groups: Eight groups of ten mice per group. Groups 1-4 were inoculated s.c. and groups 5-8 were inoculated i.v. with CT26 colon carcinoma cells transfected with human EphA2, as shown in Table 6 below:

<u>Treatment Group</u>	<u>Number of Mice per Groups</u>
1. Control - HBSS	10
2. L4029 – control <i>Listeria monocytogenes</i>	10
3. L4029-EphA2 exFlag – <i>Listeria monocytogenes</i> expressing extracellular domain of human EphA2	10
4. L4029 – AH1 <i>Listeria monocytogenes</i>	10
5. Control - HBSS	10
6. L4029 – control <i>Listeria monocytogenes</i>	10
7. L4029-EphA2 exFlag – <i>Listeria monocytogenes</i> expressing extracellular domain of human EphA2	10
8. L4029 – AH1 <i>Listeria monocytogenes</i>	10

**TABLE 6**

[00405] Schedule: Animals received i.v. administrations of the agents listed above in 200µl bolus on Day 0 and Day 10. On Day 14, animals were inoculated with CT26 colon carcinoma cells transfected with human EphA2 (L4029EphA2-exFlag), *Listeria* control (L4029), or *Listeria* positive control containing the AH1 protein (L4029-AH1) (5 × 10<sup>5</sup> cells

in 100 $\mu$ l volume) either subcutaneously or intravenously (experimental lung metastases model). Tumor volume was measured bi-weekly (s.c inoculation only) and animal weights assessed on a weekly basis. Any animals possessing tumors greater than 2000 mm<sup>3</sup> or demonstrating signs of morbidity (hunched posture, impaired breathing, decreases mobility, greater than 20% weight loss, etc.) were humanely euthanized. The experimental schedule is summarized in Table 7 below:

Group	Cell Inoculation Route (5 x10 <sup>5</sup> cell in 100 $\mu$ l) (Day 14)	Primary Vaccination (Day 0)	Boost Vaccination (Day 10)
1. Control	s.c.	HBSS	HBSS
2. L4029	s.c.	2x10 <sup>7</sup> CFU	2x10 <sup>7</sup> CFU
3. L4029 EphA2-exFlag	s.c.	2x10 <sup>7</sup> CFU	2x10 <sup>7</sup> CFU
4. L4029 –AH1	s.c.	2x10 <sup>7</sup> CFU	2x10 <sup>7</sup> CFU
5. Control	i.v.	HBSS	HBSS
6. L4029	i.v.	2x10 <sup>7</sup> CFU	2x10 <sup>7</sup> CFU
7. L4029 EphA2-exFlag	i.v.	2x10 <sup>7</sup> CFU	2x10 <sup>7</sup> CFU
8. L4029 – AH1	i.v.	2x10 <sup>7</sup> CFU	2x10 <sup>7</sup> CFU

**TABLE 7**

**[00406]** In this study, vaccination with *Listeria*-huEphA2 exFlag demonstrated a significant anti-tumor effect in both the s.c. and experimental lung metastases models (i.v.). In the s.c. model, a significant reduction in tumor growth was achieved with 3 mice remaining tumor-free. This response was also specific compared to the control *Listeria* and vehicle treated animals. In the experimental lung metastases model, vaccination with *Listeria* huEphA2-exFlag also demonstrated efficacy compared to the vehicle treated group.

**[00407]** Figures 11A-11D illustrate results of the preventive experiments. Figure 11A shows that the tumor volume of mice inoculated with CT26 cells expressing the ECD of huEphA2 was significantly reduced when compared to vehicle (HBSS), *Listeria* (L4029) and *Listeria* positive (L4029-AH1) controls starting at day 21 and continued until day 32 post inoculation. Figure 11B also depicts results of the preventive experiments, showing again that the tumor volume of mice inoculated with CT26 cells expressing the ECD of huEphA2 (L4029-EphA2 exFlag) was significantly reduced when compared to the *Listeria* (L4029) control starting at day 21 and continued until day 32 post inoculation. Figure 11C illustrates the results of the prevention study in the s.c. model, measuring percent survival of the mice post CT26 tumor cell inoculation. Compared to all control groups, the L4029-EphA2 exFlag group had the most significant survival rate (indicated by triangles). Figure 11D illustrates the results of the prevention study in the lung metastases model, measuring

the percent survival of the mice post tumor cell inoculation. Compared to all control groups, the L4029-EphA2 exFlag group had the most significant survival rate.

[00408] The foregoing data demonstrate that preventative immunization with *Listeria* expressing the ECD of hEphA2 suppresses CT26-hEphA2 tumor growth and increases survival.

#### **6.5.3.2. Effect of Immunization with Listeria Expressing ICD of huEphA2 on the Survival of Mice Inoculated with RenCa-hEphA2**

[00409] Preventive studies were performed utilizing a pool of RenCa cells (American Type Culture Collection, Manassas, VA) expressing huEphA2 generated and screened by the methods described above. Groups of ten Balb/c mice per group were inoculated subcutaneously with RenCa renal cell carcinoma cells expressing human EphA2 ("RenCa-hEphA2 cells"). The mice were immunized with 0.1 LD50 *Listeria* control or *Listeria* expressing the ICD of hEphA2 in a 200µl bolus. The immunizations were performed 18 and 4 days prior to RenCa-hEphA2 cell tumor challenge. Tumor volume measurements were obtained twice weekly for the course of the study to determine an anti-tumor effect of the vaccinations.

[00410] Figure 12 demonstrates the anti-tumor efficacy of *Listeria* expressing the ICD of hEphA2 against s.c. inoculations of huEphA2-expressing RenCA cells as compared to the negative controls. A significant anti-tumor response, as assessed by increased survival via Kaplan-Meier analysis, was observed in animals vaccinated with *Listeria* expressing the ICD of hEphA2 as compared to animals vaccinated with *Listeria* alone (\*p=0.0079).

#### **6.5.4 Therapeutic EphA2 Vaccinations**

[00411] Therapeutic studies were performed utilizing a pool of CT26 cells expressing huEphA2 generated by the single cell FACS assays described above.

[00412] A representative therapeutic study was performed as follows:

[00413] Groups: Six groups of ten mice per group. Groups 1-3 were inoculated s.c. and groups 4-6 were inoculated i.v. with CT26 murine colon carcinoma cells, as shown in

Table 8 below:

<b>Treatment Group</b>	<b>Number of Mice per Groups</b>
1. Control - HBSS	10
2. L4029 – control <i>Listeria monocytogenes</i>	10
3. L4029-EphA2 exFlag – <i>Listeria monocytogenes</i> expressing extracellular domain of human EphA2	10
4. Control - HBSS	10
5. L4029 – control <i>Listeria monocytogenes</i>	10
6. L4029-EphA2 exFlag – <i>Listeria monocytogenes</i> expressing extracellular domain of human EphA2	10

**TABLE 8**

**[00414]** Schedule: Animals were inoculated with CT26 colon carcinoma cells transfected with human EphA2 (L4029-EphA2 exFlag), *Listeria* control (L4029-control) or vehicle (HBSS) ( $5 \times 10^5$  cells in 100ml volume) either subcutaneously or intravenously (experimental lung metastases model). Three days after cell inoculation, animals received i.v. administrations of the agents listed above in 200ml bolus. Two weeks following the first administration, the animals received a booster vaccination. Tumor volume was measured bi-weekly (s.c inoculation only) and animal weights assessed on a weekly basis. Any animals possessing tumors greater than 2000 mm<sup>3</sup> or demonstrating signs of morbidity (hunched posture, impaired breathing, decreases mobility, greater than 20% weight loss, etc.) were humanely euthanized. The schedule is summarized in Table 9 below.

<b>Group</b>	<b>Cell Inoculation Route</b> ( $5 \times 10^5$ cell in 100μl)	<b>Primary Vaccination</b> (Day 3)	<b>Boost Vaccination</b> (Day 17)
1. Control	s.c.	HBSS	HBSS
2. L4029	s.c.	$6 \times 10^5$ to $2 \times 10^7$ CFU	$6 \times 10^5$ to $2 \times 10^7$ CFU
3. L4029 EphA2-exFlag	s.c.	$6 \times 10^5$ to $2 \times 10^7$ CFU	$6 \times 10^5$ to $2 \times 10^7$ CFU
4. Control	i.v.	HBSS	HBSS
5. L4029	i.v.	$6 \times 10^5$ to $2 \times 10^7$ CFU	$6 \times 10^5$ to $2 \times 10^7$ CFU
6. L4029 EphA2-exFlag	i.v.	$6 \times 10^5$ to $2 \times 10^7$ CFU	$6 \times 10^5$ to $2 \times 10^7$ CFU

**TABLE 9**

**[00415]** Figures 13A-13C illustrate the results of a typical therapeutic study. In Figure 13A, tumor volume was measured at several intervals post inoculation. Compared to the HBSS and *Listeria* controls, the mice inoculated with CT26 cells expressing the ECD of huEphA2 had a significantly lower tumor volume after day 14 and continued onto day 28. Figure 13B depicts the mean tumor volume of mice inoculated with CT26 cells containing either *Listeria* control or huEphA2. Compared to control, the mice inoculated

with CT26 cells expressing huEphA2 had a reduced mean tumor volume. Figure 13C represents the results of the therapeutic study using the lung metastases model, measuring percent survival of the mice post inoculation with CT26 cells with either HBSS or *Listeria* control, or *Listeria* expressing the ECD of huEphA2. Animals inoculated with CT26 cells expressing the ECD of huEphA2 (depicted by triangles) showed a higher percent survival rate compared to controls.

[00416] In another study, groups of ten Balb/c mice per group were inoculated s.c. or i.v. with CT26 colon carcinoma cells transfected with human EphA2 ("CT26-hEphA2"). The mice were immunized with 0.1 LD<sub>50</sub> actA *Listeria* control or *Listeria* expressing the ICD of hEphA2 in a 200µl bolus. In one regimen, the immunizations were performed 6 and 14 days post s.c. CT26-hEphA2 tumor inoculation. In another regimen, the immunizations were performed 3 and 14 days post i.v. CT26-hEphA2 tumor inoculation. Anti-tumor efficacy was determined from twice weekly tumor measurements and survival.

[00417] Significant anti-tumor efficacy was observed in the *Listeria*-hEphA2 vaccinated animals (p=0.0035).

[00418] Figure 14A demonstrates the tumor measurements of immunized animals. This data is summarized in Table 10 below:

Vaccination Group	Tumor Volume (mm <sup>3</sup> ± s.e.m.) (Day 21)	P vs. HBSS	P vs. <i>Listeria</i> Control
HBSS	1827 (± 518)	-	0.961
<i>Listeria</i> Control	1799 (± 267)	0.961	-
<i>Listeria</i> -AH1/A5	0	0.0005	0.000003
<i>Listeria</i> -hEphA2-ICD-1	694 (± 232)	0.0054	0.006
<i>Listeria</i> -hEphA2-ICD-2	731 (± 176)	0.052	0.004

**TABLE 10**

[00419] Figure 14B demonstrates the survival time of immunized animals. This data is summarized in Table 11 below:

Vaccination Group	Median Survival (Days)	P vs. HBSS	# Survivors (Day 65)
HBSS	19	-	0
<i>Listeria</i> Control	20	Ns	0
<i>Listeria</i> -hEphA2-ICD-1	>65	0.0035	3
<i>Listeria</i> -hEphA2-ICD-2	>65	0.0035	4
<i>Listeria</i> -hEphA2-ICD-3	>65	0.0035	4

**TABLE 11**

[00420] Immunization of Balb/C mice bearing CT26.24 (huEphA2+) lung tumors with recombinant *Listeria* encoding OVA.AH1 (MMTV gp70 immunodominant epitope) or

OVA.AH1-A5 (MMTV gp70 immunodominant epitope, with heteroclitic change for enhanced T-cell receptor binding) confers long-term survival (Figure 14C).

**[00421]** The EphA2 CO domain is strongly immunogenic, and a significant long term increase in survival of Balb/C mice bearing CT26.24 (huEphA2+) lung tumors was observed when immunized with recombinant *Listeria* encoding codon-optimized or native EphA2 CO domain sequence (Figure 14D).

**[00422]** The EphA2 EX2 domain is poorly immunogenic, and increased survival of Balb/C mice bearing CT26.24 (huEphA2+) lung tumors was observed only when immunized with recombinant *Listeria* encoding codon-optimized secA1 signal peptide fused with the codon-optimized EphA2 EX2 domain sequence. Therapeutic efficacy was not observed in mice when immunized with recombinant *Listeria* encoding native secA1 signal peptide fused with the codon-optimized EphA2 EX2 domain sequence (Figure 14E). The desirability of using both codon-optimized secA1 signal peptide and EphA2 EX2 domain sequences was supported by statistically significant therapeutic anti-tumor efficacy, as shown in the table below:

**[00423]** A comparison by log-rank test of survival curves shown in Figure 14E and summarized in Table 12 below:

Experimental Group	Median Survival (Days)	Significance versus HBSS cohort (p value)	Significance versus actA-native secA1/EphA2 EX2 cohort (p value)
HBSS	19	-	-
ActA	20	NS	NS
actA-native secA1-EphA2 EX2 (native)	19	NS	-
actA-native secA1-EphA2 EX2 (CodOp)	24	0.0035	NS
actA-CodOp secA1-EphA2 EX2 (CodOp)	37	0.0035	0.0162
actA-native secA1-EphA2 CO (CodOp)	>99	0.0035	0.0015

**TABLE 12**

**[00424]** Significantly, even though pCDNA4-EphA2 plasmid transfected 293 cells yielded very high levels of protein expression, immunization of Balb/C mice bearing CT26.24 (huEphA2+) lung tumors with the pCDNA4-EphA2 plasmid did not result in any observance of therapeutic anti-tumor efficacy (Figure 14F).

**[00425]** For therapeutic *in vivo* tumor studies, female Balb/C mice were implanted IV with  $5 \times 10^5$  CT26 cells stably expressing EphA2. Three days later, mice were randomized and vaccinated IV with various recombinant *Listeria* strains encoding EphA2. In some cases (noted in figures) mice were vaccinated with 100  $\mu$ g of pCDNA4 plasmid or

pCDNA4-EphA2 plasmid in the *tibialis* anterior muscle. As a positive control, mice were vaccinated IV with recombinant *Listeria* strains encoding OVA.AHI or OVA.AHI-A5 protein chimeras. Mice were vaccinated on days 3 and 14 following tumor cell implantation. Mice injected with Hanks Balanced Salt Solution (HBSS) buffer or unmodified *Listeria* served as negative controls. All experimental cohorts contained 5 mice. For survival studies mice were sacrificed when they started to show any signs of stress or labored breathing.

[00426] The foregoing data demonstrate that therapeutic immunization with *Listeria* expressing the hEphA2 suppresses established CT26-hEphA2 tumor growth and increases survival.

#### **6.6 EXAMPLE 6: Long-term Suppression of CT26-hEphA2 Tumor Growth Upon Rechallenge**

[00427] Balb/c mice failing to form tumors after preventative immunization with *Listeria* expressing either the ICD or ECD of hEphA2 against CT26-hEphA2 tumor challenged, were re-challenged (s.c.) with both CT26 parental cell line and CT26-hEphA2 cells on opposite flanks 56 days after initial tumor challenge and 60 days after the last immunization. Age-matched mice were used as a control in this experiment.

[00428] Re-challenge with parental CT26 cells showed no statistically significant differences in tumor growth between groups (data not shown). However, as shown in Figure 15, both groups vaccinated with *Listeria* expressing either the ICD or ECD of hEphA2 demonstrated a significant suppression of tumor growth upon re-challenge (\*p<0.041).

#### **6.7 EXAMPLE 7: Immunization with *Listeria* Expressing hEphA2 Elicits an EphA2-Specific CD8+ T Cell Response.**

[00429] Balb/c mice (n=3) were immunized with *Listeria* L461T expressing the intracellular domain of hEphA2 (hEphA2-ICD) or *ActA* expressing codon optimized extracellular domain of hEphA2 (hEphA2-ECD) two weeks apart. Mice were euthanized, and spleens harvested and pooled 6 days after the last immunization. For the ELISPOT assay, the cells were re-stimulated *in vitro* with P815 cells expressing full-length hEphA2 or cell lysates prepared from these cells. The parental P815 cells or cell lysates served as a negative control. Cells were also stimulated with recombinant hEphA2 Fc fusion protein. IFN-gamma positive spot forming colonies (SFCs) were measured using a 96 well spot reader.

[00430] As shown in Figure 16, increased IFN-gamma SFCs were observed with spleen cells derived from mice vaccinated with *Listeria*-hEphA2. Both hEphA2 expressing



cells or cell lysates stimulation resulted in an increase in IFN-gamma SFC which suggests an EphA2-specific CD8+ as well as CD4+ T cell response. Spleen cells from mice vaccinated with the parental *Listeria* control did not demonstrate an increase in IFN-gamma SFC.

5 **6.8 EXAMPLE 8: Both CD4+ and CD8+ T Cell Responses are Required for Maximal hEphA2-Directed Anti-Tumor Efficacy.**

[00431] Balb/c mice (n=10) were inoculated i.v. with  $2 \times 10^5$  CT26-hEphA2 on day 0. CD4+ cells and CD8+ T-cells were depleted by injecting 200  $\mu$ g anti-CD4 (ATCC hybridoma GK1.5) or anti-CD8 (ATCC hybridoma 2.4-3) on Days 1 and 3, which was confirmed by FACS analysis (data not shown). Mice were then immunized i.v. with 0.1 LD<sub>50</sub> *Listeria* L461T expressing hEphA2 ICD on Day 4 and monitored for survival.

[00432] As shown in Figure 17, both CD4+ and CD8+ depleted groups failed to demonstrate the degree of anti-tumor response seen in the non-T cell depleted animals. The data are summarized in Table 13 below:

Vaccination Group	Median Survival (Days)	P vs. HBSS	# Survivors (Day 67)
HBSS	17	-	0
<i>Listeria</i> -hEphA2-ICD	>67	<0.0001	7
<i>Listeria</i> -hEphA2-ICD + anti-CD4	19	0.03	2
<i>Listeria</i> -hEphA2-ICD + anti-CD8	24	0.0002	0

15 **TABLE 13**

[00433] The foregoing data indicate a requirement for both CD4+ and CD8+ T cells in optimal suppression of tumor growth.

20 **6.9 EXAMPLE 9: Therapeutic Vaccination with *Listeria* Expressing Human EphA2 ICD Enhances CD45+ Tumor Infiltrate**

[00434] Balb/c mice (n=3) were immunized with 0.1 LD<sub>50</sub> *actA*-*Listeria* control or *Listeria* expressing the either the ECD or ICD of hEphA2, 6 days post s.c. CT26-hEphA2 tumor inoculation. 9 days post-vaccination, tumors were harvested, fixed in 10% neutral buffered formalin, embedded in paraffin and sectioned at 4 $\mu$ m. Microscope slides were prepared from the tumor sections. The tissues on the slides were deparaffinized and rehydrated as follows: 4 changes with xylene, 5 minutes each; 2 changes with absolute alcohol, 5 minutes each; 1 change with 95% alcohol for 5 minutes; 1 change with 70% for 5 minutes; and two changes with distilled water.

- [00435]** Steam antigen retrieval was performed in a Black and Decker Rice steamer using target antigen retrieval (TAR) solution (DakoCytomation, Carpinteria, CA) using a modification of the manufacture's protocol. The slides were placed into TAR solution preheated to just below boiling temperature and incubated for 20 minutes. The slides were then removed from the TAR solution and allowed to cool at room temperature for 20 minutes, and rinsed twice in TBS assay buffer.
- [00436]** Staining of the slides with biotinylated antibody was performed as follows.
- [00437]** Endogenous peroxidase was blocked by immersing the slides in solution of 3% Hydrogen Peroxide in Methanol, for 10 minutes, followed with 2 changes of distilled water, 5 minutes each. Protein was blocked by immersing the slides in a solution of 5% Bovine Serum Albumin (BSA) in 1x Tris buffered saline with 0.01% Tween 20 (TBST) for at least 30 minutes.
- [00438]** After wiping excess BSA solution from the slide, creating a "pool", centered around tissue, the slide is laid flat in humid chamber and biotinylated rat anti-mouse CD45/B220 (Pharmingen) at 1:100 dilution in a solution of 1%BSA/TBST is applied. The slide is incubated a in humid chamber overnight at room temperature with care taken to prevent drying of the tissue sections.
- [00439]** The next morning, the slides are washed with 2 changes of TBST, the second one lasting 10 minutes. Streptavidin conjugated with either HRP or AP is applied, incubating for 30 minutes at room temp. The slides are washed with two changes of TBST, visualized with an appropriate substrate chromagen (for Strep-HRP, DAB is used). After a wash in distilled water, the slides are counterstained with Mayers Hematoxylin by immersing the slides in dye for 2 minutes. The slides are then washed in running tap water until water runs clear, immersed in bluing agent (Scotts substitute tap water) for 30 seconds, and washed again in tap water. The slides are dehydrated and cleared in graded alcohols through xylene (or xylene substitute) by the following washes: 95% alcohol for 1 minute, 3 changes absolute alcohol for 1 minute each, and 4 changes xylenes for 1 minute each.
- [00440]** Mounting media is applied to the cover slips (for xylene, DPX mountant is used) and the slides are allowed to dry over night prior to visualization.
- [00441]** The sections were visualized on a Nikon Eclipse E400 and images captured with a Nikon DXM1200 digital camera (Figure 18A). Data was further normalized to tumor volume (Figure 18B).
- [00442]** The results demonstrate that tumor associated infiltrating lymphocytes are increased following therapeutic vaccination.

## **7. EQUIVALENTS**

[00443] Those skilled in the art will recognize, or be able to ascertain using no more than routine experimentation, many equivalents to the specific embodiments of the invention described herein. Such equivalents are intended to be encompassed by the following claims.

[00444] All publications, patents and patent applications mentioned in this specification are herein incorporated by reference into the specification to the same extent as if each individual publication, patent or patent application was specifically and individually indicated to be incorporated herein by reference.

We claim:

1. A method of eliciting an immune response against an EphA2-expressing cell, said method comprising administering to an individual a composition comprising an EphA2 antigenic polypeptide in an amount effective to elicit an immune response against an  
5 EphA2-expressing cell.
2. The method of claim 1, wherein the composition further comprises an adjuvant.
3. The method of claim 1, wherein the composition comprises a heat shock protein bound to said EphA2 antigenic polypeptide.
4. The method of claim 1, where the polypeptide further comprises a protein transduction  
10 domain.
5. The method of claim 4, wherein the protein transduction domain is the Antennapedia or the HIV tat protein transduction domain.
6. A method of eliciting an immune response against an EphA2-expressing cell, said method comprising administering to an individual a composition comprising an EphA2  
15 antigenic polypeptide expression vehicle in an amount effective to elicit an immune response against an EphA2-expressing cell.
7. The method of claim 6, wherein the expression vehicle is a nucleic acid encoding said EphA2 antigenic polypeptide operably linked to a promoter.
8. The method of claim 7, wherein the nucleic acid is DNA.
- 20 9. The method of claim 8, wherein the DNA is conjugated to a carrier.
10. The method of claim 7, wherein the carrier is asialoglycoprotein.
11. The method of claim 7, wherein the carrier is transferrin.
12. The method of claim 7, wherein the carrier is polymeric IgA.
13. The method of claim 6, wherein the expression vehicle is an infectious agent  
25 comprising a nucleic acid, said nucleic acid comprising a nucleotide sequence encoding said EphA2 antigenic polypeptide operably linked to a promoter.

32. The method of claim 30, wherein the mammalian cell is encapsulated within a membrane.
33. The method of claim 32, wherein said administering is by means of implantation.
34. A method of eliciting an immune response against an EphA2-expressing cell, said  
5 method comprising administering to an individual a composition comprising antigen presenting cells sensitized with an EphA2 antigenic polypeptide.
35. The method of claim 34, further comprising prior to said administration the step of sensitizing the antigen presenting cells.
36. The method of claim 35, wherein the antigen presenting cells are sensitized by a  
10 method comprising: contacting the cells with a composition comprising one or more EphA2 antigenic peptides in an amount effective to sensitize the cells.
37. The method of claim 37, wherein the composition further comprises a heat shock protein.
38. The method of claim 37, wherein the heat shock protein is hsp70, gp96, or hsp90.
- 15 39. The method of claim 34, wherein the antigen presenting cells are autologous to the individual.
40. The method of claim 34, wherein the antigen presenting cells are non-autologous to the individual.
41. The method of claim 34, wherein the antigen presenting cells are macrophages.
- 20 42. The method of claim 34, wherein the antigen presenting cells are dendritic cells.
43. The method of claim 1, 6, or 34, wherein the individual has cancer.
44. The method of claim 43, wherein said cancer is of an epithelial cell origin.
45. The method of claim 43, wherein said cancer comprises cells that overexpress EphA2 relative to non-cancer cells having the tissue type of said cancer cells.
- 25 46. The method of claim 43, wherein said cancer is a cancer of the skin, lung, colon, ovary, esophagus, breast, prostate, bladder or pancreas or is a renal cell carcinoma or melanoma.

14. The method of claim 13, wherein the sequence encoding said EphA2 antigenic polypeptide is codon-optimized for expression in said infectious agent.
15. The method of claim 13, wherein the infectious agent is coated with a reagent that targets the infectious agent to EphA2-expressing cells.
- 5 16. The method of claim 15, wherein the reagent is an anti-EphA2 antibody.
17. The method of claim 13, wherein the infectious agent is coated with a reagent that targets the infectious agent to antigen-presenting cells.
18. The method of claim 13, wherein the infectious agent is a bacterium.
19. The method of claim 18, wherein the bacterium is attenuated.
- 10 20. The method of claim 18, wherein the nucleic acid comprises a nucleotide sequence encoding a secretory signal operatively linked to the sequence encoding the EphA2 antigenic polypeptide.
21. The method of claim 20, wherein the secretory signal is a SecA secretory signal.
22. The method of claim 18, wherein the bacterium is *Pseudomonas aeruginosa*.
- 15 23. The method of claim 13, wherein the infectious agent is a virus.
24. The method of claim 23, wherein the virus is a retrovirus.
25. The method of claim 24, wherein the retrovirus is a lentivirus.
26. The method of claim 23, wherein the virus is an adenovirus.
27. The method of claim 23, wherein the virus is an adeno-associated virus.
- 20 28. The method of claim 23, wherein the virus is herpes simplex virus.
29. The method of claim 23, wherein the virus is attenuated.
30. The method of claim 6, wherein the expression vehicle is a mammalian cell comprising a recombinant nucleic acid, said nucleic acid comprising a nucleotide sequence encoding said EphA2 antigenic polypeptide.
- 25 31. The method of claim 31, wherein the mammalian cell is a human cell.

47. The method of claim 1, 6, 34, or 50, wherein the individual has a non-neoplastic hyperproliferative disorder.
48. The method of claim 47, wherein the hyperproliferative disorder is an epithelial cell disorder.
- 5 49. The method of claim 48, wherein the hyperproliferative is asthma, chronic pulmonary obstructive disease, lung fibrosis, bronchial hyper responsiveness, psoriasis, and seborrheic dermatitis.
- 10 50. A method of eliciting an immune response against an EphA2-expressing cell, said method comprising administering to an individual a composition comprising an anti-idiotypic antibody or antigen-binding fragment thereof which immunospecifically binds to an idiotype of an anti-EphA2 antibody in an amount effective to elicit an immune response against an EphA2-expressing cell.
- 15 51. A method of treating a human individual having a hyperproliferative disorder of EphA2-expressing cells, said method comprising administering to the individual a composition comprising an EphA2 antigenic polypeptide in an amount effective to treat a hyperproliferative disorder of EphA2-expressing cells.
- 20 52. A method of treating a human individual having a hyperproliferative disorder of EphA2-expressing cells, said method comprising administering to the individual a composition comprising an EphA2 expression vehicle in an amount effective to treat a hyperproliferative disorder of EphA2-expressing cells.
- 25 53. A method of treating a human individual having a hyperproliferative disorder of EphA2-expressing cells, said method comprising administering to the individual a composition comprising antigen presenting cells sensitized with an EphA2 antigenic polypeptide in an amount effective to treat a hyperproliferative disorder of EphA2-expressing cells.
- 30 54. A method of treating a human individual having a hyperproliferative disorder of EphA2-expressing cells, said method comprising administering to an individual a composition comprising an anti-idiotypic antibody or antigen-binding fragment thereof which immunospecifically binds to an idiotype of an anti-EphA2 antibody in an amount effective to elicit treat a hyperproliferative disorder of EphA2-expressing cells.

55. The method of claim 51, 52, 53, or 54, wherein the individual has cancer.
56. The method of claim 55, wherein said cancer is of an epithelial cell origin.
57. The method of claim 55, wherein said cancer comprises cells that overexpress EphA2 relative to non-cancer cells having the tissue type of said cancer cells.
58. The method of claim 55, wherein said cancer is a cancer of the skin, lung, colon, breast, prostate, bladder or pancreas or is a renal cell carcinoma or melanoma.
59. The method of claim 51, 52, 53, or 54, wherein the individual has a non-neoplastic hyperproliferative disorder.
60. The method of claim 59, wherein the hyperproliferative disorder is an epithelial cell disorder.
61. The method of claim 60, wherein the hyperproliferative disorder is asthma, chronic pulmonary obstructive disease, lung fibrosis, bronchial hyper responsiveness, psoriasis, and seborrheic dermatitis.
62. The method of any one of claims 1, 6, 34, 51, 52, and 53, wherein the EphA2 polypeptide comprises full length EphA2.
63. The method of any one of claims 1, 6, 34, 51, 52, and 53, wherein the EphA2 polypeptide comprises the extracellular domain of EphA2.
64. The method of any one of claims 1, 6, 34, 51, 52, and 53, wherein the EphA2 polypeptide comprises the intracellular domain of EphA2.
65. The method of any one of claims 1, 6, 34, 51, 52, and 53, wherein the EphA2 polypeptide is a chimeric polypeptide comprising at least an antigenic portion of EphA2 and a second polypeptide.
66. The method of claim 50 or 54, wherein the EphA2 antibody immunospecifically binds to an epitope in the extracellular domain of EphA2.
67. The method of claim 50 or 54, wherein the EphA2 antibody immunospecifically binds to an epitope the intracellular domain of EphA2.



68. The method of any one of claims 1, 6, 34, 51, 52, and 53, wherein the EphA2 polypeptide is a chimeric polypeptide comprising at least an antigenic portion of EphA2 and a second polypeptide.
- 5 69. The method of claim 1, or 51, wherein the composition comprises a plurality of EphA2 antigenic polypeptides.
70. The method of claim 6 or 52, wherein the composition comprises a plurality of EphA2 antigenic polypeptide expression vehicles.
71. The method of claim 6 or 52, wherein the expression vehicle expresses a plurality of EphA2 antigenic polypeptides.
- 10 72. The method of claim 34 or 53, wherein the antigen presenting cells are sensitized with a plurality of EphA2 antigenic polypeptides
73. The method of any one of claims 1, 6, 34, 50, 51, 52, 53, and 54, further comprising administering an additional anti-cancer therapy.
74. The method of claim 73, wherein the additional anti-cancer therapy is an agonistic  
15 EphA2 antibody.
75. The method of claim 73, wherein the additional anti-cancer therapy is chemotherapy, biological therapy, immunotherapy, radiation therapy, hormonal therapy, or surgery.
76. The method of any one of claims 1, 6, 34, 50, 51, 52, 53, and 54, wherein said administering is mucosal, intranasal, parenteral, intramuscular, intravenous, oral or  
20 intraperitoneal.
77. The method of any one of claims 1, 6, 34, 50, 51, 52, 53, and 54, wherein the administration elicits a CD4<sup>+</sup> T-cell response, a CD8<sup>+</sup> T-cell response, an innate immune response, an antibody response, or a combination of one or more of the foregoing.
78. The method of claim m 80, wherein the administration elicits both a CD4<sup>+</sup> T-cell  
25 response and a CD8<sup>+</sup> T-cell response.
79. The method of claim 1, 6, 37 or 53, wherein the subject has a disease involving aberrant angiogenesis.

80. The method of claim 82, wherein the wherein the disease is macular degeneration, diabetic retinopathy, retinopathy of prematurity, vascular restenosis, infantile hemangioma, verruca vulgaris, psoriasis, Kaposi's sarcoma, neurofibromatosis, recessive dystrophic epidermolysis bullosa, rheumatoid arthritis, ankylosing spondylitis, systemic lupus, psoriatic arthropathy, Reiter's syndrome, and Sjogren's syndrome, endometriosis, preeclampsia, atherosclerosis or coronary artery disease.

81. A method of treating a human individual having a disease involving aberrant angiogenesis, said method comprising administering to the individual a composition comprising an EphA2 antigenic polypeptide in an amount effective to treat a disease involving aberrant angiogenesis.

82. A method of treating a human individual having a a disease involving aberrant angiogenesis, said method comprising administering to the individual a composition comprising an EphA2 expression vehicle in an amount effective to treat a disease involving aberrant angiogenesis.

83. A method of treating a human individual having a disease involving aberrant angiogenesis, said method comprising administering to the individual a composition comprising antigen presenting cells sensitized with an EphA2 antigenic polypeptide in an amount effective to treat a disease involving aberrant angiogenesis.

84. A method of treating a human individual having a a disease involving aberrant angiogenesis, said method comprising administering to an individual a composition comprising an anti-idiotypic antibody or antigen-binding fragment thereof which immunospecifically binds to an idio type of an anti-EphA2 antibody in an amount effective to elicit treat a disease involving aberrant angiogenesis.

85. The method of claim 84, 85 86, or 87 wherein the wherein the disease is macular degeneration, diabetic retinopathy, retinopathy of prematurity, vascular restenosis, infantile hemangioma, verruca vulgaris, psoriasis, Kaposi's sarcoma, neurofibromatosis, recessive dystrophic epidermolysis bullosa, rheumatoid arthritis, ankylosing spondylitis, systemic lupus, psoriatic arthropathy, Reiter's syndrome, and Sjogren's syndrome, endometriosis, preeclampsia, atherosclerosis or coronary artery disease.

## ABSTRACT

[00445] The present invention relates to methods and compositions designed for the treatment, management, or prevention of cancer, particularly metastatic cancer, and hyperproliferative diseases involving EphA2-expressing cells. In one embodiment, the

5 methods of the invention comprise the administration of an effective amount of an EphA2 antigenic peptide to elicit an immune response against the EphA2-expressing cells. In other embodiments, the methods of the invention entail the use of an EphA2 expression vehicle, such as a naked nucleic acid or viral vector. In yet other embodiments, the methods of the invention comprise the use of adoptive immunotherapy with autologous or non-autologous  
10 antigen presenting cells that are sensitized with one or more EphA2 antigenic peptides. The invention also provides pharmaceutical compositions comprising one or more EphA2 antigenic peptides, expression vehicles or antigen-presenting cells of the invention either alone or in combination with one or more other agents useful for cancer therapy.





FIGURE 1




FIGURE 2




FIGURE 3





FIGURE 4

# Human EphA2 Expression in CT26 Subclones


## *Generation of "the Super Clone"*

**FIGURE 5**

- Subclones of transfected clones obtained by FACS sorting into 96 well plates



# Human EphA2 Protein Expression in CT26 Murine Colon Carcinoma Cells Following FACS Sorting



•pooled populations of transfected cells  
sorted by FACS

FIGURE 6

# Human EphA2 Protein Expression in B16F10 Cells

## Round 3 – Success!




FIGURE 7




FIGURE 8

FIGURE 9





FIGURE 10A




FIGURE 10B


\*  $p=0.0017$

FIGURE 11A

# Effect of *L. monocytogenes* Expressing ECD of huEphA2 on CT26 huEphA2 Tumor Growth *Prevention Study*




FIGURE 11B

**Effect of *L. monocytogenes* Expressing ECD of huEphA2 on  
CT26 huEphA2 Tumor Growth  
Prevention Study**

**Listeria Control      Listeria-ECD-huEphA2**






FIGURE 11C

# Animal Survival in CT26 huEphA2 S.C. Model

## Prevention Study




FIGURE 11D

# **Animal Survival in CT26 huEphA2 Lung Metastases Model *Prevention Study***




Figure 12




Figure 13A

**Effect of *L. monocytogenes* Expressing  
ECD of huEphA2 on CT26 Tumor Growth**

**Therapeutic Study**




Figure 13B

# Effect of *L. monocytogenes* Expressing ECD of huEphA2 on CT26 Tumor Growth *Therapeutic Study*




Figure 13C

# Animal Survival in CT26 Experimental Lung Metastases Model





Figure 14A



\* p < 0.0061








Figure 14C

Figure 14D





Figure 14E






Figure 14F

Figure 15




Figure 16




Figure 17




Figure 18 A



*Listeria* Control*Listeria*-hEphA2 ECD*Listeria*-hEphA2 ICD



Figure 18B



**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

**BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☒ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☒ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**

# SEQUENCE LISTING

```

<110>  Kinch, Michael S.
<120>  EphA2 VACCINES
<130>  10271-143-888
<160>  72
<170>  PatentIn version 3.2

<210>  1
<211>  3963
<212>  DNA
<213>  Homo sapiens

<220>
<221>  CDS
<222>  (138)..(3068)

<400>  1
attaaggact cggggcagga ggggcagaag ttgcgcgcag gccggcgggc gggagcggac      60
accgaggccg gcgtgcagcg gtgcgggtgt gcgggagccg ggctcggggg gatcgggaccg      120
agagcgagaa gcgcggc atg gag ctc cag gca gcc cgc gcc tgc ttc gcc      170
              Met Glu Leu Gln Ala Ala Arg Ala Cys Phe Ala
              1              5              10

ctg ctg tgg ggc tgt gcg ctg gcc gcg gcc gcg gcg gcg cag gcc aag      218
Leu Leu Trp Gly Cys Ala Leu Ala Ala Ala Ala Gln Gly Lys
              15              20              25

gaa gtg gta ctg ctg gac ttt gct gca gct gga ggg gag ctc gcc tgg      266
Glu Val Val Leu Leu Asp Phe Ala Ala Ala Gly Gly Glu Leu Gly Trp
              30              35              40

ctc aca cac ccg tat ggc aaa ggg tgg gac ctg atg cag aac atc atg      314
Leu Thr His Pro Tyr Gly Lys Gly Trp Asp Leu Met Gln Asn Ile Met
              45              50              55

aat gac atg ccg atc tac atg tac tcc gtg tgc aac gtg atg tct ggc      362
Asn Asp Met Pro Ile Tyr Met Tyr Ser Val Cys Asn Val Met Ser Gly
              60              65              70              75

gac cag gac aac tgg ctc cgc acc aac tgg gtg tac cga gga gag gct      410
Asp Gln Asp Asn Trp Leu Arg Thr Asn Trp Val Tyr Arg Gly Glu Ala
              80              85              90

gag cgt atc ttc att gag ctc aag ttt act gta cgt gac tgc aac agc      458
Glu Arg Ile Phe Ile Glu Leu Lys Phe Thr Val Arg Asp Cys Asn Ser
              95              100              105

ttc cct ggt ggc gcc agc tcc tgc aag gag act ttc aac ctc tac tat      506
Phe Pro Gly Gly Ala Ser Ser Cys Lys Glu Thr Phe Asn Leu Tyr Tyr
              110              115              120

gcc gag tgc gac ctg gac tac gcc acc aac ttc cag aag cgc ctg ttc      554
Ala Glu Ser Asp Leu Asp Tyr Gly Thr Asn Phe Gln Lys Arg Leu Phe

```

125	130	135	
acc aag att gac Thr Lys Ile Asp 140	acc att gcg ccc gat Thr Ile Ala Pro Asp 145	gag atc acc gtc agc agc gac Ile Thr Val Ser Ser Asp 150	602
ttc gag gca cgc Phe Glu Ala Arg 160	cac gtg aag ctg aac His Val Lys Leu Asn 165	gag gag cgc tcc gtg ggg Val Glu Glu Arg Ser Val Gly 170	650
ccg ctc acc cgc Pro Leu Thr Arg 175	aaa ggc ttc tac Lys Gly Phe Tyr 180	ctg gcc ttc cag gat atc ggt gcc Ala Phe Gln Asp Ile Gly Ala 185	698
tgt gtg gcg ctg Cys Val Ala Leu 190	ctc tcc gtc cgt Leu Ser Val Arg 195	tac tac aag aag tgc ccc gag Tyr Tyr Lys Lys Cys Pro Glu 200	746
ctg ctg cag ggc Leu Leu Gln Gly 205	ctg gcc cac ttc cct Leu His Phe Pro 210	gag acc atc gcc ggc tct gat Glu Thr Ile Ala Gly Ser Asp 215	794
gca cct tcc ctg Ala Pro Ser Leu 220	gcc act gtg gcc ggc Val Ala Gly Thr 225	acc tgt gtg gac cat gcc gtg Cys Val Asp His Ala Val 230	842
gtg cca ccg ggg Val Pro Pro Gly 240	ggg ggt gaa gag ccc Glu Glu Pro Arg 245	cgt atg cac tgt gca gtg gat ggc Met His Cys Ala Val Asp Gly 250	890
gag tgg ctg gtg Glu Trp Leu Val 255	ccc att ggg cag tgc Pro Ile Gly Gln Cys 260	ctg tgc cag gca ggc tac gag Leu Cys Gln Ala Gly Tyr Glu 265	938
aag gtg gag gat Lys Val Glu Asp 270	gcc tgc cag gcc tgc Ala Cys Gln Ala Cys 275	cct gga ttt ttt aag ttt Pro Gly Phe Phe Lys Phe 280	986
gag gca tct gag Glu Ala Ser Glu 285	agc ccc tgc ttg gag tgc Ser Pro Cys Leu Glu Cys 290	cct gag cac acg ctg cca Pro Glu His Thr Leu Pro 295	1034
tcc cct gag ggt Ser Pro Glu Gly 300	gcc acc tcc tgc gag tgt Ala Thr Ser Cys Glu Cys 305	gag gaa ggc ttc ttc cgg Glu Glu Gly Phe Phe Arg 310	1082
gca cct cag gac Ala Pro Gln Asp 320	cca gcg tgc atg cct Pro Ala Ser Met Pro 325	tgc aca cga ccc ccc tcc gcc Thr Arg Pro Pro Ser Ala 330	1130
cca cac tac ctc Pro His Tyr Leu 335	aca gcc gtg ggc atg ggt Leu Thr Ala Val Gly Met 340	gcc aag gtg gag ctg cgc Ala Lys Val Glu Leu Arg 345	1178
tgg acg ccc cct Trp Thr Pro Pro 350	cag gac agc ggg ggc Gln Asp Ser Gly Gly 355	gag gac att gtc tac agc Glu Asp Ile Val Tyr Ser 360	1226
gtc acc tgc gaa Val Thr Cys Glu 365	cag tgc tgg ccc gag Gln Cys Trp Pro 370	tgc ggg gaa tgc ggg ccg tgt Glu Cys Gly Pro Cys 375	1274

gag gcc agt gtg cgc tac tgc gag cct cct cac gga ctg acc cgc acc Glu Ala Ser Val Arg Tyr Ser Glu Pro His Gly Leu Thr Arg Thr 380 385 390 395	1322
agt gtg aca gtg agc gac ctg gag ccc cac atg aac tac acc ttc acc Ser Val Thr Val Ser Asp Leu Glu Pro His Met Asn Tyr Thr Phe Thr 400 405 410	1370
gtg gag gcc cgc aat ggc gtc tca ggc ctg gta acc agc cgc agc ttc Val Glu Ala Arg Asn Gly Val Ser Gly Leu Val Thr Ser Arg Ser Phe 415 420 425	1418
cgt act gcc agt gtc agc atc aac cag aca gag ccc ccc aag gtg agg Arg Thr Ala Ser Val Ser Ile Asn Gln Thr Glu Pro Lys Val Arg 430 435 440	1466
ctg gag ggc cgc agc acc acc tgc ctt agc gtc tcc tgg agc atc ccc Leu Glu Gly Arg Ser Thr Thr Ser Leu Ser Val Ser Trp Ser Ile Pro 445 450 455	1514
ccg ccg cag cag agc cga gtg tgg aag tac gag gtc act tac cgc aag Pro Pro Gln Gln Ser Arg Val Trp Lys Tyr Glu Val Thr Tyr Arg Lys 460 465 470 475	1562
aag gga gac tcc aac agc tac aat gtg cgc cgc acc gag ggt ttc tcc Lys Gly Asp Ser Asn Ser Tyr Asn Val Arg Arg Thr Glu Gly Phe Ser 480 485 490	1610
gtg acc ctg gac gac ctg gcc cca gac acc acc tac ctg gtc cag gtg Val Thr Leu Asp Asp Leu Ala Pro Thr Thr Tyr Leu Val Gln Val 495 500 505	1658
cag gca ctg acg cag gag ggc cag ggg gcc gcc agc aag gtg cac gaa Gln Ala Leu Thr Gln Glu Gly Gln Gly Ala Gly Ser Lys Val His Glu 510 515 520	1706
ttc cag acg ctg tcc ccg gag gga tct ggc aac ttg gcg gtg att ggc Phe Gln Thr Leu Ser Pro Glu Gly Ser Gly Asn Ala Val Ile Gly 525 530 535	1754
ggc gtg gct gtc ggt gtg gtc ctg ctt ctg gtg ctg gca gga gtt ggc Gly Val Ala Val Gly Val Val Leu Leu Leu Val Leu Ala Gly Val Gly 540 545 550 555	1802
ttc ttt atc cac cgc agg agg aag aac cag cgt gcc cgc cag tcc ccg Phe Phe Ile His Arg Arg Arg Lys Asn Gln Arg Ala Arg Gln Ser Pro 560 565 570	1850
gag gac gtt tac ttc tcc aag tca gaa caa ctg aag ccc ctg aag aca Glu Asp Val Tyr Ser Ser Lys Ser Glu Gln Leu Lys Pro Leu Lys Thr 575 580 585	1898
tac gtg gac ccc cac aca tat gag gac ccc aac cag gct gtg ttg aag Tyr Val Asp Pro His Thr Tyr Glu Asp Pro Asn Gln Ala Leu Lys 590 595 600	1946
ttc act acc gag atc cat cca tcc tgt gtc act cgg cag aag gtg atc Phe Thr Thr Glu Ile His Pro Ser Cys Val Thr Arg Gln Lys Val Ile 605 610 615	1994

gga gca gga gag ttt ggg gag gtg tac aag ggc atg ctg aag aca tcc Gly Ala Gly Glu Phe Gly Glu Val Tyr Lys Gly Met Leu Lys Thr Ser 620 625 630 635	2042
tcg ggg aag aag gag gtg ccg gtg gcc atc aag acg ctg aaa gcc ggc Ser Gly Lys Lys Glu Val Pro Val Ala Ile Lys Thr Leu Lys Ala Gly 640 645 650	2090
tac aca gag aag cag cga gtg gac ttc ctc ggc gag gcc ggc atc atg Tyr Thr Glu Lys Glu Arg Val Asp Phe Leu Gly Glu Ala Gly Ile Met 655 660 665	2138
ggc cag ttc agc cac cac aac atc atc cgc cta gag ggc gtc atc tcc Gly Gln Phe Ser His His Asn Ile Ile Arg Leu Glu Gly Val Ile Ser 670 675 680	2186
aaa tac aag ccc atg atg atc atc act gag tac atg gag aat ggg gcc Lys Tyr Lys Pro Met Met Ile Ile Thr Glu Tyr Met Glu Asn Gly Ala 685 690 695	2234
ctg gac aag ttc ctt cgg gag aag gat ggc gag ttc agc gtg ctg cag Leu Asp Lys Phe Leu Arg Glu Lys Asp Gly Phe Ser Val Leu Gln 700 705 710 715	2282
ctg gtg ggc atg ctg cgg ggc atc gca gct ggc atg aag tac ctg gcc Leu Val Gly Met Leu Arg Gly Ile Ala Ala Gly Met Lys Tyr Leu Ala 720 725 730	2330
aac atg aac tat gtg cac cgt gac ctg gct gcc cgc aac atc ctc gtc Asn Met Asn Tyr Val His Arg Asp Leu Ala Ala Arg Asn Ile Leu Val 735 740 745	2378
aac agc aac ctg gtc tgc aag gtg tct gac ttt ggc ctg tcc cgc gtg Asn Ser Asn Leu Val Cys Lys Val Ser Asp Phe Gly Leu Ser Arg Val 750 755 760	2426
ctg gag gac gac ccc gag gcc acc tac acc acc agt ggc ggc aag atc Leu Glu Asp Asp Pro Glu Ala Thr Tyr Thr Thr Ser Gly Gly Lys Ile 765 770 775	2474
ccc atc cgc tgg acc gcc cgg gag gcc att tcc tac cgg aag ttc acc Pro Ile Arg Trp Thr Ala Pro Glu Ala Ile Ser Tyr Arg Lys Phe Thr 780 785 790 795	2522
tct gcc agc gac gtg tgg agc ttt ggc att gtc atg tgg gag gtg atg Ser Ala Ser Asp Val Trp Ser Phe Gly Ile Val Met Trp Glu Val Met 800 805 810	2570
acc tat ggc gag cgg ccc tac tgg gag ttg tcc aac cac gag gtg atg Thr Tyr Gly Glu Arg Pro Tyr Trp Glu Leu Ser Asn His Glu Val Met 815 820 825	2618
aaa gcc atc aat gat ggc ttc cgg ctc ccc aca ccc atg gac tgc ccc Lys Ala Ile Asn Asp Gly Phe Arg Leu Pro Thr Pro Met Asp Cys Pro 830 835 840	2666
tcc gcc atc tac cag ctc atg atg cag tgc tgg cag cag gag cgt gcc Ser Ala Ile Tyr Gln Leu Met Met Gln Cys Trp Gln Gln Glu Arg Ala 845 850 855	2714
cgc cgc ccc aag ttc gct gac atc gtc agc atc ctg gac aag ctc att	2762

Arg	Arg	Pro	Lys	Phe	Ala	Asp	Ile	Val	Ser	Ile	Leu	Asp	Lys	Leu	Ile				
860					865					870					875				
cg	gc	cc	ga	tc	tc	aa	ac	ct	gc	ga	tt	ga	cc	cg	gt	2810			
Arg	Ala	Pro	Asp	Ser	Leu	Lys	Thr	Leu	Ala	Asp	Phe	Asp	Pro	Arg	Val				
				880				885						890					
tc	at	cg	ct	cc	ag	ac	gc	gg	tc	ga	gg	gt	cc	tt	cg	2858			
Ser	Ile	Arg	Leu	Pro	Ser	Thr	Ser	Gly	Ser	Glu	Gly	Val	Pro	Phe	Arg				
			895					900					905						
ac	gt	tc	ga	tg	ct	ga	tc	at	aa	at	ca	ca	ta	ac	ga	2906			
Thr	Val	Ser	Glu	Trp	Leu	Glu	Ser	Ile	Lys	Met	Gln	Gln	Tyr	Thr	Glu				
			910					915					920						
ca	tt	at	gc	gc	gg	ta	ac	gc	at	ga	aa	gt	gt	ca	at	2954			
His	Phe	Met	Ala	Ala	Gly	Tyr	Thr	Ala	Ile	Glu	Lys	Val	Val	Gln	Met				
			925				930					935							
ac	aa	ga	ga	at	aa	ag	at	gg	gt	cg	ct	cc	gg	ca	ca	3002			
Thr	Asn	Asp	Asp	Ile	Lys	Arg	Ile	Gly	Val	Arg	Leu	Pro	Gly	His	Gln				
				945					950						955				
aa	cg	at	gc	ta	ag	ct	ct	ga	ct	aa	ga	ca	gt	aa	ac	3050			
Lys	Arg	Ile	Ala	Tyr	Ser	Leu	Leu	Gly	Leu	Lys	Asp	Gln	Val	Asn	Thr				
				960					965										
gt	gg	at	cc	at	ta	gc	tc	ga	ca	gc	gc	cc	gc	gc	gc	3098			
Val	Gly	Ile	Pro	Ile															
				975															
aa	ga	ata	ct	ga	aa	ac	ag	ag	tg	gc	tc	at	gc	tg	gc	act	ggg	gact	3158
tt	at	tt	at	tt	ct	gt	tt	ct	ct	cc	cc	ct	gc	ca	act	tc	gc	gc	3218
cc	tg	gc	ct	gc	ta	act	ga	gg	ga	ta	gc	ac	gg	ga	ta	gc	gc	gc	3278
cg	ca	ca	gc	gc	ta	gc	ca	ct	ta	gc	ca	ct	gc	ta	gc	gc	gc	gc	3338
cc	gc	ca	gc	gc	ta	gc	ca	ct	ta	gc	ca	ct	gc	ta	gc	gc	gc	gc	3398
tc	cc	ca	gc	gc	ta	gc	ca	ct	ta	gc	ca	ct	gc	ta	gc	gc	gc	gc	3458
cc	ta	gc	gc	gc	ta	gc	ca	ct	ta	gc	ca	ct	gc	ta	gc	gc	gc	gc	3518
aa	gc	gc	gc	gc	ta	gc	ca	ct	ta	gc	ca	ct	gc	ta	gc	gc	gc	gc	3578
ac	tg	gc	ca	aa	cc	ct	gc	ct	ta	gc	ca	ct	ta	gc	ca	aa	gc	gc	3638
aa	at	at	at	at	tt	gt	ac	tt	tt	gt	gt	gt	gt	gt	gt	gt	gt	gt	3698
gc	tg	gg	gg	ga	ga	gg	gt	gc	ta	gc	gg	gg	gc	gc	gc	gc	gc	gc	3758
ag	tg	tc	tc	gc	cc	at	gc	cc	gc	gc	gc	gc	gc	gc	gc	gc	gc	gc	3818
tg	tc	ag	tg	tt	aa	g	at	tt	tt	gt	tg	ga	ca	tt	tt	tt	tt	tt	3878
tt	tt	at	at	tt	at	tg	ta	ga	aa	gc	ta	gc	ta	gc	ta	gc	ta	gc	3938
ca	aa	cc	gc	aa	aaaa	aaaa	aaaa	aaaa	aaaa	aaaa	aaaa	aaaa	aaaa	aaaa	aaaa	aaaa	aaaa	aaaa	3963

<210> 2  
 <211> 976  
 <212> PRT  
 <213> Homo sapiens

<400> 2

```
Met Glu Leu Gln Ala Ala Arg Ala Cys Phe Ala Leu Leu Trp Gly Cys
1                               5                               10                               15

Ala Leu Ala Ala Ala Ala Ala Ala Gln Gly Lys Glu Val Val Leu Leu
20                               25                               30

Asp Phe Ala Ala Ala Gly Gly Glu Leu Gly Trp Leu Thr His Pro Tyr
35                               40                               45

Gly Lys Gly Trp Asp Leu Met Gln Asn Ile Met Asn Asp Met Pro Ile
50                               55                               60

Tyr Met Tyr Ser Val Cys Asn Val Met Ser Gly Asp Gln Asp Asn Trp
65                               70                               75                               80

Leu Arg Thr Asn Trp Val Tyr Arg Gly Glu Ala Glu Arg Ile Phe Ile
85                               90                               95

Glu Leu Lys Phe Thr Val Arg Asp Cys Asn Ser Phe Pro Gly Gly Ala
100                              105                              110

Ser Ser Cys Lys Glu Thr Phe Asn Leu Tyr Tyr Ala Glu Ser Asp Leu
115                              120                              125

Asp Tyr Gly Thr Asn Phe Gln Lys Arg Leu Phe Thr Lys Ile Asp Thr
130                              135                              140

Ile Ala Pro Asp Glu Ile Thr Val Ser Ser Asp Phe Glu Ala Arg His
145                              150                              155                              160

Val Lys Leu Asn Val Glu Glu Arg Ser Val Gly Pro Leu Thr Arg Lys
165                              170                              175

Gly Phe Tyr Leu Ala Phe Gln Asp Ile Gly Ala Cys Val Ala Leu Leu
180                              185                              190

Ser Val Arg Val Tyr Tyr Lys Lys Cys Pro Glu Leu Leu Gln Gly Leu
195                              200                              205

Ala His Phe Pro Glu Thr Ile Ala Gly Ser Asp Ala Pro Ser Leu Ala
```



210	215	220
Thr Val Ala Gly Thr Cys Val Asp His Ala Val Val Pro Pro Gly Gly		
225	230	235 240
Glu Glu Pro Arg Met His Cys Ala Val Asp Gly Glu Trp Leu Val Pro		
	245	250 255
Ile Gly Gln Cys Leu Cys Gln Ala Gly Tyr Glu Lys Val Glu Asp Ala		
	260	265 270
Cys Gln Ala Cys Ser Pro Gly Phe Phe Lys Phe Glu Ala Ser Glu Ser		
	275	280 285
Pro Cys Leu Glu Cys Pro Glu His Thr Leu Pro Ser Pro Glu Gly Ala		
	290	295 300
Thr Ser Cys Glu Cys Glu Glu Gly Phe Phe Arg Ala Pro Gln Asp Pro		
305	310	315 320
Ala Ser Met Pro Cys Thr Arg Pro Pro Ser Ala Pro His Tyr Leu Thr		
	325	330 335
Ala Val Gly Met Gly Ala Lys Val Glu Leu Arg Trp Thr Pro Pro Gln		
	340	345 350
Asp Ser Gly Gly Arg Glu Asp Ile Val Tyr Ser Val Thr Cys Glu Gln		
	355	360 365
Cys Trp Pro Glu Ser Gly Glu Cys Gly Pro Cys Glu Ala Ser Val Arg		
	370	375 380
Tyr Ser Glu Pro Pro His Gly Leu Thr Arg Thr Ser Val Thr Val Ser		
385	390	395 400
Asp Leu Glu Pro His Met Asn Tyr Thr Phe Thr Val Glu Ala Arg Asn		
	405	410 415
Gly Val Ser Gly Leu Val Thr Ser Arg Ser Phe Arg Thr Ala Ser Val		
	420	425 430
Ser Ile Asn Gln Thr Glu Pro Pro Lys Val Arg Leu Glu Gly Arg Ser		
	435	440 445
Thr Thr Ser Leu Ser Val Ser Trp Ser Ile Pro Pro Pro Gln Gln Ser		

450		455		460
Arg Val Trp Lys Tyr Glu Val Thr Tyr Arg Lys Lys Gly Asp Ser Asn				
465		470		480
Ser Tyr Asn Val Arg Arg Thr Glu Gly Phe Ser Val Thr Leu Asp Asp				
	485		490	495
Leu Ala Pro Asp Thr Thr Tyr Leu Val Gln Val Gln Ala Leu Thr Gln				
	500		505	510
Glu Gly Gln Gly Ala Gly Ser Lys Val His Glu Phe Gln Thr Leu Ser				
	515		520	525
Pro Glu Gly Ser Gly Asn Leu Ala Val Ile Gly Gly Val Ala Val Gly				
	530		535	540
Val Val Leu Leu Leu Val Leu Ala Gly Val Gly Phe Phe Ile His Arg				
	545		550	555
Arg Arg Lys Asn Gln Arg Ala Arg Gln Ser Pro Glu Asp Val Tyr Phe				
	565		570	575
Ser Lys Ser Glu Gln Leu Lys Pro Leu Lys Thr Tyr Val Asp Pro His				
	580		585	590
Thr Tyr Glu Asp Pro Asn Gln Ala Val Leu Lys Phe Thr Thr Glu Ile				
	595		600	605
His Pro Ser Cys Val Thr Arg Gln Lys Val Ile Gly Ala Gly Glu Phe				
	610		615	620
Gly Glu Val Tyr Lys Gly Met Leu Lys Thr Ser Ser Gly Lys Lys Glu				
	625		630	635
Val Pro Val Ala Ile Lys Thr Leu Lys Ala Gly Tyr Thr Glu Lys Gln				
	645		650	655
Arg Val Asp Phe Leu Gly Glu Ala Gly Ile Met Gly Gln Phe Ser His				
	660		665	670
His Asn Ile Ile Arg Leu Glu Gly Val Ile Ser Lys Tyr Lys Pro Met				
	675		680	685
Met Ile Ile Thr Glu Tyr Met Glu Asn Gly Ala Leu Asp Lys Phe Leu				
	690		695	700

Arg Glu Lys Asp Gly Glu Phe Ser Val Leu Gln Leu Val Gly Met Leu  
 705 710 715 720

Arg Gly Ile Ala Ala Gly Met Lys Tyr Leu Ala Asn Met Asn Tyr Val  
 725 730 735

His Arg Asp Leu Ala Ala Arg Asn Ile Leu Val Asn Ser Asn Leu Val  
 740 745 750

Cys Lys Val Ser Asp Phe Gly Leu Ser Arg Val Leu Glu Asp Asp Pro  
 755 760 765

Glu Ala Thr Tyr Thr Thr Ser Gly Gly Lys Ile Pro Ile Arg Trp Thr  
 770 775 780

Ala Pro Glu Ala Ile Ser Tyr Arg Lys Phe Thr Ser Ala Ser Asp Val  
 785 790 795 800

Trp Ser Phe Gly Ile Val Met Trp Glu Val Met Thr Tyr Gly Glu Arg  
 805 810 815

Pro Tyr Trp Glu Leu Ser Asn His Glu Val Met Lys Ala Ile Asn Asp  
 820 825 830

Gly Phe Arg Leu Pro Thr Pro Met Asp Cys Pro Ser Ala Ile Tyr Gln  
 835 840 845

Leu Met Met Gln Cys Trp Gln Gln Glu Arg Ala Arg Arg Pro Lys Phe  
 850 855 860

Ala Asp Ile Val Ser Ile Leu Asp Lys Leu Ile Arg Ala Pro Asp Ser  
 865 870 875 880

Leu Lys Thr Leu Ala Asp Phe Asp Pro Arg Val Ser Ile Arg Leu Pro  
 885 890 895

Ser Thr Ser Gly Ser Glu Gly Val Pro Phe Arg Thr Val Ser Glu Trp  
 900 905 910

Leu Glu Ser Ile Lys Met Gln Gln Tyr Thr Glu His Phe Met Ala Ala  
 915 920 925

Gly Tyr Thr Ala Ile Glu Lys Val Val Gln Met Thr Asn Asp Asp Ile  
 930 935 940

Lys Arg Ile Gly Val Arg Leu Pro Gly His Gln Lys Arg Ile Ala Tyr  
 945 950 955 960

Ser Leu Leu Gly Leu Lys Asp Gln Val Asn Thr Val Gly Ile Pro Ile  
 965 970 975

<210> 3  
 <211> 12  
 <212> PRT  
 <213> Homo sapiens  
 <400> 3

Thr Leu Ala Asp Phe Asp Pro Arg Val Pro Arg Thr  
 1 5 10

<210> 4  
 <211> 9  
 <212> PRT  
 <213> Homo sapiens  
 <400> 4

Val Leu Leu Leu Val Leu Ala Gly Val  
 1 5

<210> 5  
 <211> 9  
 <212> PRT  
 <213> Homo sapiens  
 <400> 5

Val Leu Ala Gly Val Gly Phe Phe Ile  
 1 5

<210> 6  
 <211> 9  
 <212> PRT  
 <213> Homo sapiens  
 <400> 6

Ile Met Asn Asp Met Pro Ile Tyr Met  
 1 5

<210> 7  
 <211> 9  
 <212> PRT  
 <213> Homo sapiens  
 <400> 7

Ser Leu Leu Gly Leu Lys Asp Gln Val  
1 5

<210> 8  
<211> 9  
<212> PRT  
<213> Homo sapiens

<400> 8

Trp Leu Val Pro Ile Gly Gln Cys Leu  
1 5

<210> 9  
<211> 9  
<212> PRT  
<213> Homo sapiens

<400> 9

Leu Leu Trp Gly Cys Ala Leu Ala Ala  
1 5

<210> 10  
<211> 9  
<212> PRT  
<213> Homo sapiens

<400> 10

Gly Leu Thr Arg Thr Ser Val Thr Val  
1 5

<210> 11  
<211> 9  
<212> PRT  
<213> Homo sapiens

<400> 11

Asn Leu Tyr Tyr Ala Glu Ser Asp Leu  
1 5

<210> 12  
<211> 9  
<212> PRT  
<213> Homo sapiens

<400> 12

Lys Leu Asn Val Glu Glu Arg Ser Val  
1 5

<210> 13

<211> 9  
<212> PRT  
<213> Homo sapiens

<400> 13

Ile Met Gly Gln Phe Ser His His Asn  
1 5

<210> 14  
<211> 9  
<212> PRT  
<213> Homo sapiens

<400> 14

Tyr Ser Val Cys Asn Val Met Ser Gly  
1 5

<210> 15  
<211> 9  
<212> PRT  
<213> Homo sapiens

<400> 15

Met Gln Asn Ile Met Asn Asp Met Pro  
1 5

<210> 16  
<211> 15  
<212> PRT  
<213> Homo sapiens

<400> 16

Glu Ala Gly Ile Met Gly Gln Phe Ser His His Asn Ile Ile Arg  
1 5 10 15

<210> 17  
<211> 13  
<212> PRT  
<213> Homo sapiens

<400> 17

Pro Ile Tyr Met Tyr Ser Val Cys Asn Val Met Ser Gly  
1 5 10

<210> 18  
<211> 16  
<212> PRT  
<213> Homo sapiens

<400> 18

Asp Leu Met Gln Asn Ile Met Asn Asp Met Pro Ile Tyr Met Tyr Ser  
 1 5 10 15

<210> 19  
 <211> 3105  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Fusion construct

<400> 19  
 atgaaaaaaa taatgctagt ttttattaca cttatattag ttagtctacc aattgcgcaa 60  
 caaactgaag caaaggatgc atctgcattc aataaagaaa attcaatttc atccatggca 120  
 ccaccagcat ctccgcctgc aagtcctaag acgccaatcg aaaagaaca cgcggatetc 180  
 gagctccagg cagcccgccg ctgcttcgcc ctgctgtggg gctgtgcgct ggccgcggcc 240  
 gcggcgccgc agggcaagga agtggtagct ctggactttg ctgcagctgg aggggagetc 300  
 ggctggctca cacaccgta tggcaagggt tgggacctga tgcagaacat catgaatgac 360  
 atgcgcgatct acatgtactc cgtgtgcaac gtgagtgtct gcgaccagga caactggctc 420  
 cgcaccaact ggggtgtacc aggagaggct gagcgtatct tcattgagct caagtttact 480  
 gtacgtgact gcaacagctt ccctggtggc gccagctcct gcaaggagac tttaacctc 540  
 tactatgccg agtcggacct ggactacggc accaacttcc agaagcgctt gttcaccaag 600  
 attgacacca ttgcgcccga tgagatcacc gtcagcagcg acttcgaggc acgccacgtg 660  
 aagctgaacg tggaggagcg ctccgtgggg ccgctcacc gcaagcgctt ctacctggcc 720  
 ttccaggata tcggtgcctg tgtggcgctg ctctccgtcc gtgtctacta caagaagtgc 780  
 cccgagctgc tgcaggacct ggcaccttc cctgagacca tcgccggctc tgatgcacct 840  
 tccctggcca ctgtggccgg cacctgtgtg gaccatgccg tgggtgccacc ggggggtgaa 900  
 gagccccgta tgcactgtgc agtggatggc gagggtgggtg tgccattggc gcagtgcctg 960  
 tgccaggcag gctacgagaa ggtggaggat gcctgccagg cctgctcgcc tggatttttt 1020  
 aagtttgagg catctgagag ccctcgcttg gaggtccttg agcacacgct gccatccctt 1080  
 gaggggtgcca cctcctgcga gtgtgaggaa ggcttcttcc gggcacctca ggacccagcg 1140  
 tcgatgcctt gcacacgacc ccctccgcc ccacactacc tcacagccgt gggcatgggt 1200  
 gccaaagttg agctgcgctg gacgccccct caggacagcg ggggcccgga ggacattgtc 1260  
 tacagcgtea cctgcgaaca gtgctggccc gagtctgggg aatgcggggc gtgtgaggcc 1320  
 agtgtgcgct actcggagcc tcttcacgga ctgacccgca ccagtgtgac agtgagcgac 1380  
 ctggagcccc acatgaacta caccctcacc gtggaggccc gcaatggcgt ctcaggcctg 1440

gtaaccagcc gcagcttccg tactgccagt gtcagcatca accagacaga gcccccaag 1500  
 gtgaggctgg agggcccgag caccacctcg cttagcgtct cctggagcat cccccgcgg 1560  
 cagcagagcc gagtgtgaa gtacgaggtc acttaccgca agaagggaga ctccaacagc 1620  
 tacaatgtgc gccgcaccga gggtttctcc gtgacctggg acgacctggc ccagacacc 1680  
 acctacctgg tccaggtgca ggcactgacg caggaggggc agggggccgg cagcagggtg 1740  
 cacgaattcc agacgctgtc ccggaggga tctggcaact tggcggatgat tggcggcgtg 1800  
 gctgtcgggtg tggctcgtct tctgggtgctg gcaggagttg gcttctttat ccaccgcagg 1860  
 aggaagaacc agcgtgcccc ccagtcctcg gaggacgttt acttctccaa gtcagaacaa 1920  
 ctgaagcccc tgaagacata cgtggacccc cacacatatg aggaccccaa ccaggctgtg 1980  
 ttgaagtcca ctaccgagat ccatccatcc tgtgtcactc ggcagaaggt gatcggagca 2040  
 ggagagtttg gggagggtga caaggccatg ctgaagacat cctcggggaa gaaggagggtg 2100  
 ccggtggcca tcaagacgtc gaaagccggc tacacagaga agcagcgagt ggacttctc 2160  
 ggcgaggccg gcacatcggg ccagttcagc caccacaaca tcatccgctc agagggcgtc 2220  
 atctccaaat acaagcccat gatgatcacc actgagtaca tggagaatgg ggcctggac 2280  
 aagttctctc gggagaagga tggcgagttc agcgtgctgc agctggtggg catgtgctg 2340  
 ggcatcgcag ctggcatgaa gtacctggcc aacatgaact atgtgcaccg tgacctggct 2400  
 gccgcgaaca tctcgtctaa cagcaacctg gtcgtcaagg tgtctgactt tggcctgtcc 2460  
 cgctgtctgg aggcagacc ccaggccacc tacaccacca gtggcggcaa gatcccatc 2520  
 cgctggaccg ccccgaggc catttcttac cggaagtcca cctctgccag cgacgtgtg 2580  
 agctttggca ttgtcatgtg ggagggtgat acctatggcg agcggcccta ctgggagttg 2640  
 tccaaccacg aggtgatgaa agccatcaat gatgcttcc ggctccccac acccatggac 2700  
 tgccctcccg ccactaccga gctcatgatg cagtgcctggc agcaggagcg tgcgcccg 2760  
 cccaagttcg ctgacatcgt cagcatcctg gacaagctca ttcgtgcccg tgactccctc 2820  
 aagaccctgg ctgactttga ccccgcggtg tctatccggc tccccagcac gagcggctcg 2880  
 gaggggggtgc ccttccgcac ggtgtccgag tggctggagt ccatcaagat gcagcagtat 2940  
 acggagcact tcatggcgcg cggtacact gccatcgaga aggtggtgca gatgaccaac 3000  
 gacgacatca agaggattgg ggtgcggctg cccggccacc agaagcgcat cgctacagc 3060  
 ctgctgggac tcaaggacca ggtgaacct gtggggatcc ccatc 3105

<210> 20  
 <211> 1035  
 <212> PRT



<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Predicted fusion protein

<400> 20

Met Lys Lys Ile Met Leu Val Phe Ile Thr Leu Ile Leu Val Ser Leu  
1 5 10 15

Pro Ile Ala Gln Gln Thr Glu Ala Lys Asp Ala Ser Ala Phe Asn Lys  
20 25 30

Glu Asn Ser Ile Ser Ser Met Ala Pro Pro Ala Ser Pro Pro Ala Ser  
35 40 45

Pro Lys Thr Pro Ile Glu Lys Lys His Ala Asp Leu Glu Leu Gln Ala  
50 55 60

Ala Arg Ala Cys Phe Ala Leu Leu Trp Gly Cys Ala Leu Ala Ala Ala  
65 70 75 80

Ala Ala Ala Gln Gly Lys Glu Val Val Leu Leu Asp Phe Ala Ala Ala  
85 90 95

Gly Gly Glu Leu Gly Trp Leu Thr His Pro Tyr Gly Lys Gly Trp Asp  
100 105 110

Leu Met Gln Asn Ile Met Asn Asp Met Pro Ile Tyr Met Tyr Ser Val  
115 120 125

Cys Asn Val Met Ser Gly Asp Gln Asp Asn Trp Leu Arg Thr Asn Trp  
130 135 140

Val Tyr Arg Gly Glu Ala Glu Arg Ile Phe Ile Glu Leu Lys Phe Thr  
145 150 155 160

Val Arg Asp Cys Asn Ser Phe Pro Gly Gly Ala Ser Ser Cys Lys Glu  
165 170 175

Thr Phe Asn Leu Tyr Tyr Ala Glu Ser Asp Leu Asp Tyr Gly Thr Asn  
180 185 190

Phe Gln Lys Arg Leu Phe Thr Lys Ile Asp Thr Ile Ala Pro Asp Glu  
195 200 205

Ile Thr Val Ser Ser Asp Phe Glu Ala Arg His Val Lys Leu Asn Val  
210 215 220

Glu Glu Arg Ser Val Gly Pro Leu Thr Arg Lys Gly Phe Tyr Leu Ala  
 225 230 235 240  
 Phe Gln Asp Ile Gly Ala Cys Val Ala Leu Leu Ser Val Arg Val Tyr  
 245 250 255  
 Tyr Lys Lys Cys Pro Glu Leu Leu Gln Gly Leu Ala His Phe Pro Glu  
 260 265 270  
 Thr Ile Ala Gly Ser Asp Ala Pro Ser Leu Ala Thr Val Ala Gly Thr  
 275 280 285  
 Cys Val Asp His Ala Val Val Pro Pro Gly Gly Glu Glu Pro Arg Met  
 290 295 300  
 His Cys Ala Val Asp Gly Glu Trp Leu Val Pro Ile Gly Gln Cys Leu  
 305 310 315 320  
 Cys Gln Ala Gly Tyr Glu Lys Val Glu Asp Ala Cys Gln Ala Cys Ser  
 325 330 335  
 Pro Gly Phe Phe Lys Phe Glu Ala Ser Glu Ser Pro Cys Leu Glu Cys  
 340 345 350  
 Pro Glu His Thr Leu Pro Ser Pro Glu Gly Ala Thr Ser Cys Glu Cys  
 355 360 365  
 Glu Glu Gly Phe Phe Arg Ala Pro Gln Asp Pro Ala Ser Met Pro Cys  
 370 375 380  
 Thr Arg Pro Pro Ser Ala Pro His Tyr Leu Thr Ala Val Gly Met Gly  
 385 390 395 400  
 Ala Lys Val Glu Leu Arg Trp Thr Pro Pro Gln Asp Ser Gly Gly Arg  
 405 410 415  
 Glu Asp Ile Val Tyr Ser Val Thr Cys Glu Gln Cys Trp Pro Glu Ser  
 420 425 430  
 Gly Glu Cys Gly Pro Cys Glu Ala Ser Val Arg Tyr Ser Glu Pro Pro  
 435 440 445  
 His Gly Leu Thr Arg Thr Ser Val Thr Val Ser Asp Leu Glu Pro His  
 450 455 460

Met Asn Tyr Thr Phe Thr Val Glu Ala Arg Asn Gly Val Ser Gly Leu  
 465 470 475 480

Val Thr Ser Arg Ser Phe Arg Thr Ala Ser Val Ser Ile Asn Gln Thr  
 485 490 495

Glu Pro Pro Lys Val Arg Leu Glu Gly Arg Ser Thr Thr Ser Leu Ser  
 500 505 510

Val Ser Trp Ser Ile Pro Pro Pro Gln Gln Ser Arg Val Trp Lys Tyr  
 515 520 525

Glu Val Thr Tyr Arg Lys Lys Gly Asp Ser Asn Ser Tyr Asn Val Arg  
 530 535 540

Arg Thr Glu Gly Phe Ser Val Thr Leu Asp Asp Leu Ala Pro Asp Thr  
 545 550 555 560

Thr Tyr Leu Val Gln Val Gln Ala Leu Thr Gln Glu Gly Gln Gly Ala  
 565 570 575

Gly Ser Arg Val His Glu Phe Gln Thr Leu Ser Pro Glu Gly Ser Gly  
 580 585 590

Asn Leu Ala Val Ile Gly Gly Val Ala Val Gly Val Val Leu Leu Leu  
 595 600 605

Val Leu Ala Gly Val Gly Phe Phe Ile His Arg Arg Arg Lys Asn Gln  
 610 615 620

Arg Ala Arg Gln Ser Pro Glu Asp Val Tyr Phe Ser Lys Ser Glu Gln  
 625 630 635 640

Leu Lys Pro Leu Lys Thr Tyr Val Asp Pro His Thr Tyr Glu Asp Pro  
 645 650 655

Asn Gln Ala Val Leu Lys Phe Thr Thr Glu Ile His Pro Ser Cys Val  
 660 665 670

Thr Arg Gln Lys Val Ile Gly Ala Gly Glu Phe Gly Glu Val Tyr Lys  
 675 680 685

Gly Met Leu Lys Thr Ser Ser Gly Lys Lys Glu Val Pro Val Ala Ile  
 690 695 700

Lys Thr Leu Lys Ala Gly Tyr Thr Glu Lys Gln Arg Val Asp Phe Leu  
 705 710 715 720  
 Gly Glu Ala Gly Ile Met Gly Gln Phe Ser His His Asn Ile Ile Arg  
 725 730 735  
 Leu Glu Gly Val Ile Ser Lys Tyr Lys Pro Met Met Ile Ile Thr Glu  
 740 745 750  
 Tyr Met Glu Asn Gly Ala Leu Asp Lys Phe Leu Arg Glu Lys Asp Gly  
 755 760 765  
 Glu Phe Ser Val Leu Gln Leu Val Gly Met Leu Arg Gly Ile Ala Ala  
 770 775 780  
 Gly Met Lys Tyr Leu Ala Asn Met Asn Tyr Val His Arg Asp Leu Ala  
 785 790 795 800  
 Ala Arg Asn Ile Leu Val Asn Ser Asn Leu Val Cys Lys Val Ser Asp  
 805 810 815  
 Phe Gly Leu Ser Arg Val Leu Glu Asp Asp Pro Glu Ala Thr Tyr Thr  
 820 825 830  
 Thr Ser Gly Gly Lys Ile Pro Ile Arg Trp Thr Ala Pro Glu Ala Ile  
 835 840 845  
 Ser Tyr Arg Lys Phe Thr Ser Ala Ser Asp Val Trp Ser Phe Gly Ile  
 850 855 860  
 Val Met Trp Glu Val Met Thr Tyr Gly Glu Arg Pro Tyr Trp Glu Leu  
 865 870 875 880  
 Ser Asn His Glu Val Met Lys Ala Ile Asn Asp Gly Phe Arg Leu Pro  
 885 890 895  
 Thr Pro Met Asp Cys Pro Ser Ala Ile Tyr Gln Leu Met Met Gln Cys  
 900 905 910  
 Trp Gln Gln Glu Arg Ala Arg Arg Pro Lys Phe Ala Asp Ile Val Ser  
 915 920 925  
 Ile Leu Asp Lys Leu Ile Arg Ala Pro Asp Ser Leu Lys Thr Leu Ala  
 930 935 940

Asp Phe Asp Pro Arg Val Ser Ile Arg Leu Pro Ser Thr Ser Gly Ser  
945 950 955 960

Glu Gly Val Pro Phe Arg Thr Val Ser Glu Trp Leu Glu Ser Ile Lys  
965 970 975

Met Gln Gln Tyr Thr Glu His Phe Met Ala Ala Gly Tyr Thr Ala Ile  
980 985 990

Glu Lys Val Val Gln Met Thr Asn Asp Asp Ile Lys Arg Ile Gly Val  
995 1000 1005

Arg Leu Pro Gly His Gln Lys Arg Ile Ala Tyr Ser Leu Leu Gly  
1010 1015 1020

Leu Lys Asp Gln Val Asn Thr Val Gly Ile Pro Ile  
1025 1030 1035

<210> 21  
<211> 1506  
<212> DNA  
<213> Homo sapiens

<400> 21  
cagggcaagg aagtgggtact gctggacttt gctgcagctg gaggggagct cggctggctc 60  
acacacccgt atggcaaaagg gtgggacctg atgcagaaca tcatgaatga catgccgatc 120  
taccatgtact ccgtgtgcaa cgtgatgtct ggcgaccagg acaactggct ccgcaccaac 180  
tgggtgtacc gagagaggc tgagcgtatc ttcattgagc tcaagtttac tgtacgtgac 240  
tgcaacagct tccctgggtg cgccagctcc tgcaaggaga ctttcaacct ctactatgcc 300  
gagtcggacc tggactacgg caccaacttc cagaagcgcc tgttcaccaa gattgacacc 360  
attgcgccc atgagatcac cgtcagcagc gacttcgagg cagccacgt gaagctgaac 420  
gtggaggagc gctccgtggg gccgctcacc cgcaaaggct tctacctggc ctccaggat 480  
atcgggtgct gtgtggcgct gctctccgct cgtgtctact acaagaagtg ccccgagctg 540  
ctgcagggcc tggcccaactt ccctgagacc atcgccggct ctgatgcc tccctggcc 600  
actgtggccc gcacctgtgt ggaccatgcc gtgggtgccac cgggggggtga agagcccgct 660  
atgcactgtg cagtggatgg cagtggtgctg gtgcccattg ggcagtgctt gtgccaggca 720  
ggctacgaga aggtggagga tgccctgccag gcctgctcgc ctggattttt taagtgtgag 780  
gcactctgaga gccctgtcct ggagtgcctt gagcacacgc tgccatcccc tgagggtgcc 840  
acctcctgcg agtgtgagga aggccttctc cgggcacctc aggaccacgc gtcgatgcct 900

tgccacacgac cccctccgc cccacactac ctcacagccg tgggcatggg tgccaaggtg	960
gagctgcgct ggacgcccc tcaggacagc gggggccgcg agggacattgt ctacacgctc	1020
acctgcgaac agtgcctggc cgagctctggg gaatgcgggc cgtgtgagcg cagtgtgcgc	1080
tactcggagc ctcctcaagg aetgacccgc accagtgta cagtgcgca cctggagccc	1140
cacatgaact acaccttcac cgtggaggcc cgcaatggcg tctcaggcct ggtaaccagc	1200
cgcagcttcc gtactgccag tgtcagcatc aaccagacag agccccccaa ggtgaggctg	1260
gagggccgca gcaccacctc gcttagcgct tcctggagca tccccccgc gcagcagagc	1320
cgagtgtgga agtacgaggt cacttacgc aagaaggag actccaacag ctacaatgtg	1380
cgccgcacgg agggtttctc cgtgacctg gacgacctgg cccagacac cacctacctg	1440
gtccaggtgc aggcactgac gcaggagggc cagggggccg gcagcagggt gcacgaatc	1500
cagacg	1506

<210> 22  
 <211> 1506  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Human sequence optimized for codon usage in *Listeria*

<400> 22	
caaggtaaag aagttgtttt attagatttt gcagcagcag gtggtgaatt aggttggtta	60
acacatccat atggtaaagg ttgggattta atgcaaaata ttatgaatga tatgccaatt	120
tatatgtata gtgtttgtaa tgttatgagt ggtgatcaag ataattggtt acgtacaaat	180
tgggtttatc gtggtgaagc agaacgtatt tttattgaat taaaatttac agttcgtgat	240
tgtaaatagt ttccagtggt tgcaagtagt tgtaagaaa catttaattt atattatgca	300
gaaagtgtatt tagattatgg tacaaatttt caaaaacggt tatttacaaa aattgatata	360
attgcaccag atgaaattac agtttagtagt gatattgaag cacgtcatgt taaattaaat	420
gttgaaagac gtatgtgttg tccattaaca cgtaaagggt tttatttagc atttcaagat	480
attggtgcac gtgttgcat attaagtgtt cgtgtttatt ataaaaatg tccagaatta	540
ttacaagggt tagcacattt tccagaaaca attgcaggta gtgatgcacc aagtttagca	600
acagttgcag gtacatgtgt tgatcatgca gttgttccac caggtgggta agaaccacgt	660
atgcattgtg cagttgatgg tgaatggta gttccaattg gtcaatgttt atgtcaagca	720
ggttatgaaa aagttgaaga tgcattgtca gcatgtagtc caggtttttt taaatttgaa	780
gcaagtgaat gtccatgttt agaattgtca gaacatacat taccaagtcc agaaggtgca	840

acaagttgtg aatgtgaaga aggttttttt cgtgcaccac aagatccagc aagtatgcc 900  
 tgtacacgtc caccaagtc accacattat ttaacagcag ttggtatggg tgcaaaagtt 960  
 gaattacgtt ggacaccacc acaagatagt ggtggtcgtg aagatatgtt ttatagtgtt 1020  
 acatgtgaac aatgttgccc agaaagtggg gaatgtggtc catgtgaagc aagtgttcgt 1080  
 tatagtgaac caccacatgg tttaacacgt acaagtgtta cagttagtga tttagaacca 1140  
 catatgaatt atacatttac agttgaagca cgtaatgggt ttagtggttt agttacaagt 1200  
 cgtagttttc gtacagcaag tgtagtatt aatcaaacag aaccacaaaa agttcgttta 1260  
 gaaggtcgta gtacaacaag tttaagtgtt agttggagta ttccaccacc acaacaaagt 1320  
 cgtgtttgga aatatgaagt tacatatcgt aaaaaaggtg atagtaatag ttataatgtt 1380  
 cgtcgtacag aagggttttg tgttacatta gatgatttag caccagatac aacatattta 1440  
 gttcaagttc aagcattaac acaagaaggt caaggtgcag gtagtcgtgt tcatgaattt 1500  
 caaaca 1506

<210> 23  
 <211> 502  
 <212> PRT  
 <213> Homo sapeins

<400> 23

Gln Gly Lys Glu Val Val Leu Leu Asp Phe Ala Ala Ala Gly Gly Glu  
 1 5 10 15

Leu Gly Trp Leu Thr His Pro Tyr Gly Lys Gly Trp Asp Leu Met Gln  
 20 25 30

Asn Ile Met Asn Asp Met Pro Ile Tyr Met Tyr Ser Val Cys Asn Val  
 35 40 45

Met Ser Gly Asp Gln Asp Asn Trp Leu Arg Thr Asn Trp Val Tyr Arg  
 50 55 60

Gly Glu Ala Glu Arg Ile Phe Ile Glu Leu Lys Phe Thr Val Arg Asp  
 65 70 75 80

Cys Asn Ser Phe Pro Gly Gly Ala Ser Ser Cys Lys Glu Thr Phe Asn  
 85 90 95

Leu Tyr Tyr Ala Glu Ser Asp Leu Asp Tyr Gly Thr Asn Phe Gln Lys  
 100 105 110

Arg Leu Phe Thr Lys Ile Asp Thr Ile Ala Pro Asp Glu Ile Thr Val  
 115 120 125

Ser Ser Asp Phe Glu Ala Arg His Val Lys Leu Asn Val Glu Glu Arg  
 130 135 140

Ser Val Gly Pro Leu Thr Arg Lys Gly Phe Tyr Leu Ala Phe Gln Asp  
 145 150 155 160

Ile Gly Ala Cys Val Ala Leu Leu Ser Val Arg Val Tyr Tyr Lys Lys  
 165 170 175

Cys Pro Glu Leu Leu Gln Gly Leu Ala His Phe Pro Glu Thr Ile Ala  
 180 185 190

Gly Ser Asp Ala Pro Ser Leu Ala Thr Val Ala Gly Thr Cys Val Asp  
 195 200 205

His Ala Val Val Pro Pro Gly Gly Glu Glu Pro Arg Met His Cys Ala  
 210 215 220

Val Asp Gly Glu Trp Leu Val Pro Ile Gly Gln Cys Leu Cys Gln Ala  
 225 230 235 240

Gly Tyr Glu Lys Val Glu Asp Ala Cys Gln Ala Cys Ser Pro Gly Phe  
 245 250 255

Phe Lys Phe Glu Ala Ser Glu Ser Pro Cys Leu Glu Cys Pro Glu His  
 260 265 270

Thr Leu Pro Ser Pro Glu Gly Ala Thr Ser Cys Glu Cys Glu Glu Gly  
 275 280 285

Phe Phe Arg Ala Pro Gln Asp Pro Ala Ser Met Pro Cys Thr Arg Pro  
 290 295 300

Pro Ser Ala Pro His Tyr Leu Thr Ala Val Gly Met Gly Ala Lys Val  
 305 310 315 320

Glu Leu Arg Trp Thr Pro Pro Gln Asp Ser Gly Gly Arg Glu Asp Ile  
 325 330 335

Val Tyr Ser Val Thr Cys Glu Gln Cys Trp Pro Glu Ser Gly Glu Cys  
 340 345 350

Gly Pro Cys Glu Ala Ser Val Arg Tyr Ser Glu Pro Pro His Gly Leu



355

360

365

Thr Arg Thr Ser Val Thr Val Ser Asp Leu Glu Pro His Met Asn Tyr  
370 375 380

Thr Phe Thr Val Glu Ala Arg Asn Gly Val Ser Gly Leu Val Thr Ser  
385 390 395 400

Arg Ser Phe Arg Thr Ala Ser Val Ser Ile Asn Gln Thr Glu Pro Pro  
405 410 415

Lys Val Arg Leu Glu Gly Arg Ser Thr Thr Ser Leu Ser Val Ser Trp  
420 425 430

Ser Ile Pro Pro Pro Gln Gln Ser Arg Val Trp Lys Tyr Glu Val Thr  
435 440 445

Tyr Arg Lys Lys Gly Asp Ser Asn Ser Tyr Asn Val Arg Arg Thr Glu  
450 455 460

Gly Phe Ser Val Thr Leu Asp Asp Leu Ala Pro Asp Thr Thr Tyr Leu  
465 470 475 480

Val Gln Val Gln Ala Leu Thr Gln Glu Gly Gln Gly Ala Gly Ser Arg  
485 490 495

Val His Glu Phe Gln Thr  
500

<210> 24

<211> 1689

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Fusion protein construct

<400> 24

atgaaaaaaa taatgctagt ttttattaca cttatattag ttagtctacc aattgcgcaa 60

caaaactgaag caaaggatgc atctgcattc aataaagaaa attcaatttc atccatggca 120

ccaccagcat ctccgcctgc aagtcctaag acgccaatcg aaaagaaaca cgcggatctc 180

gagcagggca aggaagtggc actgctggac ttgtctgcag ctggagggga gctcggctgg 240

ctcacacacc cgtatggcaa aggggtggac ctgatgcaga acatcatgaa tgacatgccg 300

atctacatgt actccgtgtg caacgtgatg tctggcgacc aggacaactg gctccgcacc 360

aactgggtgt accgaggaga ggctgagcgt atcttcattg agctcaagtt tactgtacgt 420

gactgcaaca gcttccctgg tggcgccagc tectgcaagg agactttcaa ccttactat 480  
gccgagtcgg acctggacta cggcaccaac ttcagaagc gcctgttcac caagattgac 540  
accattgcgc ccgatgagat caccgtcagc agcgacttcg aggcacgcc a cgtgaagctg 600  
aacgtggagg agcgcctcgt ggggcgcgtc acccgcaaa gcttctacct ggcttccag 660  
gatatcgggt cctgtgtggc gctgctctcc gtccgtgtct actacaagaa gtgccccgag 720  
ctgctgcagg gcctggccca ctccctgag accatgcgc gctctgatgc accttccctg 780  
gccactgtgg ccggcacctg tgtggacat gccgtggtgc caccgggggg tgaagagccc 840  
cgtatgcact gtgcagtggg tggcgagtgg ctgggtgccca ttgggcagtg cctgtgccag 900  
gcaggetacg agaaggtgga ggatgcctgc caggcctgct cgcctggatt ttttaagttt 960  
gaggcatctg agagccctcg cttggagtgc cctgagcaca cgctgccate ccctgagggg 1020  
gccacctctc gcgagtgtga ggaaggett ctcggggcac ctcaggaccc agcgctgatg 1080  
ccttgccacg gacccccctc cgcgccacac tacctcacag ccgtgggcag ggggtgccaa 1140  
gtggagctgc gctggacgcc cctcaggac agcggggggc gcgaggacat tgtctacagc 1200  
gtccactcgc aacagtgtgc gcccgagtct ggggaatgcg ggccgtgtga ggcagtggtg 1260  
cgctactcgg agcctcctca cggactgacc cgcaccagtg tgacagttag cgcactggag 1320  
ccccacatga actacacctt caccgtggag gcccgcaatg cgtctcagg cctggtaacc 1380  
agccgcagct tccgtactgc cagtgtcagc atcaaccaga cagagcccc caaggtgagg 1440  
ctggaggggc gcagcaccac ctgccttagc gtctcctgga gcaccccc gccgcagcag 1500  
agccgagtg ggaagtacga ggtcacttac cgcaagaagg gagactccaa cagctacaat 1560  
gtgcgccga ccgaggggtt ctccgtgacc ctggacgacc tggccccaga caccacctac 1620  
ctgttcagg tgcaggcact gacgcaggag ggccaggggg ccggcagcag ggtgcacgaa 1680  
ttccagacg 1689

<210> 25

<211> 563

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Predicted fusion protein

<400> 25

Met Lys Lys Ile Met Leu Val Phe Ile Thr Leu Ile Leu Val Ser Leu  
1 5 10 15

Pro Ile Ala Gln Gln Thr Glu Ala Lys Asp Ala Ser Ala Phe Asn Lys

20

25

30

Glu Asn Ser Ile Ser Ser Met Ala Pro Pro Ala Ser Pro Pro Ala Ser  
 35 40 45  
 Pro Lys Thr Pro Ile Glu Lys Lys His Ala Asp Leu Glu Gln Gly Lys  
 50 55 60  
 Glu Val Val Leu Leu Asp Phe Ala Ala Ala Gly Gly Glu Leu Gly Trp  
 65 70 75 80  
 Leu Thr His Pro Tyr Gly Lys Gly Trp Asp Leu Met Gln Asn Ile Met  
 85 90 95  
 Asn Asp Met Pro Ile Tyr Met Tyr Ser Val Cys Asn Val Met Ser Gly  
 100 105 110  
 Asp Gln Asp Asn Trp Leu Arg Thr Asn Trp Val Tyr Arg Gly Glu Ala  
 115 120 125  
 Glu Arg Ile Phe Ile Glu Leu Lys Phe Thr Val Arg Asp Cys Asn Ser  
 130 135 140  
 Phe Pro Gly Gly Ala Ser Ser Cys Lys Glu Thr Phe Asn Leu Tyr Tyr  
 145 150 155 160  
 Ala Glu Ser Asp Leu Asp Tyr Gly Thr Asn Phe Gln Lys Arg Leu Phe  
 165 170 175  
 Thr Lys Ile Asp Thr Ile Ala Pro Asp Glu Ile Thr Val Ser Ser Asp  
 180 185 190  
 Phe Glu Ala Arg His Val Lys Leu Asn Val Glu Glu Arg Ser Val Gly  
 195 200 205  
 Pro Leu Thr Arg Lys Gly Phe Tyr Leu Ala Phe Gln Asp Ile Gly Ala  
 210 215 220  
 Cys Val Ala Leu Leu Ser Val Arg Val Tyr Tyr Lys Lys Cys Pro Glu  
 225 230 235 240  
 Leu Leu Gln Gly Leu Ala His Phe Pro Glu Thr Ile Ala Gly Ser Asp  
 245 250 255  
 Ala Pro Ser Leu Ala Thr Val Ala Gly Thr Cys Val Asp His Ala Val

260	265	270
Val Pro Pro Gly Gly Glu Glu Pro Arg Met His Cys Ala Val Asp Gly		
275	280	285
Glu Trp Leu Val Pro Ile Gly Gln Cys Leu Cys Gln Ala Gly Tyr Glu		
290	295	300
Lys Val Glu Asp Ala Cys Gln Ala Cys Ser Pro Gly Phe Phe Lys Phe		
305	310	315
Glu Ala Ser Glu Ser Pro Cys Leu Glu Cys Pro Glu His Thr Leu Pro		
	325	330
Ser Pro Glu Gly Ala Thr Ser Cys Glu Cys Glu Glu Gly Phe Phe Arg		
	340	345
Ala Pro Gln Asp Pro Ala Ser Met Pro Cys Thr Arg Pro Pro Ser Ala		
	355	360
Pro His Tyr Leu Thr Ala Val Gly Met Gly Ala Lys Val Glu Leu Arg		
370	375	380
Trp Thr Pro Pro Gln Asp Ser Gly Gly Arg Glu Asp Ile Val Tyr Ser		
385	390	395
Val Thr Cys Glu Gln Cys Trp Pro Glu Ser Gly Glu Cys Gly Pro Cys		
	405	410
Glu Ala Ser Val Arg Tyr Ser Glu Pro Pro His Gly Leu Thr Arg Thr		
	420	425
Ser Val Thr Val Ser Asp Leu Glu Pro His Met Asn Tyr Thr Phe Thr		
	435	440
Val Glu Ala Arg Asn Gly Val Ser Gly Leu Val Thr Ser Arg Ser Phe		
450	455	460
Arg Thr Ala Ser Val Ser Ile Asn Gln Thr Glu Pro Pro Lys Val Arg		
465	470	475
Leu Glu Gly Arg Ser Thr Thr Ser Leu Ser Val Ser Trp Ser Ile Pro		
	485	490
Pro Pro Gln Gln Ser Arg Val Trp Lys Tyr Glu Val Thr Tyr Arg Lys		
500	505	510

Lys Gly Asp Ser Asn Ser Tyr Asn Val Arg Arg Thr Glu Gly Phe Ser  
515 520 525

Val Thr Leu Asp Asp Leu Ala Pro Asp Thr Thr Tyr Leu Val Gln Val  
530 535 540

Gln Ala Leu Thr Gln Glu Gly Gln Gly Ala Gly Ser Arg Val His Glu  
545 550 555 560

Phe Gln Thr

<210> 26  
<211> 1989  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Fusion protein construct

<400> 26  
ggtacctcct ttgattagta tattcctatc tttaaagttac ttttatgtgg aggcattaac 60  
atattgttaat gacgtcaaaa ggatagcaag actagaataa agctataaag caagcatata 120  
atattgcgtt tcattctttag aagcgaattt cgccaatatt ataattatca aaagagaggg 180  
gtggcaaacg gtatttggca ttattaggtt aaaaaatgta gaaggagatg gaaacccatg 240  
aaaaaaaaaa tgctagtttt tattacactt atattagtta gtctaccaat tgcgcaacaa 300  
actgaagcaa aggatgcacg tgcattcaat aaagaaaatt caatttcacg catggcacca 360  
ccagcatctc cgcctgcaag tcctaagacg ccaatcgaaa agaaacacg ggatggatcc 420  
gattataaag atgatgatga taacaaggtt aaagaagttg ttttattaga ttttcgacga 480  
  
gcagggtggtg aattaggttg gtaaacacat ccatatggta aagggtggga tttaatgcaa 540  
aatattatga atgatatgcc aatttatatg tatagtgttt gtaatgttat gagtggatgat 600  
caagataaatt ggttacgtac aaattgggtt tatcgtgggtg aagcagaacg tatttttatt 660  
gaattaaaat ttacagttcg tgattgtaat agttttccag tgggtgcaag tagttgtaaa 720  
gaaacattta atttatatta tgcagaaagt gatttagatt atggtacaaa ttttcaaaaa 780  
cgttttatta caaaaattga tacaattgca ccagatgaaa ttacagttag tagtgatttt 840  
gaagcacgtc atgttaaaat aaatgttgaa gaacgtagtg ttggtccatt aacacgtaaa 900  
ggtttttatt tagcatttca agatattggt gcatgtgttg cattattaag tgttcgtggt 960

tattataaaa aatgtccaga attattacaa ggtttagcac attttccaga aacaattgca 1020  
 ggtagtgatg caccaagttt agcaacagtt gcagggtacat gtgttgatca tgcagttggt 1080  
 ccaccaggtg gtgaagaacc acgtatgcat tgtgcagttg atggtgaatg gttagtcca 1140  
 attggccaat gtttatgtca agcagggttat gaaaaagttg aagatgcatg tcaagcatgt 1200  
 agtcagggtt tttttaaatt tgaagcaagt gaaagttccat gtttagaatg tccagaacat 1260  
 acattaccaa gtccagaagg tgcaacaagt tgtgaatgtg aagaagggtt ttttcgtgca 1320  
 ccacaagatc cagcaagtat gccatgtaca cgtccaccaa gtgcaccaca ttatttaaca 1380  
 gcagttggta tgggtgcaaa agttgaatta cgttggacac caccacaaga tagtgggtggt 1440  
 cgtgaagata ttgtttatag tgttacatgt gaacaatgtt ggccagaaaag tggtagaatgt 1500  
 ggtccatgtg aagcaagtgt tcgttatagt gaaccaccac atggtttaac acgtacaagt 1560  
 gttacagtta gtgatttaga accacatatg aattatacat ttacagttag agcacgtaat 1620  
 ggtgttagtg gtttagttac aagtcgtagt ttctgtacag caagtgttag tattaatcaa 1680  
 acagaaccac caaaagtctg tttagaaggt cgtagtacaa caagtttaag tgttagttgg 1740  
 agtattccac caccacaaca aagtcgtggt tggaaatatg aagttacata tcgtaaaaaa 1800  
 ggtgatagta atagttataa tgttcgtcgt acagaagggt ttagtgttac attagatgat 1860  
 ttagaccag atacaacata tttagttaa gtccaagcat taacacaaga aggtcaaggt 1920  
 gcaggtagtc gtgttcatga atttcaaaca gaacaaaaat taattagtga agaagattta 1980  
 tgagagctc 1989

<210> 27  
 <211> 581  
 <212> PRT  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Predicted fusion protein

<400> 27

Met Lys Lys Ile Met Leu Val Phe Ile Thr Leu Ile Leu Val Ser Leu  
 1 5 10 15

Pro Ile Ala Gln Gln Thr Glu Ala Lys Asp Ala Ser Ala Phe Asn Lys  
 20 25 30

Glu Asn Ser Ile Ser Ser Met Ala Pro Pro Ala Ser Pro Pro Ala Ser  
 35 40 45

Pro Lys Thr Pro Ile Glu Lys Lys His Ala Asp Gly Ser Asp Tyr Lys  
 50 55 60

Asp Asp Asp Asp Lys Gln Gly Lys Glu Val Val Leu Leu Asp Phe Ala  
 65 70 75 80

Ala Ala Gly Gly Glu Leu Gly Trp Leu Thr His Pro Tyr Gly Lys Gly  
 85 90 95

Trp Asp Leu Met Gln Asn Ile Met Asn Asp Met Pro Ile Tyr Met Tyr  
 100 105 110

Ser Val Cys Asn Val Met Ser Gly Asp Gln Asp Asn Trp Leu Arg Thr  
 115 120 125

Asn Trp Val Tyr Arg Gly Glu Ala Glu Arg Ile Phe Ile Glu Leu Lys  
 130 135 140

Phe Thr Val Arg Asp Cys Asn Ser Phe Pro Gly Gly Ala Ser Ser Cys  
 145 150 155 160

Lys Glu Thr Phe Asn Leu Tyr Tyr Ala Glu Ser Asp Leu Asp Tyr Gly  
 165 170 175

Thr Asn Phe Gln Lys Arg Leu Phe Thr Lys Ile Asp Thr Ile Ala Pro  
 180 185 190

Asp Glu Ile Thr Val Ser Ser Asp Phe Glu Ala Arg His Val Lys Leu  
 195 200 205

Asn Val Glu Glu Arg Ser Val Gly Pro Leu Thr Arg Lys Gly Phe Tyr  
 210 215 220

Leu Ala Phe Gln Asp Ile Gly Ala Cys Val Ala Leu Leu Ser Val Arg  
 225 230 235 240

Val Tyr Tyr Lys Lys Cys Pro Glu Leu Leu Gln Gly Leu Ala His Phe  
 245 250 255

Pro Glu Thr Ile Ala Gly Ser Asp Ala Pro Ser Leu Ala Thr Val Ala  
 260 265 270

Gly Thr Cys Val Asp His Ala Val Val Pro Pro Gly Gly Glu Glu Pro  
 275 280 285

Arg Met His Cys Ala Val Asp Gly Glu Trp Leu Val Pro Ile Gly Gln  
 290 295 300

Cys Leu Cys Gln Ala Gly Tyr Glu Lys Val Glu Asp Ala Cys Gln Ala  
 305 310 315 320

Cys Ser Pro Gly Phe Phe Lys Phe Glu Ala Ser Glu Ser Pro Cys Leu  
 325 330 335

Glu Cys Pro Glu His Thr Leu Pro Ser Pro Glu Gly Ala Thr Ser Cys  
 340 345 350

Glu Cys Glu Glu Gly Phe Phe Arg Ala Pro Gln Asp Pro Ala Ser Met  
 355 360 365

Pro Cys Thr Arg Pro Pro Ser Ala Pro His Tyr Leu Thr Ala Val Gly  
 370 375 380

Met Gly Ala Lys Val Glu Leu Arg Trp Thr Pro Pro Gln Asp Ser Gly  
 385 390 395 400

Gly Arg Glu Asp Ile Val Tyr Ser Val Thr Cys Glu Gln Cys Trp Pro  
 405 410 415

Glu Ser Gly Glu Cys Gly Pro Cys Glu Ala Ser Val Arg Tyr Ser Glu  
 420 425 430

Pro Pro His Gly Leu Thr Arg Thr Ser Val Thr Val Ser Asp Leu Glu  
 435 440 445

Pro His Met Asn Tyr Thr Phe Thr Val Glu Ala Arg Asn Gly Val Ser  
 450 455 460

Gly Leu Val Thr Ser Arg Ser Phe Arg Thr Ala Ser Val Ser Ile Asn  
 465 470 475 480

Gln Thr Glu Pro Pro Lys Val Arg Leu Glu Gly Arg Ser Thr Thr Ser  
 485 490 495

Leu Ser Val Ser Trp Ser Ile Pro Pro Pro Gln Gln Ser Arg Val Trp  
 500 505 510

Lys Tyr Glu Val Thr Tyr Arg Lys Lys Gly Asp Ser Asn Ser Tyr Asn  
 515 520 525

Val Arg Arg Thr Glu Gly Phe Ser Val Thr Leu Asp Asp Leu Ala Pro  
 530 535 540



Asp Thr Thr Tyr Leu Val Gln Val Gln Ala Leu Thr Gln Glu Gly Gln  
545 550 555 560

Gly Ala Gly Ser Arg Val His Glu Phe Gln Thr Glu Gln Lys Leu Ile  
565 570 575

Ser Glu Glu Asp Leu  
580

<210> 28  
<211> 1989  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Construct for fusion protein

<400> 28  
ggtacctcct ttgattagta tattectatc ttaaagttac ttttatgtgg aggcattaac 60  
atttgtaaat gacgtcaaaa ggatagcaag actagaataa agctataaag caagcatata 120  
atattgcgtt tcatcttttg aagcgaattt cgccaatatt ataattatca aaagagaggg 180  
gtggcaaacg gtatttggca ttattaggtt aaaaaatgta gaaggagagt gaaacccatg 240  
aaaaaaatta tgtagttttt tattacatta attttagtta gtttaccatg tgcacaacaa 300  
acagaagcaa aagatgcaag tgcattttaat aaagaaaata gtattagtag tatggcacca 360  
ccagcaagtc caccagcaag tccaaaaaca ccaattgaaa aaaaacatgc agatggatcc 420  
gattataaag atgatgatga taaacaaggt aaagaagttg ttttattaga ttttcgacga 480  
gcaggtgggtg aattaggttg gttacacat ccatatggta aaggttggga tttaatgcaa 540  
aatattatga atgatatgcc aatttatatg tatagtgttt gtaatgttat gagtgggtgat 600  
caagataaatt ggttacgtac aaattgggtt tatcgtgggt aagcagaacg tatttttatt 660  
gaattaaaaa ttacagtctg tgattgtaat agttttccag gtggtgcaag tagttgtaaa 720  
gaaacattta atttatatta tgcagaaagt gatttagatt atggtaacaa ttttcaaaaa 780  
cgtttattta caaaaattga tacaattgca ccagatgaaa ttacagttag tagtgatttt 840  
gaagcacgtc atgttaaat aaatgttgaa gaacgtatgt ttggtccatt aacacgtaaa 900  
ggtttttatt tagcatttca agatattgggt gcatgtgttg cattattaag tgttcgtgtt 960  
tattataaaa aatgtccaga attattacaa ggtttagcac attttccaga aacaattgca 1020  
ggtagtgtat caccaagttt agcaacagtt gcaggtacat gtgttgatca tgcagttgtt 1080  
ccaccaggtg gtgaagaacc acgtatgcat tgtgcagttg atggtgaaat gtttagttcca 1140  
attggtcaat gtttatgtca agcaggttat gaaaaagttg aagatgcagt tcaagcatgt 1200

agtccaggtt tttttaaatt tgaagcaagt gaaagtcctat gtttagaatg tccagaacat 1260  
 acattaccaa gtccagaagg tgcaacaagt tgtgaatgtg aagaagggtt ttttcgtgca 1320  
 ccacaagatc cagcaagtat gccatgtaca cgtccaccaa gtgcaccaca ttatttaaca 1380  
 gcagttggta tgggtgcaaa agttgaatta cgttggacac caccacaaga tagtggtggt 1440  
 cgtgaagata ttgtttatag tgttacatgt gaacaatggt ggccagaaaag tggatgaatgt 1500  
 ggtccatgtg aagcaagtgt tcgttatagt gaaccaccac atgggtttaac acgtacaagt 1560  
 gttacagtta gtgatttaga accacatatg aattatacat ttacagttag agcacgtaat 1620  
 ggtgttagtg gtttagttac aagtcgtagt tttcgtacag caagtgttag tattaatcaa 1680  
 acagaaccac caaaagttcg tttagaaggt cgtagtacaa caagtttaag tgtagtattg 1740  
 agtattccac caccacaaca aagtcgtggt tggaaatatg aagttacata tcgtaaaaaa 1800  
 ggtgatagta atagttataa tgttcgtcgt acagaagggt ttagtgttac attagatgat 1860  
 tttagcaccag atacaacata tttagttcaa gttcaagcat taacacaaga aggtcaagggt 1920  
 gcaggtagtc gtgttcatga atttcaaaca gaacaaaaat taattagtga agaagattta 1980  
 tgagagctc 1989

<210> 29  
 <211> 581  
 <212> PRT  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Predicted Fusion protein

<400> 29

Met Lys Lys Ile Met Leu Val Phe Ile Thr Leu Ile Leu Val Ser Leu  
 1 5 10 15

Pro Ile Ala Gln Gln Thr Glu Ala Lys Asp Ala Ser Ala Phe Asn Lys  
 20 25 30

Glu Asn Ser Ile Ser Ser Met Ala Pro Pro Ala Ser Pro Pro Ala Ser  
 35 40 45

Pro Lys Thr Pro Ile Glu Lys Lys His Ala Asp Gly Ser Asp Tyr Lys  
 50 55 60

Asp Asp Asp Asp Lys Gln Gly Lys Glu Val Val Leu Leu Asp Phe Ala  
 65 70 75 80

Ala Ala Gly Gly Glu Leu Gly Trp Leu Thr His Pro Tyr Gly Lys Gly  
 85 90 95

Trp Asp Leu Met Gln Asn Ile Met Asn Asp Met Pro Ile Tyr Met Tyr  
 100 105 110

Ser Val Cys Asn Val Met Ser Gly Asp Gln Asp Asn Trp Leu Arg Thr  
 115 120 125

Asn Trp Val Tyr Arg Gly Glu Ala Glu Arg Ile Phe Ile Glu Leu Lys  
 130 135 140

Phe Thr Val Arg Asp Cys Asn Ser Phe Pro Gly Gly Ala Ser Ser Cys  
 145 150 155 160

Lys Glu Thr Phe Asn Leu Tyr Tyr Ala Glu Ser Asp Leu Asp Tyr Gly  
 165 170 175

Thr Asn Phe Gln Lys Arg Leu Phe Thr Lys Ile Asp Thr Ile Ala Pro  
 180 185 190

Asp Glu Ile Thr Val Ser Ser Asp Phe Glu Ala Arg His Val Lys Leu  
 195 200 205

Asn Val Glu Glu Arg Ser Val Gly Pro Leu Thr Arg Lys Gly Phe Tyr  
 210 215 220

Leu Ala Phe Gln Asp Ile Gly Ala Cys Val Ala Leu Leu Ser Val Arg  
 225 230 235 240

Val Tyr Tyr Lys Lys Cys Pro Glu Leu Leu Gln Gly Leu Ala His Phe  
 245 250 255

Pro Glu Thr Ile Ala Gly Ser Asp Ala Pro Ser Leu Ala Thr Val Ala  
 260 265 270

Gly Thr Cys Val Asp His Ala Val Val Pro Pro Gly Gly Glu Glu Pro  
 275 280 285

Arg Met His Cys Ala Val Asp Gly Glu Trp Leu Val Pro Ile Gly Gln  
 290 295 300

Cys Leu Cys Gln Ala Gly Tyr Glu Lys Val Glu Asp Ala Cys Gln Ala  
 305 310 315 320

Cys Ser Pro Gly Phe Phe Lys Phe Glu Ala Ser Glu Ser Pro Cys Leu  
 325 330 335

Glu Cys Pro Glu His Thr Leu Pro Ser Pro Glu Gly Ala Thr Ser Cys  
 340 345 350

Glu Cys Glu Glu Gly Phe Phe Arg Ala Pro Gln Asp Pro Ala Ser Met  
 355 360 365

Pro Cys Thr Arg Pro Pro Ser Ala Pro His Tyr Leu Thr Ala Val Gly  
 370 375 380

Met Gly Ala Lys Val Glu Leu Arg Trp Thr Pro Pro Gln Asp Ser Gly  
 385 390 395 400

Gly Arg Glu Asp Ile Val Tyr Ser Val Thr Cys Glu Gln Cys Trp Pro  
 405 410 415

Glu Ser Gly Glu Cys Gly Pro Cys Glu Ala Ser Val Arg Tyr Ser Glu  
 420 425 430

Pro Pro His Gly Leu Thr Arg Thr Ser Val Thr Val Ser Asp Leu Glu  
 435 440 445

Pro His Met Asn Tyr Thr Phe Thr Val Glu Ala Arg Asn Gly Val Ser  
 450 455 460

Gly Leu Val Thr Ser Arg Ser Phe Arg Thr Ala Ser Val Ser Ile Asn  
 465 470 475 480

Gln Thr Glu Pro Pro Lys Val Arg Leu Glu Gly Arg Ser Thr Thr Ser  
 485 490 495

Leu Ser Val Ser Trp Ser Ile Pro Pro Pro Gln Gln Ser Arg Val Trp  
 500 505 510

Lys Tyr Glu Val Thr Tyr Arg Lys Lys Gly Asp Ser Asn Ser Tyr Asn  
 515 520 525

Val Arg Arg Thr Glu Gly Phe Ser Val Thr Leu Asp Asp Leu Ala Pro  
 530 535 540

Asp Thr Thr Tyr Leu Val Gln Val Gln Ala Leu Thr Gln Glu Gly Gln  
 545 550 555 560

Gly Ala Gly Ser Arg Val His Glu Phe Gln Thr Glu Gln Lys Leu Ile  
 565 570 575

Ser Glu Glu Asp Leu  
580

<210> 30  
<211> 1968  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Fusion protein construct

<400> 30  
ggtagctcct ttgattagta tattcctatc ttaaagttac ttttatgtgg aggcattaac 60  
atttgtaaat gacgtcaaaa ggatagcaag actagaataa agctataaag caagcatata 120  
atatgcgttt tcattctttag aagcgaattt cgccaatatt ataattatca aaagagaggg 180  
gtggcaaacg gtattttggca ttatttaggtt aaaaaatgta gaaggagagt gaaacccatg 240  
gcatacgaca gtcgttttga tgaatgggta cagaaactga aagaggaaag ctttcaaaac 300  
aatacgtttg accgccgcaa atttattcaa ggagcgggga agattgcagg actttctctt 360  
ggattaacga ttgcccgatc ggttggggcc tttggatccg attataaaga tgatgatgat 420  
aaacaaggtg aagaagttgt tttattagat tttgcagcag caggtgtgta attaggttgg 480  
ttaacacatc catatggtaa aggttgggat ttaatgcaa atattatgaa tgatagcca 540  
atttatatgt atagtgtttg taatgttatg agtgggtgatc aagataaatt gttacgtaca 600  
aattgggttt atcgtgttga agcagaacgt atttttattg aattaaaatt tacagttcgt 660  
gattgtaata gttttccagg tgggtcaagt agttgtaaa aacattttaa tttatattat 720  
gcagaaagtg atttagatta tggtaacaa tttcaaaaac gtttatttac aaaaattgat 780  
acaattgcac cagatgaaat tacagttagt agtgattttg aagcacgtca tggtaaatga 840  
aatgttgaag aacgtagtgt tgggtccatta acacgtaaa gtttttattt agcatttcaa 900  
gatattgggt catgtgttgc attattaagt gttcgtgttt attataaaaa atgtccagaa 960  
ttattacaag gtttagcaca ttttccagaa acaattgcag gtatgtatgc accaagttaa 1020  
gcaacagttg caggtacatg tgttgatcat gcagttgttc caccaggtgg tgaagaacca 1080  
cgtatgcatt gtgcagttga tgggtgaatgg ttagttccaa ttggtcaatg tttatgtcaa 1140  
gcaggttatg aaaaagttga agatgcatgt caagcatgta gtccaggttt ttttaaatgt 1200  
gaagcaagtg aaagtccatg tttagaatgt ccagaacata cattaccaag tccagaaggt 1260  
gcaacaagtt gtgaatgtga agaaggtttt tttcgtgcac cacaagatcc agcaagtatg 1320  
ccatgtacac gtccaccaag tgcaccacat tatttaacag cagttggtat ggggtcaaaa 1380  
gttgaaattc gttggacacc accacaagat agtgggtgtc gtgaagatat tgtttatagt 1440

Ser Glu Glu Asp Leu  
580

<210> 30  
<211> 1968  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Fusion protein construct

<400> 30  
ggtacctect ttgattagta tattcctatc ttaaggttac ttttatgtgg aggcattaac 60  
at ttgttaat gacgtcaaaa ggatagcaag actagaataa agctataaag caagcatata 120  
atattgcggt tcatcttttag aagcgaattt cgccaatatt ataattatca aaagagaggg 180  
gtggcaaacg gtatttggca ttattagggt aaaaaatgta gaaggagagt gaaacccatg 240  
gcatacgaca gtcgttttga tgaatgggta cagaaactga aagaggaaaag ctttcaaaac 300  
aatacgtttg accgccgcaa atttattcaa ggagcgggga agattgcagg actttctctt 360  
ggattaacga ttgccagtc ggttggggcc tttggatccg attataaaga tgatgatgat 420  
aaacaaggta aagaagttgt tttattagat ttgcagcag caggtgggtga attaggttgg 480  
ttaacacatc catatggtaa aggttgggat ttaatgcaaa atattatgaa tgatagcca 540  
at ttatatgt atagtgtttg taatgttatg agtgggtgatc aagataatg gttacgtaca 600  
aattgggttt atcgtgggtg agcagaacgt atttttatg aattaaaaat tacagttcgt 660  
gattgtaata gttttccagg tgggtgcaagt agttgtaaa aaacatttaa tttatattat 720  
gcagaaagtg atttagatta tgggtacaaat ttcaaaaaac gttttttac aaaaattgat 780  
acaattgcac cagatgaaat tacagttagt agtgattttg aagcacgtca tgttaaatta 840  
aatgttgaag aacgtagtgt tgggtccatta acacgtaaaag gtttttattt agcatttcaa 900  
gatattgggt catgtgttgc attattaagt gttcgtgttt attataaaaa atgtccagaa 960  
ttattacaag gtttagcaca tttccagaa acaattgcag gtatgtatgc accaagttta 1020  
gcaacagtgt caggtacatg tgttgatcat gcagttgttc caccaggtgt tgaagaacca 1080  
cgtatgcatt gtgcagttga tgggtgaatg ttagttccaa ttggtcaatg tttatgtcaa 1140  
gcaggttatg aaaaagttag agatgcattg caagcatgta gtccagggtt ttttaaaatt 1200  
gaagcaagtg aaagtccatg tttagaatgt ccagaacata cattaccaag tccagaaggt 1260  
gcaacaagtt gtgaatgtga agaaggtttt tttcgtgcac cacaagatcc agcaagtatg 1320  
ccatgtacac gtccaccaag tgcaccacat tatttaacag cagttgggat ggggtcaaaa 1380  
gttgaattac gttggacacc accacaagat agtgggtggc gtgaagatat tgtttatagt 1440

gttacatgtg aacaatgttg gccagaaagt ggtgaatgtg gtccatgtga agcaagtgtt 1500  
 cggtatagtg aaccaccaca tggtttaaca cgtacaagtg ttacagttag tgatttagaa 1560  
 ccacatatga attatacatt tacagttgaa gcacgtaatg gtgttagtgg tttagtata 1620  
 agtcgtagtt ttcgtacagc aagtgttagt attaatcaaa cagaaccacc aaaagtctgt 1680  
 ttagaaggtc gtagtacaac aagtttaagt gttagttgga gtattccacc accacaacaa 1740  
 agtcgtgttt ggaaatatga agttacatat cgtaaaaaag gtgatagtaa tagttataat 1800  
 gttcgtcgta cagaagggtt tagtggtaca ttagatgatt tagcaccaga tacaacatat 1860  
 ttagttcaag ttcaagcatt aacacaagaa ggtcaagggt caggtagtcg tgttcatgaa 1920  
 tttcaaacag aacaaaaatt aattagttaa gaagatttat gagagctc 1968

<210> 31  
 <211> 574  
 <212> PRT  
 <213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Predicted Fusion Protein

<400> 31

Met Ala Tyr Asp Ser Arg Phe Asp Glu Trp Val Gln Lys Leu Lys Glu  
 1 5 10 15

Glu Ser Phe Gln Asn Asn Thr Phe Asp Arg Arg Lys Phe Ile Gln Gly  
 20 25 30

Ala Gly Lys Ile Ala Gly Leu Ser Leu Gly Leu Thr Ile Ala Gln Ser  
 35 40 45

Val Gly Ala Phe Gly Ser Asp Tyr Lys Asp Asp Asp Lys Gln Gly  
 50 55 60

Lys Glu Val Val Leu Leu Asp Phe Ala Ala Ala Gly Gly Glu Leu Gly  
 65 70 75 80

Trp Leu Thr His Pro Tyr Gly Lys Gly Trp Asp Leu Met Gln Asn Ile  
 85 90 95

Met Asn Asp Met Pro Ile Tyr Met Tyr Ser Val Cys Asn Val Met Ser  
 100 105 110

Gly Asp Gln Asp Asn Trp Leu Arg Thr Asn Trp Val Tyr Arg Gly Glu  
 115 120 125

Ala Glu Arg Ile Phe Ile Glu Leu Lys Phe Thr Val Arg Asp Cys Asn  
130 135 140

Ser Phe Pro Gly Gly Ala Ser Ser Cys Lys Glu Thr Phe Asn Leu Tyr  
145 150 155 160

Tyr Ala Glu Ser Asp Leu Asp Tyr Gly Thr Asn Phe Gln Lys Arg Leu  
165 170 175

Phe Thr Lys Ile Asp Thr Ile Ala Pro Asp Glu Ile Thr Val Ser Ser  
180 185 190

Asp Phe Glu Ala Arg His Val Lys Leu Asn Val Glu Glu Arg Ser Val  
195 200 205

Gly Pro Leu Thr Arg Lys Gly Phe Tyr Leu Ala Phe Gln Asp Ile Gly  
210 215 220

Ala Cys Val Ala Leu Leu Ser Val Arg Val Tyr Tyr Lys Lys Cys Pro  
225 230 235 240

Glu Leu Leu Gln Gly Leu Ala His Phe Pro Glu Thr Ile Ala Gly Ser  
245 250 255

Asp Ala Pro Ser Leu Ala Thr Val Ala Gly Thr Cys Val Asp His Ala  
260 265 270

Val Val Pro Pro Gly Gly Glu Glu Pro Arg Met His Cys Ala Val Asp  
275 280 285

Gly Glu Trp Leu Val Pro Ile Gly Gln Cys Leu Cys Gln Ala Gly Tyr  
290 295 300

Glu Lys Val Glu Asp Ala Cys Gln Ala Cys Ser Pro Gly Phe Phe Lys  
305 310 315 320

Phe Glu Ala Ser Glu Ser Pro Cys Leu Glu Cys Pro Glu His Thr Leu  
325 330 335

Pro Ser Pro Glu Gly Ala Thr Ser Cys Glu Cys Glu Glu Gly Phe Phe  
340 345 350

Arg Ala Pro Gln Asp Pro Ala Ser Met Pro Cys Thr Arg Pro Pro Ser  
355 360 365



Ala Pro His Tyr Leu Thr Ala Val Gly Met Gly Ala Lys Val Glu Leu  
370 375 380

Arg Trp Thr Pro Pro Gln Asp Ser Gly Gly Arg Glu Asp Ile Val Tyr  
385 390 395 400

Ser Val Thr Cys Glu Gln Cys Trp Pro Glu Ser Gly Glu Cys Gly Pro  
405 410 415

Cys Glu Ala Ser Val Arg Tyr Ser Glu Pro Pro His Gly Leu Thr Arg  
420 425 430

Thr Ser Val Thr Val Ser Asp Leu Glu Pro His Met Asn Tyr Thr Phe  
435 440 445

Thr Val Glu Ala Arg Asn Gly Val Ser Gly Leu Val Thr Ser Arg Ser  
450 455 460

Phe Arg Thr Ala Ser Val Ser Ile Asn Gln Thr Glu Pro Pro Lys Val  
465 470 475 480

Arg Leu Glu Gly Arg Ser Thr Thr Ser Leu Ser Val Ser Trp Ser Ile  
485 490 495

Pro Pro Pro Gln Gln Ser Arg Val Trp Lys Tyr Glu Val Thr Tyr Arg  
500 505 510

Lys Lys Gly Asp Ser Asn Ser Tyr Asn Val Arg Arg Thr Glu Gly Phe  
515 520 525

Ser Val Thr Leu Asp Asp Leu Ala Pro Asp Thr Thr Tyr Leu Val Gln  
530 535 540

Val Gln Ala Leu Thr Gln Glu Gly Gln Gly Ala Gly Ser Arg Val His  
545 550 555 560

Glu Phe Gln Thr Glu Gln Lys Leu Ile Ser Glu Glu Asp Leu  
565 570

<210> 32

<211> 1254

<212> DNA

<213> Homo sapiens

<400> 32

caccgcagga ggaagaacca gcgtgcccgc cagtccccgg aggacgttta cttctccaag

60

tcagacaac	tgaagccct	gaagacatac	gtggaccccc	acacatatga	ggaccccaac	120
caggctgtgt	tgaagttcac	taccgagatc	catccatcct	gtgtcactcg	gcagaagggtg	180
atcggagcag	gagagtttgg	ggaggtgtac	aaggcgatgc	tgaagacatc	ctcgggggaag	240
aaggaggtgc	cggtagccat	caagacgctg	aaagccggct	acacagagaa	gcagcgagtg	300
gacttcctcg	gcgagggccg	catcatgggc	cagttcagcc	accacaacat	catccgccta	360
gagggcgta	tctccaaata	caagcccatg	atgatcatca	ctgagtacat	ggagaatggg	420
gccctggaca	agttccttcg	ggagaaggat	ggcgagttca	gcgtgctgca	gctggtgggc	480
atgctgcggg	gcacgcgacg	tggcatgaag	tacctggcca	acatgaacta	tgtgcaccgt	540
gacctggctg	cccgcacatc	cctcgtcaac	agcaacctgg	tctgcaaggt	gtctgacttt	600
ggcctgtccc	gcgtgctgga	ggagaccccc	gaggccacct	acaccaccag	tggcggaag	660
atccccatcc	gctggaccgc	cccgagggcc	atttctacc	ggaagttcac	ctctgccacg	720
gacgtgtgga	gctttggcat	tgatcatgtg	gaggtgatga	cctatggcca	gcggccctac	780
tgggagttgt	ccaaccacga	ggtgatgaaa	gccatcaatg	atggcttcgc	gctccccaca	840
cccatggact	gccctccgc	catctaccag	ctcatgatgc	agtgtctggc	gcaggagcgt	900
gcccgcgcgc	ccaagttcgc	tgacatcgtc	agcatcctgg	acaagctcat	tcgtgccctt	960
gactccctca	agaccctggc	tgactttgac	ccccgcgtgt	ctatccggct	ccccagcacg	1020
agcggtctcg	aggggggtgcc	cttccgcacg	gtgtccgagt	ggctggagtc	catcaagatg	1080
cagcagtata	cggagcactt	catggcggcc	ggctacactg	ccatcgagaa	gggtgtgcag	1140
atgaccaacg	acgacatcaa	gaggattggg	gtgcgggtgc	ccggccacca	gaagcgcatc	1200
gcctacagcc	tgctgggcat	caaggaccag	gtgaacactg	tggggatccc	catc	1254

<210> 33

<211> 1254

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Sequence Optimized for codon usage in *Listeria*

<400> 33

cacagacgta	gaaaaaatca	acgtgctcga	caatccccag	aagatgtgta	tttttcgaaa	60
agtgaacaat	taaaaccatt	aaaaacttat	gttgatccgc	atcgtacga	agacccaaat	120
caagcagtat	taaaatttac	aacagaaata	cacccaagtt	gtgttacaga	acaaaagtt	180
attgagcag	gtgaattcgg	agaggtatat	aaaggtatgt	taaaaacatc	atcaggtaaa	240
aaagaagttc	cggttgcaat	taaaacctta	aaggcaggat	atacagaaaa	acagcgagtt	300

gatttttttag gtgaagcagg aattatgggt caatttagcc atcataatat tattcgtttg 360  
 gaaggagtaa taagtaaata taaaccaatg atgattatta cagaatacat ggaaaacggg 420  
 gctttagata aatttttacg tgaagaggat ggtgaattta gtgttttaca attggttggt 480  
 atgttaagag gaattgctgc aggtatgaaa tatttagcta atatgaatta tgttcaccgt 540  
 gatttggcag caagaaatat cctagtcaat tccaatttag tatgtaaagt tagtgatttt 600  
 ggtttaagca gagtattaga agacgatcca gaggcaacct atacaacatc gggagggtaaa 660  
 attcctattc gttggacagc accagaagct atcagttacc gtaaatttac aagtgcattc 720  
 gacgtgtgga gttttgggat tgtaatgtgg gaagttatga catatggaga aagaccatat 780  
 tgggaattaa gtaatcatga agttatgaaa gcaattaacg atggatttag attaccaact 840  
 ccgatggatt gtccatctgc catttatcaa ctaatgatgc aatgttggca acaagaaaga 900  
 gcacgagctc caaattttgc agatattgtt agtatttttag acaaattaat tcgtgcacca 960  
 gatagtttaa aaacttttag agactttgat cctcgtgtta gtattcgatt accaagtacg 1020  
 tcaggttccg aaggagttcc atttcgcaca gtctccgaat ggttggaatc aattaaaaag 1080  
 caacaatata ccgaacactt tatggcgaca ggttacacag caatcgaaaa agttgttcaa 1140  
 atgacaaatg atgatattaa acgtattgga gttagattac caggccacca gaaacgtatt 1200  
 gcatattctt tatttagggtt aaaagatcaa gttaataaccg tgggaattcc aatt 1254

<210> 34  
 <211> 456  
 <212> PRT  
 <213> Homo sapiens

<400> 34

Val His Glu Phe Gln Thr Leu Ser Pro Glu Gly Ser Gly Asn Leu Ala  
 1 5 10 15

Val Ile Gly Gly Val Ala Val Gly Val Val Leu Leu Leu Val Leu Ala  
 20 25 30

Gly Val Gly Phe Phe Ile His Arg Arg Arg Lys Asn Gln Arg Ala Arg  
 35 40 45

Gln Ser Pro Glu Asp Val Tyr Phe Ser Lys Ser Glu Gln Leu Lys Pro  
 50 55 60

Leu Lys Thr Tyr Val Asp Pro His Thr Tyr Glu Asp Pro Asn Gln Ala  
 65 70 75 80

Val Leu Lys Phe Thr Thr Glu Ile His Pro Ser Cys Val Thr Arg Gln

Lys Val Ile Gly Ala Gly Glu Phe Gly Glu Val Tyr Lys Gly Met Leu  
 100 105 110  
 Lys Thr Ser Ser Gly Lys Lys Glu Val Pro Val Ala Ile Lys Thr Leu  
 115 120 125  
 Lys Ala Gly Tyr Thr Glu Lys Gln Arg Val Asp Phe Leu Gly Glu Ala  
 130 135 140  
 Gly Ile Met Gly Gln Phe Ser His His Asn Ile Ile Arg Leu Glu Gly  
 145 150 155 160  
 Val Ile Ser Lys Tyr Lys Pro Met Met Ile Ile Thr Glu Tyr Met Glu  
 165 170 175  
 Asn Gly Ala Leu Asp Lys Phe Leu Arg Glu Lys Asp Gly Glu Phe Ser  
 180 185 190  
 Val Leu Gln Leu Val Gly Met Leu Arg Gly Ile Ala Ala Gly Met Lys  
 195 200 205  
 Tyr Leu Ala Asn Met Asn Tyr Val His Arg Asp Leu Ala Ala Arg Asn  
 210 215 220  
 Ile Leu Val Asn Ser Asn Leu Val Cys Lys Val Ser Asp Phe Gly Leu  
 225 230 235 240  
 Ser Arg Val Leu Glu Asp Asp Pro Glu Ala Thr Tyr Thr Thr Ser Gly  
 245 250 255  
 Gly Lys Ile Pro Ile Arg Trp Thr Ala Pro Glu Ala Ile Ser Tyr Arg  
 260 265 270  
 Lys Phe Thr Ser Ala Ser Asp Val Trp Ser Phe Gly Ile Val Met Trp  
 275 280 285  
 Glu Val Met Thr Tyr Gly Glu Arg Pro Tyr Trp Glu Leu Ser Asn His  
 290 295 300  
 Glu Val Met Lys Ala Ile Asn Asp Gly Phe Arg Leu Pro Thr Pro Met  
 305 310 315 320  
 Asp Cys Pro Ser Ala Ile Tyr Gln Leu Met Met Gln Cys Trp Gln Gln  
 325 330 335

Glu Arg Ala Arg Arg Pro Lys Phe Ala Asp Ile Val Ser Ile Leu Asp  
340 345 350

Lys Leu Ile Arg Ala Pro Asp Ser Leu Lys Thr Leu Ala Asp Phe Asp  
355 360 365

Pro Arg Val Ser Ile Arg Leu Pro Ser Thr Ser Gly Ser Glu Gly Val  
370 375 380

Pro Phe Arg Thr Val Ser Glu Trp Leu Glu Ser Ile Lys Met Gln Gln  
385 390 395 400

Tyr Thr Glu His Phe Met Ala Ala Gly Tyr Thr Ala Ile Glu Lys Val  
405 410 415

Val Gln Met Thr Asn Asp Asp Ile Lys Arg Ile Gly Val Arg Leu Pro  
420 425 430

Gly His Gln Lys Arg Ile Ala Tyr Ser Leu Leu Gly Leu Lys Asp Gln  
435 440 445

Val Asn Thr Val Gly Ile Pro Ile  
450 455

<210> 35  
<211> 1437  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Fusion Protein

<400> 35  
atgaaaaaaa taatgctagt ttttattaca cttatattag ttagtctacc aattgcgcga 60  
caaaactgaag caaaggatgc atctgcattc aataaagaaa attcaatttc atccatggca 120  
ccaccagcat ctccgcctgc aagtccctaag acgccaatcg aaaagaacaa cgcggatctc 180  
gagcaccgca ggagggaagaa ccagcgtgcc cgcagtcgcc cggaggacgt ttactttctc 240  
aagtgcagaac aactgaagcc cctgaagaca tacgtggacc cccacacata tgaggacccc 300  
aaccaggctg tgttgaagtt cactaccgag atccatccat cctgtgtcac tcggcagaag 360  
gtgatcgagg caggagagtt tggggagggtg tacaagggca tgctgaagac atcctcgggg 420  
aagaaggagg tgccggtggc catcaagacg ctgaaagccg gctacacaga gaagcagcga 480  
gtggacttcc tcggcgaggc cggcatcatg ggccagttca gccaccacaa catcatccgc 540

ctagaggcgg tcactctccaa atacaagccc atgatgatca tcaactgagta catggagaat 600  
 ggggcccttg acaagttcct tcgggagaag gatggcgagt tcagcgtgct gcagctgggt 660  
 ggcatgctgc ggggcacgc agctggcatg aagtacctgg ccaacatgaa ctatgtgcac 720  
 cgtgacctgg ctgcccgaac catcctcgtc aacagcaacc tgggtctgcaa ggtgtctgac 780  
 tttggcctgt cccgcgtgct ggaggacgac cccgaggcca cctacaccac cagtggcggc 840  
 aagatcccca tccgctggac cggcccgagg gccatttctt accggaagtt cacctctgcc 900  
 agcgacgtgt ggagcttttg cattgtcatg tgggaggtga tgacctatgg cgagcggccc 960  
 tactgggagt tgtccaacca cgaggtgatg aaagccatca atgatggctt ccggctcccc 1020  
 acacccatgg actgcccctc cgccatctac cagctcatga tgcagtgtct gcagcaggag 1080  
 cgtgcccgcc gccccaagtt cgctgacatc gtcagcatcc tggacaagct cattcgtgcc 1140  
 cctgactccc tcaagacctt ggctgacttt gacccccggt tgtctatccg gctccccagc 1200  
 acgagcgggt cggagggggg gcccttcgcg acggtgtccg agtggctgga gtccatcaag 1260  
 atgcagcagt atacggagca cttcattggc gccggctaca ctgccatcga gaagtggtg 1320  
 cagatgacca acgacgacat caagaggatt ggggtgcggc tgccccgcca ccagaagcgc 1380  
 atcgccctaca gcctgctggg actcaaggac caggtgaaca ctgtggggat ccccatc 1437

<210> 36  
 <211> 479  
 <212> PRT  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Predicted Protein Sequence

<400> 36

Met Lys Lys Ile Met Leu Val Phe Ile Thr Leu Ile Leu Val Ser Leu  
 1 5 10 15

Pro Ile Ala Gln Gln Thr Glu Ala Lys Asp Ala Ser Ala Phe Asn Lys  
 20 25 30

Glu Asn Ser Ile Ser Ser Met Ala Pro Pro Ala Ser Pro Pro Ala Ser  
 35 40 45

Pro Lys Thr Pro Ile Glu Lys Lys His Ala Asp Leu Glu His Arg Arg  
 50 55 60

Arg Lys Asn Gln Arg Ala Arg Gln Ser Pro Glu Asp Val Tyr Phe Ser  
 65 70 75 80

Lys Ser Glu Gln Leu Lys Pro Leu Lys Thr Tyr Val Asp Pro His Thr  
 85 90 95

Tyr Glu Asp Pro Asn Gln Ala Val Leu Lys Phe Thr Thr Glu Ile His  
 100 105 110

Pro Ser Cys Val Thr Arg Gln Lys Val Ile Gly Ala Gly Glu Phe Gly  
 115 120 125

Glu Val Tyr Lys Gly Met Leu Lys Thr Ser Ser Gly Lys Lys Glu Val  
 130 135 140

Pro Val Ala Ile Lys Thr Leu Lys Ala Gly Tyr Thr Glu Lys Gln Arg  
 145 150 155 160

Val Asp Phe Leu Gly Glu Ala Gly Ile Met Gly Gln Phe Ser His His  
 165 170 175

Asn Ile Ile Arg Leu Glu Gly Val Ile Ser Lys Tyr Lys Pro Met Met  
 180 185 190

Ile Ile Thr Glu Tyr Met Glu Asn Gly Ala Leu Asp Lys Phe Leu Arg  
 195 200 205

Glu Lys Asp Gly Glu Phe Ser Val Leu Gln Leu Val Gly Met Leu Arg  
 210 215 220

Gly Ile Ala Ala Gly Met Lys Tyr Leu Ala Asn Met Asn Tyr Val His  
 225 230 235 240

Arg Asp Leu Ala Ala Arg Asn Ile Leu Val Asn Ser Asn Leu Val Cys  
 245 250 255

Lys Val Ser Asp Phe Gly Leu Ser Arg Val Leu Glu Asp Asp Pro Glu  
 260 265 270

Ala Thr Tyr Thr Thr Ser Gly Gly Lys Ile Pro Ile Arg Trp Thr Ala  
 275 280 285

Pro Glu Ala Ile Ser Tyr Arg Lys Phe Thr Ser Ala Ser Asp Val Trp  
 290 295 300

Ser Phe Gly Ile Val Met Trp Glu Val Met Thr Tyr Gly Glu Arg Pro  
 305 310 315 320

Tyr Trp Glu Leu Ser Asn His Glu Val Met Lys Ala Ile Asn Asp Gly  
325 330 335

Phe Arg Leu Pro Thr Pro Met Asp Cys Pro Ser Ala Ile Tyr Gln Leu  
340 345 350

Met Met Gln Cys Trp Gln Gln Glu Arg Ala Arg Arg Pro Lys Phe Ala  
355 360 365

Asp Ile Val Ser Ile Leu Asp Lys Leu Ile Arg Ala Pro Asp Ser Leu  
370 375 380

Lys Thr Leu Ala Asp Phe Asp Pro Arg Val Ser Ile Arg Leu Pro Ser  
385 390 395 400

Thr Ser Gly Ser Glu Gly Val Pro Phe Arg Thr Val Ser Glu Trp Leu  
405 410 415

Glu Ser Ile Lys Met Gln Gln Tyr Thr Glu His Phe Met Ala Ala Gly  
420 425 430

Tyr Thr Ala Ile Glu Lys Val Val Gln Met Thr Asn Asp Asp Ile Lys  
435 440 445

Arg Ile Gly Val Arg Leu Pro Gly His Gln Lys Arg Ile Ala Tyr Ser  
450 455 460

Leu Leu Gly Leu Lys Asp Gln Val Asn Thr Val Gly Ile Pro Ile  
465 470 475

<210> 37

<211> 1737

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Fusion protein sequence

<400> 37

ggtacctcct ttgattagta tattcctatc ttaaagttac ttttatgtgg aggcattaac 60

atattgttaac gacgtcaaaa ggatagcaag actagaataa agctataaag caagcatata 120

atatcgctt tcattctttag aagcgaattt cgccaatatt ataattatca aaagagaggg 180

gtggcaaacg gtatttggca ttattagggt aaaaaatgta gaaggagagt gaaacccatg 240

aaaaaaaaaa tgctagtgtt tattacactt atattagtta gtctaccaat tgcgcaacaa 300



actgaagcaa aggatgcac	tgcatccaat aaagaaaatt caatttcac	catggcacca	360
ccagcatctc cgctgcga	tcctaagacg ccaatcgaaa agaaacacgc	ggatggatcc	420
gattataaag atgatgatga	taaacacaga cgtagaaaaa atcaacgtgc	tcgacaatcc	480
ccagaagatg tgtatttttc	gaaaagtga caattaaaac cattaaaaac	ttatgttgat	540
ccgcatacgt acgaagacc	aaatcaagca gtattaaaat ttacaacaga	aatacaccca	600
agttgtgtta caagacaaaa	agttattgga gcaggtgaat tcggagaggt	atataaaggt	660
atgttaaaaa catcatcagg	taaaaaagaa gtcccggttg caattaaaac	cttaaggcca	720
ggatatacag aaaaacacgc	agttgatttt ttaggtgaag caggaattat	gggtcaattt	780
agccatcata atattattcg	tttgggaagga gtaataagta aatataaacc	aatgatgatt	840
attacagaat acatggaaa	cggtgcttta gataaaattt tacgtgaaa	ggatggtgaa	900
tttagtgttt tacaattggt	tggtatgtta agaggaattg ctgcaggtat	gaaatattta	960
gctaataatga attatgttca	ccgtgatttg gcagcaagaa atatccctagt	caattccaat	1020
ttagtatgta aagttagtga	ttttgggtta agcagagtat tagaagacga	tccagaggca	1080
acctatacaa catcgggagg	taaaattcct attcgttggga cagcaccaga	agctatcagt	1140
taccgtaaat ttacaagtgc	atcacacgtg tggagttttg ggattgtaat	gtgggaagtt	1200
atgacatatg gagaaagacc	atattgggaa ttaagtaatc atgaagttaa	gaaagcaatt	1260
aacgatggat tttagattacc	aactccgatg gattgtccat ctgccattta	tcaactaatg	1320
atgcaatggt ggcaacaaga	aagagcacga cgtccaaaaa ttgcagatat	tgtagtatatt	1380
ttagacaaat taattcgtgc	accagatagt ttaaaaaactt tagcagactt	tgatccctct	1440
gttagtatctc gattaccaag	tacgtcaggt tccgaaggag ttccatttcg	cacagctcc	1500
gaatgggttg aatcaattaa	aatgcaacaa tacaccgaac actttatggc	agcaggttac	1560
acagcaatcg aaaaagttgt	tcaaatgaca aatgatgata ttaaacgtat	tggagttaga	1620
ttaccaggcc accagaaacg	tattgcatat tctttattag gtttaaaaga	tcaagttaat	1680
accgtgggaa ttccaattga	acaaaaatta atttccgaag aagacttata	agagctc	1737

<210> 38

<211> 497

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Predicted fusion protein

<400> 38

Met Lys Lys Ile Met Leu Val Phe Ile Thr Leu Ile Leu Val Ser Leu

1

5

10

15

Pro Ile Ala Gln Gln Thr Glu Ala Lys Asp Ala Ser Ala Phe Asn Lys  
 20 25 30

Glu Asn Ser Ile Ser Ser Met Ala Pro Pro Ala Ser Pro Pro Ala Ser  
 35 40 45

Pro Lys Thr Pro Ile Glu Lys Lys His Ala Asp Gly Ser Asp Tyr Lys  
 50 55 60

Asp Asp Asp Asp Lys His Arg Arg Arg Lys Asn Gln Arg Ala Arg Gln  
 65 70 75 80

Ser Pro Glu Asp Val Tyr Phe Ser Lys Ser Glu Gln Leu Lys Pro Leu  
 85 90 95

Lys Thr Tyr Val Asp Pro His Thr Tyr Glu Asp Pro Asn Gln Ala Val  
 100 105 110

Leu Lys Phe Thr Thr Glu Ile His Pro Ser Cys Val Thr Arg Gln Lys  
 115 120 125

Val Ile Gly Ala Gly Glu Phe Gly Glu Val Tyr Lys Gly Met Leu Lys  
 130 135 140

Thr Ser Ser Gly Lys Lys Glu Val Pro Val Ala Ile Lys Thr Leu Lys  
 145 150 155 160

Ala Gly Tyr Thr Glu Lys Gln Arg Val Asp Phe Leu Gly Glu Ala Gly  
 165 170 175

Ile Met Gly Gln Phe Ser His His Asn Ile Ile Arg Leu Glu Gly Val  
 180 185 190

Ile Ser Lys Tyr Lys Pro Met Met Ile Ile Thr Glu Tyr Met Glu Asn  
 195 200 205

Gly Ala Leu Asp Lys Phe Leu Arg Glu Lys Asp Gly Glu Phe Ser Val  
 210 215 220

Leu Gln Leu Val Gly Met Leu Arg Gly Ile Ala Ala Gly Met Lys Tyr  
 225 230 235 240

Leu Ala Asn Met Asn Tyr Val His Arg Asp Leu Ala Ala Arg Asn Ile  
 245 250 255

Leu Val Asn Ser Asn Leu Val Cys Lys Val Ser Asp Phe Gly Leu Ser  
 260 265 270

Arg Val Leu Glu Asp Asp Pro Glu Ala Thr Tyr Thr Thr Ser Gly Gly  
 275 280 285

Lys Ile Pro Ile Arg Trp Thr Ala Pro Glu Ala Ile Ser Tyr Arg Lys  
 290 295 300

Phe Thr Ser Ala Ser Asp Val Trp Ser Phe Gly Ile Val Met Trp Glu  
 305 310 315 320

Val Met Thr Tyr Gly Glu Arg Pro Tyr Trp Glu Leu Ser Asn His Glu  
 325 330 335

Val Met Lys Ala Ile Asn Asp Gly Phe Arg Leu Pro Thr Pro Met Asp  
 340 345 350

Cys Pro Ser Ala Ile Tyr Gln Leu Met Met Gln Cys Trp Gln Gln Glu  
 355 360 365

Arg Ala Arg Arg Pro Lys Phe Ala Asp Ile Val Ser Ile Leu Asp Lys  
 370 375 380

Leu Ile Arg Ala Pro Asp Ser Leu Lys Thr Leu Ala Asp Phe Asp Pro  
 385 390 395 400

Arg Val Ser Ile Arg Leu Pro Ser Thr Ser Gly Ser Glu Gly Val Pro  
 405 410 415

Phe Arg Thr Val Ser Glu Trp Leu Glu Ser Ile Lys Met Gln Gln Tyr  
 420 425 430

Thr Glu His Phe Met Ala Ala Gly Tyr Thr Ala Ile Glu Lys Val Val  
 435 440 445

Gln Met Thr Asn Asp Asp Ile Lys Arg Ile Gly Val Arg Leu Pro Gly  
 450 455 460

His Gln Lys Arg Ile Ala Tyr Ser Leu Leu Gly Leu Lys Asp Gln Val  
 465 470 475 480

Asn Thr Val Gly Ile Pro Ile Glu Gln Lys Leu Ile Ser Glu Glu Asp  
 485 490 495

Leu

<210> 39  
<211> 1737  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Fusion protein construct

<400> 39  
ggtacctcct ttgattagta tattcctatc ttaaagttac ttttatgtgg aggcattaac 60  
at ttgttaat gacgtcaaaa ggatagcaag actagaataa agctataaag caagcatata 120  
atattgcgtt tcatcttttag aagcgaattt cgccaatatt ataattatca aaagagaggg 180  
gtggcaaacg gtatttggca ttattaggtt aaaaaatgta gaaggagagt gaaacccatg 240  
aaaaaatta tgttagtttt tattacatta attttagtta gtttaccatg tgcacaacaa 300  
acagaagcaa aagatgcaag tgcatttaaat aaagaaaata gtattagtat tatggcacca 360  
ccagcaagtc caccagcaag tccaaaaaca ccaattgaaa aaaacatcg agatggatcc 420  
gattataaag acgatgatga taaacacaga cgtagaaaaa atcaacgtgc tgcacaatcc 480  
ccagaagatg tgtatttttc gaaaagtga caattaaaac catataaaac ttatgttgat 540  
ccgcatacgt acgaagacc aaatcaagca gtattaaaaa ttacaacaga aatacaccca 600  
agttgtgtta caagacaaaa agttattgga gcaggtgaat tcggagaggt atataaagg 660  
atgttaaaaa catcatcagg taaaaagaa gttccggtg caattaaaac cttaaaggca 720  
ggatatacag aaaaacagcg agttgatttt ttagtgaag caggaattat gggtaattt 780  
agccatcata atattattcg ttggaagga gtaataagta aatataaacc aatgatgatt 840  
attacagaat acatggaaaa cgggtcctta gataaatttt tacgtgaaaa ggatgggtgaa 900  
tttagtgttt tacaattggg ttgtatgtta agaggaattg ctgcaggtat gaaatattta 960  
gtaaatatga attatgttca cgtgatttg gcagcaagaa atatcctagt caattccaat 1020  
ttagtatgta aagttagtga ttttggttta agcagagtat tagaagacga tccagaggca 1080  
acctatacaa catcggggag taaaattcct attcgttgga cagcaccaga agctatcagt 1140  
taccgtaaat ttacaagtc atcagacgtg tggagttttg ggattgtaat gtgggaagtt 1200  
atgacatatg gagaagagacc atattgggaa ttaagtaatc atgaagtat gaaagcaatt 1260  
aacgatggat ttgatttacc aactccgatg gattgtccat ctgccattta tcaactaatg 1320  
atgcaatgtt ggcaacaaga aagagcacga cgtccaaaat ttgcagatat tgttagtatt 1380  
ttagacaaat taattcgtgc accagatagt ttaaaaaact tagcagactt tgatcctcgt 1440

gttagtattc gattaccaag tacgtcaggt tccgaaggag ttccatttcg cacagtctcc 1500  
 gaatgggttg aatcaattaa aatgcaacaa tacaccgaac accttatggc agcagggttac 1560  
 acagcaatcg aaaaagttgt tcaaatgaca aatgatgata ttaaacgtat tggagttaga 1620  
 ttaccaggcc accagaaacg tattgcatat tctttattag gttaaaga tcaagttaat 1680  
 accgtgggaa ttccaattga aaaaaaatta atttcgaag aagacttata agagctc 1737

<210> 40  
 <211> 497  
 <212> PRT  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Predicted Fusion Protein  
 <400> 40

Met Lys Lys Ile Met Leu Val Phe Ile Thr Leu Ile Leu Val Ser Leu  
 1 5 10 15

Pro Ile Ala Gln Gln Thr Glu Ala Lys Asp Ala Ser Ala Phe Asn Lys  
 20 25 30

Glu Asn Ser Ile Ser Ser Met Ala Pro Pro Ala Ser Pro Pro Ala Ser  
 35 40 45

Pro Lys Thr Pro Ile Glu Lys Lys His Ala Asp Gly Ser Asp Tyr Lys  
 50 55 60

Asp Asp Asp Asp Lys His Arg Arg Arg Lys Asn Gln Arg Ala Arg Gln  
 65 70 75 80

Ser Pro Glu Asp Val Tyr Phe Ser Lys Ser Glu Gln Leu Lys Pro Leu  
 85 90 95

Lys Thr Tyr Val Asp Pro His Thr Tyr Glu Asp Pro Asn Gln Ala Val  
 100 105 110

Leu Lys Phe Thr Thr Glu Ile His Pro Ser Cys Val Thr Arg Gln Lys  
 115 120 125

Val Ile Gly Ala Gly Glu Phe Gly Glu Val Tyr Lys Gly Met Leu Lys  
 130 135 140

Thr Ser Ser Gly Lys Lys Glu Val Pro Val Ala Ile Lys Thr Leu Lys  
 145 150 155 160

Ala Gly Tyr Thr Glu Lys Gln Arg Val Asp Phe Leu Gly Glu Ala Gly  
165 170 175

Ile Met Gly Gln Phe Ser His His Asn Ile Ile Arg Leu Glu Gly Val  
180 185 190

Ile Ser Lys Tyr Lys Pro Met Met Ile Ile Thr Glu Tyr Met Glu Asn  
195 200 205

Gly Ala Leu Asp Lys Phe Leu Arg Glu Lys Asp Gly Glu Phe Ser Val  
210 215 220

Leu Gln Leu Val Gly Met Leu Arg Gly Ile Ala Ala Gly Met Lys Tyr  
225 230 235 240

Leu Ala Asn Met Asn Tyr Val His Arg Asp Leu Ala Ala Arg Asn Ile  
245 250 255

Leu Val Asn Ser Asn Leu Val Cys Lys Val Ser Asp Phe Gly Leu Ser  
260 265 270

Arg Val Leu Glu Asp Asp Pro Glu Ala Thr Tyr Thr Ser Gly Gly  
275 280 285

Lys Ile Pro Ile Arg Trp Thr Ala Pro Glu Ala Ile Ser Tyr Arg Lys  
290 295 300

Phe Thr Ser Ala Ser Asp Val Trp Ser Phe Gly Ile Val Met Trp Glu  
305 310 315 320

Val Met Thr Tyr Gly Glu Arg Pro Tyr Trp Glu Leu Ser Asn His Glu  
325 330 335

Val Met Lys Ala Ile Asn Asp Gly Phe Arg Leu Pro Thr Pro Met Asp  
340 345 350

Cys Pro Ser Ala Ile Tyr Gln Leu Met Met Gln Cys Trp Gln Gln Glu  
355 360 365

Arg Ala Arg Arg Pro Lys Phe Ala Asp Ile Val Ser Ile Leu Asp Lys  
370 375 380

Leu Ile Arg Ala Pro Asp Ser Leu Lys Thr Leu Ala Asp Phe Asp Pro  
385 390 395 400

Arg Val Ser Ile Arg Leu Pro Ser Thr Ser Gly Ser Glu Gly Val Pro  
405 410 415

Phe Arg Thr Val Ser Glu Trp Leu Glu Ser Ile Lys Met Gln Gln Tyr  
420 425 430

Thr Glu His Phe Met Ala Ala Gly Tyr Thr Ala Ile Glu Lys Val Val  
435 440 445

Gln Met Thr Asn Asp Asp Ile Lys Arg Ile Gly Val Arg Leu Pro Gly  
450 455 460

His Gln Lys Arg Ile Ala Tyr Ser Leu Leu Gly Leu Lys Asp Gln Val  
465 470 475 480

Asn Thr Val Gly Ile Pro Ile Glu Gln Lys Leu Ile Ser Glu Glu Asp  
485 490 495

Leu

<210> 41

<211> 1716

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Fusion protein construct

<400> 41

```

ggtaacctcct ttgattagta tattcctatc ttaaagttac ttttatgtgg aggcattaac      60
at ttgtttaat gacgtcaaaa ggatagcaag actagaataa agctataaaag caagcatata      120
atattgcggtt tcattctttag aagcgaattt cgccaatatt ataattatca aaagagagggg      180
gtggcaaacg gtattttggca ttattagggtt aaaaaatgta gaaggagagt gaaacccatg      240
gcatacgaca gtcgttttga tgaatgggta cagaactga aagaggaaaag ctttcaaac      300
aatacgtttg accgccgcaa atttattcaa ggagcgggga agattgcagg actttctctt      360
ggattaacga ttgccagtc ggttggggcc ttggatccg attataaaga tgatgatgat      420
aaacacagac gtgaaaaaaa tcaacgtgct cgacaatccc cagaagatgt gtatttttcg      480
aaaagtgaac aattaaacc attaaaaact tatgttgatc cgcatacgt cgaagaccga      540
aatcaagcag tattaaatt tacaacagaa atacacccaa gtttgtttac aagacaaaaa      600
gttatttgag caggtgaatt cggagaggta tataaaggta tgtaaaaaa atcatcaggt      660
aaaaaagaag ttccggttgc aattaaacc ttaaaggcag gatatacaga aaacagcga      720

```

gttgattttt taggtgaagc aggaattatg ggtcaattta gccatcataa tattattcgt 780  
 ttggaaggag taataagtaa atataaacca atgatgatta ttacagaata catggaaaac 840  
 ggtgccttag ataaattttt acgtgaaaag gatggtgaat ttagtgtttt acaattggtt 900  
 ggtatgttaa gaggaattgc tgcaggtatg aaattattag ctaatatgaa ttatgttcac 960  
 cgtgatttgg cagcaagaaa tatcctagtc aattccaatt tagtatgtaa agttagtgtat 1020  
 ttgggtttaa gcagagtatt agaagacgat ccagaggcaa cctatacaac atcgggaggt 1080  
 aaaattccta ttcgttggaac agcaccagaa gctatcagtt accgtaaat tacaagtgca 1140  
 tcagacgtgt ggagttttgg gattgtaatg tgggaagtta tgacatatgg agaaagacca 1200  
 tattgggaat taagtaatca tgaagttatg aaagcaatta acgatggatt tagattacca 1260  
 actccagtg attgtccatc tgccatttat caactaatga tgcaatgttg gcaacaagaa 1320  
 agagcacgac gtccaaaaatt tgcagatatt gttagtattt tagacaaatt aattcgtgca 1380  
 ccagatagtt taaaaacttt agcagacttt gatcctcgtg ttagtattcg attaccaagt 1440  
 acgtcaggtt ccgaaggagt tccatttcgc acagtctcgc aatggttgga atcaattaaa 1500  
 atgcaacaat acaccgaaca ctttatggca gcaggttaca cagcaatcga aaaagttggt 1560  
 caaatgacaa atgatgatat taaacgtatt ggagttagat taccaggcca ccagaacgt 1620  
 attgcataatt ctttattagg tttaaaagat caagttaata ccgtgggaat tccaattgaa 1680  
 caaaaaattaa ttccgaaga agacttataa gagctc 1716

<210> 42  
 <211> 490  
 <212> PRT  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Predicted fusion protein  
 <400> 42

Met Ala Tyr Asp Ser Arg Phe Asp Glu Trp Val Gln Lys Leu Lys Glu  
 1 5 10 15

Glu Ser Phe Gln Asn Asn Thr Phe Asp Arg Arg Lys Phe Ile Gln Gly  
 20 25 30

Ala Gly Lys Ile Ala Gly Leu Ser Leu Gly Leu Thr Ile Ala Gln Ser  
 35 40 45

Val Gly Ala Phe Gly Ser Asp Tyr Lys Asp Asp Asp Asp Lys His Arg  
 50 55 60



Arg Arg Lys Asn Gln Arg Ala Arg Gln Ser Pro Glu Asp Val Tyr Phe  
 65 70 75 80  
 Ser Lys Ser Glu Gln Leu Lys Pro Leu Lys Thr Tyr Val Asp Pro His  
 85 90 95  
 Thr Tyr Glu Asp Pro Asn Gln Ala Val Leu Lys Phe Thr Thr Glu Ile  
 100 105 110  
 His Pro Ser Cys Val Thr Arg Gln Lys Val Ile Gly Ala Gly Glu Phe  
 115 120 125  
 Gly Glu Val Tyr Lys Gly Met Leu Lys Thr Ser Ser Gly Lys Lys Glu  
 130 135 140  
 Val Pro Val Ala Ile Lys Thr Leu Lys Ala Gly Tyr Thr Glu Lys Gln  
 145 150 155 160  
 Arg Val Asp Phe Leu Gly Glu Ala Gly Ile Met Gly Gln Phe Ser His  
 165 170 175  
 His Asn Ile Ile Arg Leu Glu Gly Val Ile Ser Lys Tyr Lys Pro Met  
 180 185 190  
 Met Ile Ile Thr Glu Tyr Met Glu Asn Gly Ala Leu Asp Lys Phe Leu  
 195 200 205  
 Arg Glu Lys Asp Gly Glu Phe Ser Val Leu Gln Leu Val Gly Met Leu  
 210 215 220  
 Arg Gly Ile Ala Ala Gly Met Lys Tyr Leu Ala Asn Met Asn Tyr Val  
 225 230 235 240  
 His Arg Asp Leu Ala Ala Arg Asn Ile Leu Val Asn Ser Asn Leu Val  
 245 250 255  
 Cys Lys Val Ser Asp Phe Gly Leu Ser Arg Val Leu Glu Asp Asp Pro  
 260 265 270  
 Glu Ala Thr Tyr Thr Thr Ser Gly Gly Lys Ile Pro Ile Arg Trp Thr  
 275 280 285  
 Ala Pro Glu Ala Ile Ser Tyr Arg Lys Phe Thr Ser Ala Ser Asp Val  
 290 295 300  
 Trp Ser Phe Gly Ile Val Met Trp Glu Val Met Thr Tyr Gly Glu Arg

305	310	315	320
Pro Tyr Trp Glu Leu Ser Asn His Glu Val Met Lys Ala Ile Asn Asp			
325		330	335
Gly Phe Arg Leu Pro Thr Pro Met Asp Cys Pro Ser Ala Ile Tyr Gln			
340	345		350
Leu Met Met Gln Cys Trp Gln Gln Glu Arg Ala Arg Arg Pro Lys Phe			
355	360		365
Ala Asp Ile Val Ser Ile Leu Asp Lys Leu Ile Arg Ala Pro Asp Ser			
370	375		380
Leu Lys Thr Leu Ala Asp Phe Asp Pro Arg Val Ser Ile Arg Leu Pro			
385	390	395	400
Ser Thr Ser Gly Ser Glu Gly Val Pro Phe Arg Thr Val Ser Glu Trp			
	405	410	415
Leu Glu Ser Ile Lys Met Gln Gln Tyr Thr Glu His Phe Met Ala Ala			
	420	425	430
Gly Tyr Thr Ala Ile Glu Lys Val Val Gln Met Thr Asn Asp Asp Ile			
	435	440	445
Lys Arg Ile Gly Val Arg Leu Pro Gly His Gln Lys Arg Ile Ala Tyr			
	450	455	460
Ser Leu Leu Gly Leu Lys Asp Gln Val Asn Thr Val Gly Ile Pro Ile			
465	470	475	480
Glu Gln Lys Leu Ile Ser Glu Glu Asp Leu			
	485	490	

<210> 43

<211> 9808

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Fusion Protein Construct

<400> 43

ctttaaactg ggcattttt ctttaaattt atgctgacga cctttgaatt tgcctttttt 60

cttagcaatt tcgattcctt gtgcctgacg ttccttaatt ttttttcggt ctgattctgc 120

ttgatacttg tacaaattcaa tgacaaggct attaatcaaa cgccttaaat tttcatcttc 180

aataccattc attgagggta aatttaagac ttccaggggt gcccccttaa ttggaatttg	240
attcatcaat tctgttaatt ctttattatt tcgtccta atcgatctaatt cagtaacaat	300
aacaatatcc ccttcacgaa tatagttaag catagcttgt aattgtgggc gttcgaccga	360
ttgaccgcct aatttgtctg aaaagaccct agaaacgccc tgtaacgcct gtaattgccg	420
atctaagttc tgttctttgc tactgacacg tgcatatacca attttagcca ttttcaacca	480
acctctaaaa ttctctcggt tgcaataacc aatcagcaat atctactttt tcaatttcaa	540
attgcttata agaaattgtc ttttcgtaag cgataaaatc ttgcgcata tgttgcctcat	600
taaaaatagc caccacttcg tcattttcta aaactcgata aataaatttt ttcattttac	660
tcctctatt atgccaact taaatgacct attcaccaag tcaattatac tgctaaaaac	720
atattaggac aaataggat actctattga cctataaatg atagcaactt aaaagatcaa	780
gtgttcgctt cgtctcact gcccctcgac gttttagtag cctttccctc acttcggtca	840
gtccaagcca actaaaagt ttccgggtac tctctcttc tccccctaat aattaattaa	900
aatcttactc tgtatatctc tgctaatac tcaactaaaca gcaaagaaaa acaaacacgt	960
atcatagata taaatgtaat ggcatagtgc gggttttatt ttcagcctgt atcgtagcta	1020
aacaaatcga gttgtgggtc cgttttgggg cgttctgccca atttgtttag agttctctga	1080
ataaatgtac gttctaaatt aaacgaagct gtcagcgcct ttatatagct ttctcgttct	1140
tcctttttta atttaatgat cgatagcaac aatgatttaa cactagcaag ttgaatgccca	1200
ccatttcttc ctggtttaat cttaaagaaa atttctgat tcgccttcag taccttcagc	1260
aatttatcta atgtccgttc aggaatgctt agcacttctc taatctcttt ttgtgtcgtc	1320
gctaaataag gcttgtatc atcgcttttt tcgctaatat aagccattaa atcttctttc	1380
cattctgaca aatgaacacg ttgacgttcg cttctttttt tctgaattt aaaccacctt	1440
tgacggacaa ataatcttt actgggttaaa tcacttgata cccaagcttt gcaagaatg	1500
gtaatgtatt ccctattagc cccttgatag tttctggaat aggcactttt acaaattttg	1560
attacttctt ttctctctaa gggttgatct aatcgattat taaactcaaa catattatat	1620
tcgcacgttt cgattgaata gcctgaacta aagtaggcta aagagaggtt aaacataacg	1680
ctattgcgc ctactaaacc ctttctcct gaaaatttcg tttcgtgcaa taagagatta	1740
aaccaggggt catctacttg ttttttgctt tctgtaccgc ttaaaaccgt tagacttgaa	1800
cgagtaaagc ccttattatc tgtttgtttg aaagaccaat cttgccattc ttgaaagaa	1860
taacggtaat tgggatcaaa aaattctaca ttgtccgttc ttggtatagc agcaatccca	1920
aaatgattgc acgttagatc aactggcaaa gactttccaa aatattctcg gatattttgc	1980

gagattattt	tggtctgtt	gacagattta	aattctgatt	ttgaagtcac	atagactggc	2040
gtttctaaaa	caaaatatgc	ttgataacct	ttatcagatt	tgataattaa	cgtaggcata	2100
aaacctaaat	caatagctgt	tgtaaaata	tcgcttgctg	aaatagtttc	ttttccgtg	2160
tgaaatcaaa	aatcaataaa	gaaggtattg	atttgtctta	aattgttttc	agaatgtcct	2220
ttagtgtatg	aacgggtttc	gtctgcatac	gtaccataac	gataaacggt	tggtgtccaa	2280
tcgctaaatg	tatcttgatt	ttcgtgaatc	gcttcttcgg	aagtcagaac	aacgccacgt	2340
ccgccaatca	tgcttttttt	tgagcgatac	gcacaaaatg	cccctttact	tttacctggc	2400
ttggtagtga	ttgagcgaat	tttactattt	ttaaatttgt	actttaacaa	gccgtcatga	2460
agcacagttt	ctacaacaaa	agggatattc	attcagctgt	tctcctttct	tacgaaaatt	2520
aattagttag	aagctacgat	caaaagtgaa	tcacaacaaa	aaaggcaatc	aactaagttt	2580
ttcttaattg	attgcctggg	atcttcttaa	agacttgaaa	tcccctcaaa	aacccgatat	2640
aatgggttta	cagatattta	agtatctgat	taataaagta	attaataact	ttaccaaatt	2700
ttgggtctcg	acttctttta	ttgattgggt	gtaatcaatt	aaggctcgca	gttaaaattt	2760
ctcaggcttt	aactggctgt	ggctcttttt	ttgtattctt	tattcagttc	gttgtttcgt	2820
tatatctagt	atatcgcttt	ttaaaaaaat	aagcaatgat	ttcgtgcatt	attcacacga	2880
aatcattgct	tttttcttct	tccatttcta	actccaatgt	tacttgttct	gtttctgggt	2940
ctggttctgt	tggtctcatt	gggattaaat	ccactactag	cggtgagtta	gttccgtctc	3000
taaatgccgg	ttaagtaata	gccggttaag	tggtcaaaat	ttgggaaaaat	ctcaaccgcg	3060
attaagtttt	gatgccatga	caatcggttg	aaatttgaac	aaaactaatg	ctaaaaagct	3120
atctgacttt	atgagtgtag	agccacaaat	acgactttgg	gatatacttc	aaacaaagtt	3180
taaaagctaa	gcacttcaag	aaaaagttta	tatcgaatat	gacaaagtaa	aagcagatac	3240
ttgggataga	cgtaatatgc	gtgttggaatt	taatcccaat	aaactcacac	atgaagaagt	3300
gattttggtta	aaacaaaata	ttatcgacta	catggaagat	gacgggttta	caagattaga	3360
cttagctttt	gattttgaag	atgatttgag	cgattactat	gcaatgactg	ataaagcagt	3420
taagaaaact	gttttttatg	gtcgtaatgg	caagccagaa	acaaaatat	ttggtgtccg	3480
tgatagtgat	agattttatta	gaattttata	taaaaaacaa	gaacgtaaag	ataacgcaga	3540
tgttggaagt	gtgtttgaac	atttatggcg	tgtagaagtt	gaattaaaaa	gagatatggt	3600
tgattactgg	aatgattgtt	ttaatgattt	acacatcttt	gaaacctgcy	tggtgtactt	3660
tagaaaaaat	taatgagcaa	gctatgggtt	atactttgtt	gcatagaaga	agtatgtggg	3720
gaaagctaa	taagaatact	aagactaaat	ttaaaaaatt	gattagagaa	atatctccaa	3780
ttgatttaac	ggaattaatg	aaatcgactt	taaaagcgaa	cgaaaaacaa	ttgcaaaagc	3840

agattgattt ttggcaacgt gaatttaggt tttggaagta aaataagttt tatttgataa	3900
aaattgctaa ttcagtataa ttaatattta cgaggtgaca taacgtatga aaaaatcaga	3960
ggattattcc tcctaaatat aaaaatttaa aatttaggag gaagtatat atgactttta	4020
atattattga attagaaaa tgggtagaaa aagaataatt tgaacactat ttaatcagc	4080
aaactactta tagcattact aaagaaattg atattacttt gtttaaagat atgataaaaa	4140
agaaaggata tgaattttat ccctctttaa tttatgcaat tatggaagtt gtaataaaaa	4200
ataaagtgtt tagaacagga attaatagtg agaataaatt aggttattgg gataagttaa	4260
atcctttgta tacagttttt aataagcaaa ctgaaaaatt tactaacatt tggactgaat	4320
ctgataaaaa cttcattttt ttttataata attataaaaa tgacttgctt gaataataag	4380
ataagaaga aatgtttcct aaaaaaccga tacctgaaaa caccataccg atttcaatga	4440
ttccttggtg tgatttttagt tcatttaatt taaattattg taacaatagc agctttttat	4500
tgccattatt tacgataggt aaattttata gtgagaataa taaaatttat ataccagttg	4560
ctctgcaact tcatcattct gtatgtgatg gttaccatgc ttcactattt atgaatgaat	4620
ttcaagatat aattcatagg gtatagtgatt ggatttagtt tttagatttt gaaagtgaat	4680
ttaattttat acacgtaagt gatcataaaa tttatgaacg tataacaacc acattttttg	4740
gttgcttggtg gttttgattt tgaatttggt tttgaactta tggactgatt tattcagtc	4800
attttttggtg cttgcacaaa aactagcctc gcagagcaca cgcattaatg acttatgaaa	4860
cgtagtaaat aagtctagtg tgttatactt tacttggaag atgcaccgaa taaaaatat	4920
tgaagaacaa ctagcaaaaag attttaaaga gttattttat tttaaagctt tataacatga	4980
gtgaagcgaa tttttaaatt tcgatagaaa tttttacatc aaaaagcccc ctgtcaaat	5040
tgacgaaggg ggttttttgg cgcacgcttt tcgtagaaa tatacaagat tgaataatcg	5100
gtataagtgc gccctttggt ttgaacttag caggttacat caatttttta aaatgatgta	5160
taagtgcgcc cttttaaatt ttgagtgatt atattttttg agttagaaaa agggattggg	5220
aaaatttccc aaaataattt aaaaaataag caaaaatttt cgatagaaa tgtgtctatt	5280
tttgcaaaag gtgtatacct tgactgtgct tgctgttaca ttaagtttat ttttaagtta	5340
ttaaaaaaga aatagctttt aaagtttggc tcgctgtcgc tttataaagc tgattgactt	5400
ttgattgcga actacttaaa gaaaacaac tcggactatt cgttttcttc tctttggttt	5460
gaacatcagc aattatcccc tcttgattgc ctatttttagc ttgttttagaa gaaacaaaag	5520
ctaaaagctc ctcttgggtt ttaaaacgct gtgtggggct tagaacgcc ttaaacgacc	5580
cttggtttac ttttatacta gcttccacct cgaaaaaagg ttctttttta aaattctcta	5640

tggtctcctg	gcgctgaaaa	aataaggat	aagggtggcg	tttgaacacg	tcctagttaa	5700
aatgtacctt	gtacgcccc	tctgttgtaa	atttaacgta	tacaaaggcg	ttgcgttcat	5760
gccgatcaac	caatcggcaa	tttggcgtgt	ttgcgcttct	tgataaaagg	gatagtaatt	5820
cattccaggt	tgcaaaat	gaaaaccgct	tcggattaca	tctttttcta	agctattgat	5880
ccatagtctt	ttaaatgttt	tatcttttga	aaaggcattt	gctttatgga	taatcgacca	5940
ggcgataatt	tcacctttct	tgctgctatc	tggtgcaaca	ataattgtat	ttgccttttt	6000
gagaagttct	gcaacaattt	taaactgctt	tcccttatct	tttgcaactt	caaaatcgta	6060
tcgatcagga	aaaatcgcca	aagattcaag	tttccaattt	tgccactttt	cgtcataatg	6120
acctggttct	gctaattcca	ctaaatgccc	aaaaccaaag	gtgataaacg	tttcatctgt	6180
aaatagtggt	tctttgatct	caaaaataacc	gtcttttttg	gtgcttttgt	ttaaagcact	6240
tgcgtaggct	aatgcctggc	ttggtttttc	agctaaaata	accgtactca	ttactatccc	6300
ctcttttcat	tgttttttct	ttgatcgact	gtcacgttat	atcttgctcg	ataccttcta	6360
aacgttcggc	gattgatccc	agtttgttct	tcaacttctt	tatcgataaa	accattcaaa	6420
aacaaatcga	aagcatggat	gcgccgcgtg	cggtgctggg	agatggcgga	cgcatgggat	6480
atgttctgcc	aagggttggt	ttcgccattc	acagttctcc	gcaagaattg	attggctcca	6540
attcttggag	tggtgaatcc	gttagcgagg	tgccgccggc	ttccattcag	gtcgagggtg	6600
ccggctccca	tgaccgcgca	cgcaacgcgg	ggaggcgagc	aaggtatagg	gcggcgccca	6660
caatccatgc	caaccggttc	catgtgctcg	ccgagggcgc	ataaatcgcc	gtgacgatca	6720
gcggtccagt	gatcgaagtt	aggctggtaa	gagccgcgag	cgatccttga	agctgtccct	6780
gatggtcgtc	atctacctgc	ctggacagca	tggtctgcaa	cgccggccatc	ccgatgccgc	6840
cggaagcgag	aagaatcata	atgggggaagg	ccatccagcc	tcgcgtcgca	atacgactca	6900
ctataggggc	aattgggtac	cgggccccc	ctcgaggctc	acggtatcga	taagcttgat	6960
atcgaaatcc	tgacgcccgg	gggatccact	agttctagag	cgcccgccac	cgcggtggag	7020
ctccagcttt	tgctcccttt	agtggagggt	aatgctagaa	atatttttat	tgattataaa	7080
gatgatcttc	ttgagatcgt	tttgtctctc	gcgtaatctc	ttgctctgaa	aacgaaaaaa	7140
ccgccttgca	ggcggttttt	tcgaagggtc	tctgagctac	caactctttg	aaccgaggtg	7200
actggtctgg	aggagcgcga	tcacaaaacc	ttgtcctttc	agtttagcct	taaccggcgc	7260
atgacttcaa	gactaactcc	tctaaatcaa	ttaccagtgg	ctgctgccag	tggtgctttt	7320
gcatgtcttt	ccgggttgga	ctcaagacga	tagttaccgg	ataaggcgca	gcggctggac	7380
tgaaacgggg	gttcgtgcat	acagtcacgc	ttggagcgaa	ctgcctacc	ggaactgagt	7440
gtcaggcgtg	gaatgagaca	aacgcggcca	taacagcgga	atgacaccgg	taaacggaaa	7500

ggcagggaaca	ggagagcgca	cgagggagcc	gccaggggga	aacgcctggt	atctttatag	7560
tcctgtcggg	tttcgccacc	actgatttga	gcgtcagatt	tcgtgatgct	tgtcaggggg	7620
gcggagccta	tggaaaaacg	gctttgccgc	ggccctctca	cttccctgtt	aagtatcttc	7680
ctggcatctt	ccaggaaate	tcgccccgt	tcgtaagcca	ttccgcctcg	ccgcagtcga	7740
acgaccgagc	gtagcggagt	agtgagcgag	gaagcggaat	atatcctgtg	tcacatatct	7800
tgctgacgca	ccggtgcagc	ctttttttct	ctgccacatg	aagcaattca	ctgacaccct	7860
catcagtgcc	aacatagtaa	gccagtatac	actccgctag	cgctgatgct	cgccggtgct	7920
tttgccgtta	cgcaccaccc	cgtcagtagc	tgaacaggag	ggacagctga	tagaaacaga	7980
agccactgga	gcacctcaaa	aacaccatca	tacactaaat	cagtaagttg	gcagcatcac	8040
ccgacgcact	ttgcgcgcaa	taataacctg	tgacggaaga	tcacttcgca	gaataaataa	8100
atcctggtgt	ccctgttgat	accgggaagc	ectggggcaa	cttttggcga	aaatgagacg	8160
ttgatcgcca	cgtaagaggt	tcacaacttc	accataatga	aataagatca	ctaccgggcg	8220
tattttttga	gttatcgaga	ttttcaggag	ctaaggaagc	taaaatggag	aaaaaaatca	8280
ctggatatac	caccgttgat	atatcccaat	ggcatcgtaa	agaacatttt	gaggcatttc	8340
agtcagttgc	tcaatgtacc	tataaccaga	ccgttcagct	ggataattacg	gccttttttaa	8400
agaccgtaaa	gaaaaataag	cacaagtttt	atccggcctt	tattcacatt	cttgcgccgc	8460
tgatgaatgc	tcatccggaa	ttccgtatgg	caatgaaaga	cggtgagctg	gtgatatggg	8520
atagtgttca	cccttgttac	accgtttttc	atgagcaaac	tgaaacgttt	tcacgcctct	8580
ggagtgaata	ccacgacgat	ttccggcagt	ttctacacat	atattcgcaa	gatgtggcgt	8640
gttacgggtga	aaacctggcc	tatttcacct	aagggtttat	tgagaatatg	tttttcgtct	8700
cagccaatcc	ctgggtgagt	ttcaccagtt	ttgattttaa	cgtggccaat	atggacaact	8760
tcttcgcccc	cgttttcacc	atgggcaaat	attatacgca	aggcgacaag	gtgctgatgc	8820
cgctggcgat	tcaggttcat	catgccgtct	gtgatggctt	ccatgtcggc	agaatgctta	8880
atgaattaca	acagtactgc	gatgagtggc	agggcggggc	gtaatttttt	taaggcagtt	8940
attggtgccc	ttaaacgcct	ggtgctacgc	ctgaataagt	gataataagc	ggatgaatgg	9000
cagaaattcg	aaagcaaat	cgaccgggtc	gtcggttcag	ggcagggctg	ttaaatagcc	9060
gcttatgtct	attgctgggt	taccggttta	ttgactaccg	gaagcagtg	gaccgtgtgc	9120
ttctcaaatg	ctgagggcca	gtttgctcag	gctctccccg	tgagggtaat	aattgacgat	9180
atgatcattt	attctgcctc	ccagagcctg	ataaaaaacg	ttagcgcttc	gttaatatag	9240
atgtaggtgt	tcacaggggt	agccagcagc	atcctgcgat	gcagatccgg	aacataatgg	9300

tgcagggcgc ttgtttcggc gtgggtatgg tggcaggccc cgtggccggg ggactgttgg 9360  
 gcgctgccgg cacctgtctc acgagttgca tgataaagaa gacagtcata agtgccggcga 9420  
 cgatagtcac gccccgcgcc caccggaagg agctaccgga cagcgggtgcg gactgttgta 9480  
 actcagaata agaaatgagg ccgctcatgg cgttgactct cagtcatagt atcgtggtat 9540  
 caccggttgg ttccactctc tgttgcgggc aacttcagca gcacgtaggg gacttccgcg 9600  
 ttccagact ttacgaaaca cggaaaccga agaccattca tgtgttgct caggtcgcag 9660  
 acgttttgca gcagcagtcg ctccacgttc gctcggtat cggtgattca ttctgctaac 9720  
 cagtaaggca accccgccag cctagccggg tcttcaacga caggagcacg atcatgcgca 9780  
 cccgtggcca ggaccaacg ctgcccga 9808

<210> 44  
 <211> 26  
 <212> PRT  
 <213> *Listeria monocytogenes*

<400> 44

Met Lys Lys Ile Met Leu Val Phe Ile Thr Leu Ile Leu Val Ser Leu  
 1 5 10 15

Pro Ile Ala Gln Gln Thr Glu Ala Lys Asp  
 20 25

<210> 45  
 <211> 59  
 <212> PRT  
 <213> *Listeria monocytogenes*

<400> 45

Met Thr Asp Lys Lys Ser Glu Asn Gln Thr Glu Lys Thr Glu Thr Lys  
 1 5 10 15

Glu Asn Lys Gly Met Thr Arg Arg Glu Met Leu Lys Leu Ser Ala Val  
 20 25 30

Ala Gly Thr Gly Ile Ala Val Gly Ala Thr Gly Leu Gly Thr Ile Leu  
 35 40 45

Asn Val Val Asp Gln Val Asp Lys Ala Leu Thr  
 50 55

<210> 46  
 <211> 53  
 <212> PRT  
 <213> *Bacillus subtilis*



<400> 46

Met Ala Tyr Asp Ser Arg Phe Asp Glu Trp Val Gln Lys Leu Lys Glu  
1 5 10 15

Glu Ser Phe Gln Asn Asn Thr Phe Asp Arg Arg Lys Phe Ile Gln Gly  
20 25 30

Ala Gly Lys Ile Ala Gly Leu Ser Leu Gly Leu Thr Ile Ala Gln Ser  
35 40 45

Val Gly Ala Phe Gly  
50

<210> 47

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Primer

<400> 47

gtcaaaacat acgctcttat c

21

<210> 48

<211> 24

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Primer

<400> 48

acataatcag tccaaagtag atgc

24

<210> 49

<211> 29

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Primer

<400> 49

ctctggtacc tcctttgatt agtatattc

29

<210> 50

<211> 29

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Primer

<400> 50  
ctctggatcc atccgcgtgt ttcttttcg 29

<210> 51  
<211> 24  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Epitope insert

<400> 51  
gattataaag atgatgatga taaa 24

<210> 52  
<211> 8  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Epitope

<400> 52  
Asp Tyr Lys Asp Asp Asp Asp Lys  
1 5

<210> 53  
<211> 30  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Epitope insert

<400> 53  
gaacaaaaat taattagtga agaagattta 30

<210> 54  
<211> 10  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Epitope

<400> 54  
Glu Gln Lys Leu Ile Ser Glu Glu Asp Leu  
1 5 10

<210> 55  
<211> 9

<212> PRT  
 <213> Mus sp.  
  
 <400> 55  
  
 Ser Pro Ser Tyr Val Tyr His Gln Phe  
 1 5  
  
 <210> 56  
 <211> 9  
 <212> PRT  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence: Epitope  
  
 <400> 56  
  
 Ser Pro Ser Tyr Ala Tyr His Gln Phe  
 1 5  
  
 <210> 57  
 <211> 29  
 <212> DNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence: Primer  
  
 <400> 57  
 ctctgggtacc tcctttgatt agtatattc 29  
  
 <210> 58  
 <211> 36  
 <212> DNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence: Primer  
  
 <400> 58  
 caatggatcc ctcgagatca taatttactt catccc 36  
  
 <210> 59  
 <211> 32  
 <212> DNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence: Primer  
  
 <400> 59  
 atttctcgag tccatggggg gttctcatca tc 32  
  
 <210> 60  
 <211> 25

<212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Description of Artificial Sequence: Primer  
 <400> 60  
 ggtgctcgcag tgcggccgca agctt 25  
 <210> 61  
 <211> 37  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Description of Artificial Sequence: Primer  
 <400> 61  
 cgattccct agttatgttt accaccaatt tgctgca 37  
 <210> 62  
 <211> 31  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Description of Artificial Sequence: Primer  
 <400> 62  
 gcaaattggt ggtaaacata actaggggaa t 31  
 <210> 63  
 <211> 27  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Description of Artificial Sequence: Epitope insert  
 <400> 63  
 agtccaagtt atgcatatca tcaattt 27  
 <210> 64  
 <211> 33  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Description of Artificial Sequence: Primer  
 <400> 64  
 cgatagtcca agttatgcat atcatcaatt tgc 33  
 <210> 65  
 <211> 34

<212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Description of Artificial Sequence: Primer  
 <400> 65  
 gtcgcaaatt gatgatatgc ataacttgga ctat 34  
 <210> 66  
 <211> 8  
 <212> RNA  
 <213> Artificial Sequence  
 <220>  
 <223> Description of Artificial Sequence: Consensus Sequence  
 <220>  
 <221> misc\_feature  
 <222> (1)..(1)  
 <223> n is a, c, g, or u  
 <400> 66  
 naggaggu 8  
 <210> 67  
 <211> 19  
 <212> DNA  
 <213> Listeria monocytogenes  
 <400> 67  
 aaggagagtg aaacccatg 19  
 <210> 68  
 <211> 240  
 <212> DNA  
 <213> Listeria monocytogenes  
 <400> 68  
 ggtacctcct ttgattagta tattcctatc ttaaagtgac ttttatgttg aggcattaac 60  
 atttgtaac gacgataaag ggacagcagg actagaataa agctataaag caagcatata 120  
 atattgcgtt tcattcttag aagcgaattt cgccaatatt ataattatca aaagagaggg 180  
 gtggcaaacg gtatttgga ttattagggt aaaaaatgta gaaggagagt gaaacccatg 240  
 <210> 69  
 <211> 240  
 <212> DNA  
 <213> Listeria monocytogenes  
 <400> 69  
 ggtacctcct ttgattagta tattcctatc ttaaagttac ttttatgttg aggcattaac 60  
 atttgtaat gacgtcaaaa ggatagcaag actagaataa agctataaag caagcatata 120

atattgcggt tcattcttag aagcgaattt cgccaatatt ataattatca aaagagaggg 180  
 gtggcaaacg gtatttgga ttattagggt aaaaaatgta gaaggagagt gaaacccatg 240

<210> 70  
 <211> 5  
 <212> PRT  
 <213> *Listeria monocytogenes*  
 <400> 70

Thr Glu Ala Lys Asp  
 1 5

<210> 71  
 <211> 5  
 <212> PRT  
 <213> *Listeria monocytogenes*  
 <400> 71

Asp Lys Ala Leu Thr  
 1 5

<210> 72  
 <211> 5  
 <212> PRT  
 <213> *Bacillus subtilis*  
 <400> 72

Val Gly Ala Phe Gly  
 1 5